

For the Period September 19 to 25, 2017

Wet and cool weather delayed harvest for most producers this past week, according to Saskatchewan Agriculture's weekly Crop Report. Seventy-eight per cent of the crop is now in the bin, slightly up from 75 per cent last week. Harvest progress remains ahead of the five-year (2012-2016) average of 74 per cent for this time of year. Fifteen per cent of the crop is swathed or ready to straight-cut. Many producers expect to be back in the field when warmer weather returns and crops can dry sufficiently.

Harvest is most advanced in the southwestern region, where 92 per cent of the crop is now combined. The southeastern region has 88 per cent combined, the west-central region 81 per cent and the east-central region 77 per cent. The northeastern region has 53 per cent combined, while the northwestern region has 48 per cent combined.

Ninety-four per cent of the mustard, 91 per cent of the durum, 88 per cent of the chickpeas, 84 per cent of the barley, 77 per cent of the spring wheat, 68 per cent of the canola and oats, 66 per cent of the canaryseed, 37 per cent of the flax and 18 per cent of the soybeans have now been combined. Twenty-seven per cent of the canola is swathed or ready to straight-cut.

One year ago

Seventy per cent of the crop was in the bin and 22 per cent was swathed or ready to straight cut. Rain kept some producers out of the field for a few days. Crop quality remained low due to disease and issues such as bleaching, staining and sprouting.

Follow the 2017 Crop Report on Twitter @SKAgriculture

Harvest Progress in SK Per cent Combined All Crops	
Sept 25/17	78
5 year avg. (2012-2016)	74
Sept 26/16	70
Sept 28/15	74
Sept 29/14	68
Sept 23/13	71
Sept 24/12	89
10 year avg. (2007-2016)	72

Although the rain was welcomed by many producers in the drier areas of the province, those in the central and northern areas need warm and dry weather soon so that harvest can resume. Rainfall this past week ranged from nil to 36 mm in the Debden and Hafford areas.

Topsoil moisture conditions continue to improve with the recent rainfall, although the subsoil will need significant new moisture to replenish what has been lost to the hot and dry weather. Across the province, topsoil moisture on cropland is rated as 29 per cent

Saskatchewan Harvest September 25, 2017 % combined	
Winter wheat	100
Fall rye*	100
Spring wheat	77
Durum	91
Oats**	68
Barley	84
Canaryseed	66
Flax	37
Canola	68
Mustard	94
Soybeans	18
Lentils	99
Peas	99
Chickpeas	88
*includes 11 per cent 'other'	
**includes three per cent 'other'	

For further information, contact Shannon Friesen, PAg,
Cropping Management Specialist, Moose Jaw, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3592, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.saskatchewan.ca/crop-report.

adequate, 36 per cent short and 35 per cent very short. Hay land and pasture topsoil moisture is rated as 22 per cent adequate, 34 per cent short and 44 per cent very short. The extended period of hot and dry conditions this summer has negatively affected production, particularly in the southern and central regions. Canola crops suffered the most, as much of the crop was in full flowering during the hottest time of the season; there are indications that yield has been reduced by as much as 75 per cent in some areas. Soybean crops have been slow to mature and there are reports of pods not filling completely. Many lentil and cereal crops were unable to fill properly as they ran out of moisture much earlier than normal; lighter bushel weights and smaller seeds have resulted. There are also reports of reduced protein content in cereal crops. While yields have been directly affected by the extremely dry conditions, crop quality has been good to excellent with minimal disease issues.

The majority of crop damage this past week was due to wildlife, strong winds, frost and lack of moisture. Pastures and hay land have suffered greatly from the lack of moisture and will need significant rainfall.

Producers are busy combining, completing fall field work, moving cattle and hauling bales.

Saskatchewan Harvest by Crop District September 25, 2017 % combined					
1A	89	3BS	92	7A	82
1B	88	3BN	96	7B	77
2A	84	4A	98	8A	63
2B	89	4B	97	8B	59
3ASE	94	5A	77	9AE	41
3ASW	81	5B	65	9AW	58
3AN	93	6A	95	9B	39
		6B	83		

Southeastern Saskatchewan:

- Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE – Radville and Lake Alma areas

Harvest has slowed in the region due to scattered rain showers throughout the week. Eighty-eight per cent of the crop is now in the bin, slightly up from 86 per cent last week and well ahead of the five-year (2012-2016) average for this time of year of 77 per cent. Crops such as flax, sunflowers and soybeans will need more time to dry before producers can continue with harvest.

Rain was welcomed in the region last week to help replenish topsoil moisture and allow for fall cereals to germinate and emerge. However, significant rain is still needed to replenish the subsoil for next year. Rainfall last week ranged from trace amounts to 29 mm in the Maryfield area. The Alida area has reported the most precipitation (240 mm) in the region since April 1.

Topsoil moisture conditions continue to significantly improve with the recent rain. Topsoil moisture on cropland is rated as 27 per cent adequate, 38 per cent short and 35 per cent

very short. Hay land and pasture topsoil moisture is rated as 22 per cent adequate, 33 per cent short and 45 per cent very short.

Most crop damage this past week was due to strong winds and lack of moisture. Frost was reported but damage is minimal as remaining crops were advanced enough. Many pastures have been overgrazed and some producers have turned cattle out onto hay and stubble fields. Post-harvest herbicide applications continue on those fields that have active weed growth.

Producers are busy combining, completing fall work, hauling bales and cleaning corrals.

Southwestern Saskatchewan:

- Crop District 3ASW – Coronach, Assiniboia and Ogema areas
- Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B – Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 – Consul, Maple Creek and Leader areas

Small amounts of rain in the region have slowed down harvest progress for some producers. A week or more of warm and dry weather will be needed for remaining crops such as flax and soybean to dry down. Ninety-two per cent of the crop is now in the bin, up slightly from 89 per cent last week and well ahead of the five-year (2012-2016) average of 83 per cent for this time of year.

Rainfall last week ranged from trace amounts to 12 mm in the Moose Jaw area. The Moose Jaw area has reported the most precipitation (196 mm) in the region since April 1. While the rain was welcomed by most producers to help germinate fall cereals and keep the dust down, significant amounts are still needed to replenish the subsoil prior to next spring.

Cropland topsoil moisture conditions have slightly improved since last week. Topsoil moisture on cropland is rated as seven per cent adequate, 23 per cent short and 70 per cent very short. Hay land and pasture topsoil moisture is rated as four per cent adequate, 15 per cent short and 81 per cent very short. All crop districts, with the exception of 4B, are reporting that at least 97 per cent of the cropland and 98 per cent of the hay land and pasture remain short to very short topsoil moisture at this time. CD 4B has indicated that 100 per cent of cropland, hay land and pasture remains very short topsoil moisture.

Most crop damage this past week is due to strong winds and lack of moisture. A hard frost was reported in some areas but damage is expected to be minimal as remaining crops were advanced enough. Combine and grass fires remain a concern in the region due to the extremely dry conditions. Many pastures have been overgrazed and have not grown in a number of months; significant rainfall will be needed to replenish what has been lost to the hot and dry weather over the summer.

Producers are busy combining, completing fall work, hauling bales and moving cattle.

East-Central Saskatchewan:

- Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A – Lumsden, Craik, Watrous and Clavet areas

Despite a rainy week, harvest continued in parts of the region. Seventy-seven per cent of the crop is now combined, up from 72 per cent last week and well ahead of the five-year (2012-2016) average of 66 per cent for this time of year. Sixteen per cent of the crop is swathed or ready to straight-cut. Most producers hope to be back in the field soon, once weather conditions improve and crops have sufficiently dried down. There are reports of producers taking the crop off tough and placing it into aeration bins.

The region received rainfall last week ranging from 8 mm in the Saltcoats, Wynyard and Raymore areas to 32 mm in the Langenburg area. The Pelly area has reported the most precipitation (307 mm) in the region since April 1.

Topsoil moisture conditions have greatly improved with the moisture. However, significant amounts are still needed to replenish what was lost over the hot and dry summer. Topsoil moisture on cropland is rated as one per cent surplus, 38 per cent adequate, 36 per cent short and 25 per cent very short. Hay land and pasture topsoil moisture is rated as 20 per cent adequate, 46 per cent short and 34 per cent very short. Crop District 6A is reporting that 76 per cent of the cropland and 83 per cent of the hay land and pasture remain short to very short topsoil moisture at this time.

Crop damage this past week is attributed to lack of moisture and wind. There were a few nights with frost reported but damage so far has been minimal as crops were advanced enough. Some producers have been applying post-harvest herbicides, although weed growth has been slow.

Producers are busy combining, fixing fences, completing fall work and moving cattle.

West-Central Saskatchewan:

- Crop District 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A – Rosetown, Kindersley, Eston, Major
- Crop District 7B – Kerrobert, Macklin, Wilkie and Biggar areas

Another rainy week has delayed harvest for some producers in the region. Eighty-one per cent of the crop is now combined, up from 74 per cent last week and well ahead of the five-year (2012-2016) average of 76 per cent for this time of year. Thirteen per cent of the crop is swathed or ready to straight-cut. Some producers are taking crops off tough and placing them in aeration bins.

Although the moisture halted harvest in some areas, it will help replenish the topsoil and allow fall cereals to germinate. Rainfall last week ranged from trace amounts to 34 mm in the Rosthern area. The Cando area has reported receiving the most precipitation (288 mm) in the region since April 1.

Despite the moisture, topsoil moisture conditions have not improved. Topsoil moisture on cropland is rated as 12 per cent adequate, 44 per cent short and 44 per cent very short.

Hay land and pasture topsoil moisture is rated as 14 per cent adequate, 38 per cent short and 48 per cent very short. Crop District 7A is now reporting that 100 per cent of cropland, hay land and pasture remains short to very short topsoil moisture at this time.

The majority of crop damage this past week was due to strong winds, lack of moisture and frost. Several nights of light frost were reported in the region; however, damage is expected to be minimal as crops were advanced. Birds and deer have damaged some swathed crops.

Combine and grass fires remain a concern in the region due to the dry conditions. Weed growth has been limited, although some producers have been applying post-harvest herbicides.

Producers are busy combining, completing fall work and moving cattle.

Northeastern Saskatchewan:

- Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas

Harvest ground to a halt this past week, thanks to the recent rainfall. Very little combining was done and harvest progress remains at 53 per cent. The five-year (2012-2016) average for this time of year is 69 per cent. Thirty-four per cent of the crop is swathed or ready to straight-cut. Several weeks of warm and dry weather will be needed for the rest of the crop to dry down sufficiently for combining. Some crop is coming off tough and being placed in aeration bins.

Rainfall this past week ranged from 7 mm in the Melfort area to 35 mm in the Nipawin area. The Nipawin area has reported the most precipitation (554 mm) in the region since April 1. Topsoil moisture conditions have greatly improved with the recent moisture. Across the region, topsoil moisture conditions on cropland are rated as 50 per cent adequate, 45 per cent short and five per cent very short. Hay land and pasture topsoil moisture is rated as 36 per cent adequate, 56 per cent short and eight per cent very short. A hard frost hit last week, although damage is expected to be minimal as most crops were advanced enough. Strong winds have also blown some swaths around. Birds and deer have been damaging swathed crops.

Producers are busy combining, working fields and moving cattle.

Northwestern Saskatchewan:

- Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

Very little harvest progress was made last week due to scattered showers and cooler weather. Forty-eight per cent of the crop is now in the bin, up slightly from 46 per cent last week but well behind the five-year (2012-2016) average of 71 per cent for this time of year. Several weeks of warm and dry weather will be needed for remaining crops to dry

down sufficiently for combining. Some crops are being taken off tough and placed in aeration bins.

Most of the region received rainfall last week, ranging from small amounts to 36 mm in the Debden and Hafford areas. The Pierceland area has reported the most precipitation (563 mm) in both the region and the province since April 1. Topsoil moisture conditions continue to improve, thanks to the rain. Cropland topsoil moisture conditions are rated as 63 per cent adequate, 34 per cent short and three per cent very short. Hay land and pasture topsoil moisture is rated as 56 per cent adequate, 41 per cent short and three per cent very short.

Strong winds and frost are the major causes of crop damage this past week. While frost was reported over several nights, damage has been minimal as most crops were advanced enough. There have been a few reports of sprouting in some cereal crops and high green counts in canola. Wildlife such as geese and deer have also damaged some swathed crops. Post-harvest herbicide applications continue in those areas with active weed growth.

Producers are busy combining, completing fall work, hauling bales and fixing fences.


Saskatchewan Harvest Progress - September 25, 2017

*Other - crop that will not be harvested due to weather, insect or disease damage or will be greenfeed

Winter Wheat	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	0	0	0	100	
northwest	0	0	0	100	
provincial	0	0	0	100	
Fall Rye	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	0	0	0	97	3
southwest	0	0	0	87	13
east central	0	0	0	100	0
west central	0	0	0	100	0
northeast	0	0	0	81	19
northwest	0	0	0	100	0
provincial	0	0	0	89	11
Spring Wheat	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	1	1	98	
southwest	4	0	4	92	
east central	4	6	5	85	
west central	5	3	18	74	
northeast	14	6	15	65	
northwest	18	9	25	48	
provincial	7	4	12	77	
Durum	% Standing	% in swath	% ready to straight combine	% combined	
southeast	1	0	1	98	
southwest	3	0	3	94	
east central	0	1	4	95	
west central	19	0	8	73	
northeast	0	0	5	95	
northwest	N/A	N/A	N/A	N/A	
provincial	5	0	4	91	
Barley	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	1	99	
southwest	4	0	3	93	
east central	3	4	4	89	
west central	3	6	8	83	
northeast	6	9	7	78	
northwest	18	13	9	60	
provincial	5	6	5	84	
Oats	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	0	1	1	96	2
southwest	5	0	7	77	11
east central	10	13	5	70	2
west central	7	13	7	54	19
northeast	18	2	21	58	1
northwest	25	17	14	42	2
provincial	13	8	11	65	3
Canaryseed	% Standing	% in swath	% ready to straight combine	% combined	
southeast	6	2	3	89	
southwest	11	0	18	71	
east central	7	0	27	66	
west central	11	2	10	77	
northeast	48	9	6	37	
northwest	N/A	N/A	N/A	N/A	
provincial	16	2	16	66	


Flax	% Standing	% in swath	% ready to straight combine	% combined	
southeast	36	3	22	39	
southwest	32	3	21	44	
east central	31	7	13	49	
west central	52	6	13	29	
northeast	56	18	17	9	
northwest	79	0	10	11	
provincial	39	5	19	37	
Canola	% Standing	% in swath	% ready to straight combine	% combined	
southeast	3	3	3	91	
southwest	5	2	6	87	
east central	4	23	4	69	
west central	2	13	6	79	
northeast	7	41	8	44	
northwest	8	39	13	40	
provincial	5	21	6	68	
Mustard	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	1	0	3	96	
east central	0	0	0	100	
west central	5	3	13	79	
northeast	N/A	N/A	N/A	N/A	
northwest	N/A	N/A	N/A	N/A	
provincial	1	1	4	94	
Soybeans	% Standing	% in swath	% ready to straight combine	% combined	
southeast	64	0	17	19	
southwest	64	0	20	16	
east central	72	0	18	10	
west central	51	0	20	29	
northeast	62	1	33	4	
northwest	100	0	0	0	
provincial	65	0	17	18	
Field Peas	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	0	100	
east central	0	0	0	100	
west central	0	0	0	100	
northeast	0	0	2	98	
northwest	0	0	1	99	
provincial	0	0	1	99	
Lentils	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	0	0	1	99	
east central	0	0	0	100	
west central	0	0	1	99	
northeast	0	0	0	100	
northwest	0	0	0	100	
provincial	0	0	1	99	
Chickpeas	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	0	100	
southwest	9	0	4	87	
east central	0	0	0	100	
west central	N/A	N/A	N/A	N/A	
northeast	N/A	N/A	N/A	N/A	
northwest	N/A	N/A	N/A	N/A	
provincial	8	0	4	88	

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

from September 19 to September 25, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period September 19 to 25, 2017


Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	12	216	4A	49	White Valley	N/A	93	7A	287	St. Andrews	NIL	167
	3	Enniskillen	10	215		51	Reno	NIL	89.5		288	Pleasant Valley	NIL	153
	31	Storthoaks	N/A	10		79	Arlington	NIL	91		290 A	Kindersley	1	155.1
	32	Reciprocity	16	240		109	Carmichael	NIL	95		290 B	Kindersley	N/A	87.4
	33	Moose Creek	N/A	238		110	Piapot	N/A	79		290 C	Kindersley	N/A	24
	34	Browning	19	204		111	Maple Creek	N/A	95		292	Milton	N/A	196
	61	Antler	N/A	108		139 A	Gull Lake	NIL	103		317 A	Marriott	N/A	N/A
	64	Brock	15	191		139 B	Gull Lake	NIL	69.5		317 B	Marriott	3	187
	65	Tecumseh	17	220		169	Pittville	N/A	119		318	Mountain View	N/A	231
	91	Maryfield	29	231		231	Happyland	1	165		320 A	Oakdale	1	195
1B	94	Hazelwood	22.8	209.2	5A	183	Fertile Belt	20	181	7B	320 B	Oakdale	2	202
	122	Martin	27	208		211 A	Churchbridge	32	264		321	Prairiedale	NIL	200
	123	Silverwood	20	190		211 B	Churchbridge	N/A	44		347	Biggar	2	176
	124	Kingsley	17	208		213	Saltcoats	8	183		350 A	Mariposa	NIL	263.1
	125 A	Chester	13	156		241	Calder	26	167		350 B	Mariposa	2	189
	125 B	Chester	12	217		243	Wallace	14	153		351	Progress	5	231
	151	Rocanville	21	221		244	Orkney	N/A	165		352	Heart's Hill	NIL	182
	154 A	Elcapo	N/A	143		245 A	Garry	N/A	157		377	Glenside	1	216
	154 B	Elcapo	16	123		245 B	Garry	N/A	122		378	Rosemount	5	288
	155	Wolseley	3.5	167.5		245 C	Garry	N/A	117		379	Reford	3	184
2A	67	Weyburn	5	120	5B	246 A	Ituna Bon Accord	16	142	8A	381	Grass Lake	N/A	165
	68	Brokenshell	14	107		246 B	Ituna Bon Accord	17	179.5		382	Eye Hill	5	239.9
	96	Fillmore	N/A	19		247	Kellross	N/A	115		409 A	Buffalo	N/A	213
	97	Wellington	3	112		248	Touchwood	N/A	106		409 B	Buffalo	NIL	242
	127 A	Francis	10	166		271	Cote	14	220		410	Round Valley	NIL	281.4
	127 B	Francis	12.5	66.3		273	Sliding Hills	12	217		395	Porcupine	N/A	264
	129	Bratt's Lake	10.5	108		277	Emerald	15	240		397	Barrier Valley	16.8	269.8
	131 A	Baildon	7	134		305	Invermay	21	247		426	Bjorkdale	N/A	45
	131 B	Baildon	NIL	168		307	Elfros	11	204		428	Star City	12	162
	156 A	Indian Head	11	152.9		308 A	Big Quill	8	126		456	Arborefield	16	288
2B	156 B	Indian Head	6	191	6A	308 B	Big Quill	8	132	8B	457	Connaught	20	206
	159	Sherwood	14.1	111.1		331	Livingston	28	307		486	Moose Range	23	291
	160 A	Pense	12	88		335	Hazel Dell	N/A	113		487	Nipawin	35	553.5
	160 B	Pense	15	88.2		336	Sasman	12	211		369	St. Peter	N/A	143
	161	Moose Jaw	7	118		337	Lakeview	N/A	215		370 A	Humboldt	14	170
	162	Caron	9	136		338	Lakeside	12	196		370 B	Humboldt	20	157
	191	Marquis	11	91		366	Kelvington	13	305		371	Bayne	24	198
	38 A	Laurier	5	105.5		367	Ponass Lake	18	267.5		372	Grant	35	221.2
	38 B	Laurier	7.5	109		368	Spalding	16	276		400	Three Lakes	16	205
	39	The Gap	5	85	6B	190 A	Dufferin	19	159	9AE	429 A	Flett's Springs	20	162
3ASE	10	Happy Valley	2	100		190 B	Dufferin	16	152		429 B	Flett's Springs	7	202
	12	Poplar Valley	NIL	93		190 C	Dufferin	15	125		459	Kinistino	15	132
	42	Willow Bunch	4	97		190 D	Dufferin	14	86		460	Birch Hills	11.5	145.7
	43	Old Post	5	102		219 A	Longlaketon	14	95		488	Torch River	35	238
	73 A	Stonehenge	10	87.4		219 B	Longlaketon	14	145		491	Buckland	N/A	141
	73 B	Stonehenge	9	153.5		220	McKillop	20	130		520	Paddockwood	15.5	191.5
	74	Wood River	10	18.3		221 A	Sarnia	20.1	159.9		521	Lakeland	15.5	191.5
	102	Lake Johnston	8.6	105.7		221 B	Sarnia	22.9	127.1	9AW	406	Mayfield	9	212
	103	Sutton	N/A	60		222	Craik	22	116		435	Redberry	36	277
	132 A	Hillsborough	10	101		251	Big Arm	17.5	120.5		436	Douglas	22	207
3ASW	132 B	Hillsborough	12	196		252	Arm River	N/A	115		463	Duck Lake	38	258.6
	193	Eyebrow	10	103		279	Mount Hope	22.2	151.1		466	Meeting Lake	18	317
	17	Val Marie	N/A	29		282	McCraney	30	149		467 A	Round Hill	N/A	273
	18	Lone Tree	2	66		312	Morris	27	158.5		467 B	Round Hill	18	297
	75	Pinto Creek	NIL	110		313	Lost River	N/A	173		467 C	Round Hill	25	397
	76	Auvergne	NIL	83		339	Leroy	9.4	180.4		493	Shellbrook	25	68.8
	77	Wise Creek	N/A	93		340	Wolverine	25	241		494	Canwood	36	269
	78	Grassy Creek	N/A	92.5		341	Viscount	N/A	123		497	Medstead	N/A	90.5
	105	Glenbain	6	106		343	Blucher	25	183	9B	438	Battle River	NIL	233
	106	Whiska Creek	NIL	131		223 A	Huron	N/A	104		440	Hillsdale	1	316.5
3AN	107	Lac Pelletier	NIL	79.5	6B	223 B	Huron	15	97		442	Manitou Lake	4.3	334.5
	108	Bone Creek	NIL	77		284 A	Rudy	12	179		498 A	Parkdale	N/A	354
	138 A	Webb	NIL	162		284 B	Rudy	5	196.5		498 B	Parkdale	NIL	288.5
	138 B	Webb	NIL	90		284 C	Rudy	8	160		499	Mervin	4	430.1
	165	Morse	N/A	161		285	Fertile Valley	2	139		501 A	Frenchman Butte	N/A	413
	166	Excelsior	NIL	106		286	Milden	NIL	187		501 B	Frenchman Butte	2	336
	168 A	Riverside	NIL	117		314	Dundurn	24	153		501 C	Frenchman Butte	3	404
	168 B	Riverside	N/A	122.5		344	Corman Park	16	219		502	Britannia	NIL	274.5
	226	Victory	N/A	51		346	Perdue	3	148		561	Loon Lake	N/A	423
	228 A	Lacadena	N/A	148		376	Eagle Creek	6	208		588 A	Meadow Lake	32	434
3BS	228 B	Lacadena	NIL	7	6B	403	Rosthern	34	244		588 B	Meadow Lake	33	479
	257	Monet	NIL	131							622	Beaver River	15	563.4

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Municipality No: A, B, C and D - more than one reporter


Cumulative Rainfall

from April 1 to September 25, 2017



Cropland Topsoil Moisture Conditions


September 25, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Hay and Pasture Topsoil Moisture Conditions


September 25, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Minimum Temperature


from September 19 to September 25, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Maximum Temperature

from September 19 to September 25, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.