

Vaccine Preventable Disease Monitoring Report Pertussis, 2017 and 2018

Release Date: June 2019

PREPARED BY POPULATION HEALTH BRANCH, SASKATCHEWAN MINISTRY OF HEALTH

FOR MORE INFORMATION CONTACT:

Val Mann, PhD
Chief Population Health Epidemiologist
Population Health Branch, Saskatchewan Ministry of Health
cdc@health.gov.sk.ca

<p><u>Purpose:</u></p> <p>The Saskatchewan Ministry of Health's Population Health Branch provides routine surveillance of notifiable diseases at the provincial and former regional health authorities (RHAs), Athabasca Health Authority (AHA), First Nations and Inuit Health Branch - Saskatchewan (FNIHB-SK) and Northern Inter-Tribal Health Authority (NITHA) levels.</p> <p>This report presents the most recent data for reportable communicable diseases as collected by the Integrated Public Health Information System (iPHIS) and immunization coverage information as collected by Panorama. Limitations associated with these systems have been described elsewhere.</p> <p>Under <i>The Public Health Act, 1994</i> and the accompanying Disease Control Regulations, local medical health officers (MHOs) must report Categories I and II Communicable Diseases, as well as any communicable disease outbreaks to the Chief and Deputy Chief Medical Health Officers. Pertussis is a Category I disease.</p> <p><u>Report Features:</u></p> <p>Background Epidemiological Summary Surveillance Case Definition Case Counts by Year Case Characteristics Vaccine Coverage by AHA and former RHA</p> <p><u>Data Source:</u></p> <p>Panorama (as of April 1, 2019)</p>	<div> <div> <h2>Background</h2> <p>Pertussis (whooping cough) is a highly contagious bacterial disease. It often begins with mild respiratory symptoms, cough and sometimes fever, and can progress to severe coughing attacks characterized by a "whoop sound" when a breath is taken.</p> <p>Infants are most vulnerable and often infected by older siblings, parents or caregivers who may not know they have the disease. One to three deaths related to pertussis occur each year in Canada, particularly in infants too young to be immunized, or in unimmunized, or partially immunized children. Deaths were not required to be reported prior to 2014, although it was common practice to do so.</p> <p>Symptoms usually develop five to 10 days after exposure, but can take up to 21 days.</p> </div> <div> <p>Pertussis is caused by the bacteria <i>Bordetella pertussis</i>. The bacteria are easily spread by droplets from the nose or mouth or through direct contact with the respiratory secretions of an infected person.</p> <p>Pertussis becomes more active on a cyclical basis with increased cases being reported every two to five years. The variability in the number of pertussis cases from one year to the next and in different geographical regions is often because of outbreaks. One hundred forty-three (144) cases of lab-confirmed pertussis were reported in 2018. Many cases were associated with an extended outbreak in one former health region.</p> </div> </div> <div> <h2>Immunization</h2> <p>Acellular pertussis vaccine is only available in combination vaccines. The Saskatchewan Routine Childhood Immunization Schedule recommends a four dose primary series of pertussis vaccine at two, four, six and 18 months of age, one booster at four to six years of age and a second booster in Grade 8. Adults are eligible to receive one lifetime pertussis vaccine dose. Pregnant women are eligible for pertussis vaccine in every pregnancy (implemented October 2017).</p> <p>Transmission is less likely in or to people who are vaccinated. Adolescents and adults who have not received a booster are at risk of infection and are often the source of infection for infants. Infants too young for vaccination are at the greatest risk for serious pertussis complications and deaths.</p> <p>The efficacy of acellular pertussis vaccine following the primary series is estimated to be about 85%, and approximately 90% following booster immunization.</p> </div> <div> <h2>Surveillance</h2> <p>Under <i>The Public Health Act, 1994</i>, Saskatchewan health care providers are required to report cases of pertussis to the local medical health officer (MHO) who then reports the case to the Chief and Deputy Chief Medical Health Officers using the case definition in the Saskatchewan Communicable Disease Manual.</p> <p>Notifiable diseases may be undetected, therefore underreported, due to a number of factors including lack of contact with the health care system or inability of laboratory tests to identify the organism. Some communicable diseases occur rarely and therefore, rates are based on small numbers of cases which can fluctuate dramatically over time. In these situations, year to year comparisons should be interpreted with caution.</p> <p>Surveillance case definitions ensure uniform reporting to allow comparability of surveillance data. The definitions are not intended to be used for clinical or laboratory diagnosis or management of cases.</p> <p>Currently molecular epidemiology genotyping is not available for pertussis.</p> </div>
--	---

EPIDEMIOLOGY AND VACCINE COVERAGE SUMMARY

Pertussis in Saskatchewan: 2018

- One hundred forty-four (144) cases of lab-confirmed pertussis were reported in 2018. Ages ranged from five weeks old to 70 years of age.
- One protracted community outbreak in the former Five Hills Health Region (HR), declared at the end of December 2016, was finally declared over in late January, 2019. The former Five Hills HR outbreak spilled into the former Sun Country HR creating several household clusters. A large community outbreak occurred in the former Saskatoon HR among a largely under-immunized school population.
- Eight of the 13 infant cases in 2018 were too young to be adequately protected by a complete immunization series; two of these were too young to begin their primary series; another three had not started. Two of the four cases who had time to complete the series declined to do so; the others had one or two doses of vaccine. One case was fully immunized.
- One of the three cases hospitalized in 2018 was an infant.
- There were no reported deaths from pertussis.

Pertussis in Saskatchewan: 2014 to 2018

- Over one thousand (1,026) cases of pertussis ranging in age from ten days to 93 years were reported. The median age of cases was 12 years.
- Over one-quarter of the cases (30%) lived in the former Saskatoon HR compared to 12% in the former Regina Qu'Appelle HR and 15% in the former Five Hills HR. Thirteen percent (13%) of all cases lived in First Nation communities.
- Seventy-one cases (7%) were reported hospitalized for pertussis illnesses, including 35 of the 98 infants.
- One-quarter of 98 infant cases (23 cases) were less than two months old and too young to begin their primary immunization series for pertussis. Half (20) of the 41 cases old enough to receive one or two doses of vaccine had not been started. Eleven (11%) infants were up to date, that is, received one or two doses, but were not adequately protected. Seventeen of 29 infant cases, old enough to complete their series had not started (37%). Eight who had completed their primary series still became infected (11%).

Pertussis Coverage in Saskatchewan: 2014 to 2018

- From 2014 to 2018, provincial coverage rates steadily improved or remained stable up to and including five years of age.
- For children older than five years, the rates when based strictly on the number of doses received, declined between 2014 and 2016.
- However, during 2017 and 2018 when rates were based on up-to-date status (i.e., adjusting for delayed or interrupted series), coverage improved or remained stable for children older than seven years.

Table 1: Pertussis case counts by year

	2018*	2017	2016	2015	2014	Total
Saskatchewan	144	425	197	213	47	1,026
Canada	N/A	N/A	3,936	3,514	1,526	8,976

*2018 figure is preliminary
N/A = not available

Table 2: Pertussis case characteristics, Saskatchewan 2014 - 2018

	Cases	Percent of Cases
Total	1026	100
Sex	Male	498
	Female	528
Age	Less than 1 year	10
	1 - 4 years	14
	5 - 19 years	21
	20 - 49 years	50
	50 years and over	6
Hospitalized	Yes	7
	No	93
	Unknown	0
Immunization status of infants (less than 1 year) for pertussis vaccine** (n=98)	3 doses	9
	2 doses	6
	1 dose	24
	0 dose	38
	Too young	24
	Unknown	0
Genotype	Not applicable	1026

**Immunization status is monitored for infant cases only, as infants are the most vulnerable for severe outcomes of pertussis infections.

Table 3: Pertussis vaccine coverage for Saskatchewan by age, dose and year, 2014-2018

Age	Doses	2018	2017	2016 ^a	2015 ^a	2014 ^b
3 months	1	87%	86.3%	85%	85%	84.1%
5 months	2	78.2%	78%	77%	76%	73.7%
8 months	3	80.4%	79.6%	78.8%	77.3%	76.4%
12 months	3	87.2%	86.6%	85.6%	84.9%	84.7%
20 months	3	90.7%	90%	89.1%	88.5%	88.8%
24 months	4	68.2%	63.9%	60.9%	60.5%	60.1%
5 years	3	91.6%	90.6%	89.9%	89.3%	89.8%
7 years	4	80.8%	77.8%	76.8%	74.6%	75.7%
13 years	5	87.2%	85.6%	87.7%	85.1%	87.8%
15 years	5	N/C	N/C	76.2%	74.8%	77.8%
17 years	up-to-date	81.7%	82.5%	N/C	N/C	N/C
13 years	5	N/C	N/C	76.9%	77.5%	80.2%
15 years	up-to-date	78.8%	78.7%	N/C	N/C	N/C
17 years	6	N/C	N/C	67.3%	66.9%	72.2%
13 years	up-to-date	81.9%	81.6%	N/C	N/C	N/C
15 years	6	N/C	N/C	70.6%	71.1%	74.3%
17 years	up-to-date	83.9%	83.4%	N/C	N/C	N/C

^a Vaccine Preventable Disease Monitoring Report: Pertussis, 2015 and 2016 (Data source: Panorama January 12, 2017)

^b Vaccine Preventable Disease Monitoring Report: Pertussis, 2014 (Data source: SIMS January 16, 2015)

N/C = Not calculated because an up-to-date coverage rate provides a better estimate of the percentage of children protected from disease than an age-dose coverage rate, which underestimates the level of protection if children have experienced a delay or interruption in the primary series.

EPIDEMIOLOGY AND VACCINE COVERAGE SUMMARY

Table 4: Pertussis vaccine coverage by Athabasca Health Authority and former health region, 2018

Jurisdiction (with former health region by Peer Group)	Immunization coverage (% immunized), by age and dose												
	3 months	5 months	8 months	12 months	20 months		24 months		5 years	7 years	13 years	15 years	17 years
	1 dose	2 doses	3 doses	3 doses	3 doses	4 doses	3 doses	4 doses	4 doses	up-to-date	up-to-date	up-to-date	up-to-date
Saskatchewan	87	78.2	80.4	87.2	90.7	68.2	91.6	80.8	87.2	81.7	78.8	81.9	83.9
Saskatchewan Health Authority	87	78.3	80.4	87.2	90.7	68.2	91.6	80.9	87.2	81.7	78.8	81.9	83.9
Peer Group A													
Regina Qu'Appelle	91.8	79.9	82.7	88	90.5	70.6	91.4	79.2	86	82.8	76.9	82.2	84.4
Saskatoon	83.2	77.3	78.9	86.9	91.2	64.3	91.5	81.6	85	76.8	75.6	81	85.2
Peer Group D													
Cypress	84.1	77.2	84.4	91.5	92.6	74.4	93.8	84.2	92.1	91.9	84.8	88.2	87.4
Five Hills	89.7	81	83	90	92.7	76.6	93.1	83.5	90.1	84.2	84.4	84.8	85.4
Heartland	89.5	83.7	82.1	90.7	93	77.2	93	88	92.7	90	86.1	88.8	90.8
Kelsey Trail	93.6	82	88.5	92.2	96.2	83.6	95.8	90.9	95.1	87.2	83.3	86.7	86.3
Sun Country	94.4	91.8	93.4	95.4	94.8	87.4	95.8	92.8	94.7	93.4	87.4	85.8	90.8
Sunrise	85.8	79.7	80.9	85.7	90.1	69	92.4	83.3	88.6	84.7	83.1	84.1	89
Peer Group F													
Athabasca Health Authority	85.7	72.2	82.1	100	93.5	71	100	69.4	100	86.7	81.7	85.4	80.4
Keewatin Yatthé	77.6	64.1	72.1	79.7	89.6	45.4	89.2	63.7	83	78.4	68	62.1	51
Mamawetan Churchill River	81.4	65.5	73.5	89.3	94.3	71.8	95.7	82.2	98	80	83.2	66.8	64.3
Peer Group H													
Prairie North	86.1	74.3	76.6	83.2	86.2	62.3	87.6	75.3	87.4	80.6	74.9	72.4	73.8
Prince Albert Parkland	83	68.3	66.7	78.2	85.6	53.8	88	70	86.1	79.3	82.3	84.1	79.8

Table 5: Pertussis vaccine coverage by Athabasca Health Authority and former health region, 2017

Jurisdiction (with former health region by Peer Group)	Immunization coverage (% immunized), by age and dose												
	3 months	5 months	8 months	12 months	20 months		24 months		5 years	7 years	13 years	15 years	17 years
	1 dose	2 doses	3 doses	3 doses	3 doses	4 doses	3 doses	4 doses	4 doses	up-to-date	up-to-date	up-to-date	up-to-date
Saskatchewan	86.3	78	79.6	86.6	90	63.9	90.6	77.8	85.6	82.5	78.7	81.6	83.4
Saskatchewan Health Authority	86.3	78	79.6	86.6	90	64	90.5	77.8	85.6	82.5	78.7	81.6	83.3
Peer Group A													
Regina Qu'Appelle	87.4	79.1	80.1	86.9	89.7	61.7	89.4	75	85.5	83.4	76.4	82.1	83.4
Saskatoon	85.1	77.5	79.8	86.6	89.2	61.7	90.5	78.8	82.8	79.1	75.3	83	84.5
Peer Group D													
Cypress	88.4	79.1	80.2	88.3	93.3	72.4	92.4	82.8	91.1	89.1	86.1	84.9	87.9
Five Hills	88.8	83.2	82.7	86.8	92	71.6	93.3	81.1	86.3	85.3	83.4	84	84.6
Heartland	85	79.4	87.6	90.3	93.6	73.1	93.3	85.7	92.9	92.4	85.5	85.6	90.5
Kelsey Trail	89.4	84.5	88.5	93.5	95.3	77	96.7	86.2	90.9	89	85.3	84.4	86.6
Sun Country	94.3	90.7	90.8	94.4	95	84.3	94.9	89.7	93.5	92.7	85.4	88.2	88.9
Sunrise	84.5	79.7	83.2	88.8	91.9	71.5	91.7	81.1	86.8	81.5	84.3	84.3	83.9
Peer Group F													
Athabasca Health Authority	92	86.4	77.4	91.7	97.6	43.9	97.3	73	93.3	85.4	88.2	80.4	86.8
Keewatin Yatthé	77.1	57.2	52.8	72.6	85.7	42.2	84.5	62.1	86.2	77.6	80.2	50.7	53.9
Mamawetan Churchill River	83.4	70.2	66.1	82.8	93.9	62.2	93.3	73.3	95	83.6	78.8	65.3	63.6
Peer Group H													
Prairie North	85.6	74.3	74.3	82.7	87.6	60.2	88.2	72.3	84	79.9	77.8	74	74.4
Prince Albert Parkland	81.9	66.9	68.6	78.7	86	52.3	87.4	69.7	85.1	78.6	79.8	76.6	83.4

- Two years of coverage data in the 13 age-dose/up-to-date categories are provided by former RHA and Athabasca Health Authority (AHA). Yellow highlight indicates rates below the provincial coverage rate.
- Pertussis vaccine is recommended at two, four, six and 18 months, with a booster dose between four and six years of age. If the primary series is delayed or interrupted, the schedule can be adjusted to bring the child up-to-date for protection. Data for three, five, eight, 12, 20, and 24 months; and five, seven, 13, 15 and 17 years are shown with seven, 13, 15 and 17 years reported as up-to-date.
- Up-to-date at seven years:
 - Children who received the four-dose primary series at two, four, six and 18 months and a booster dose between four and six years of age AND
 - Children who received four doses before the age of four years and one booster dose after the age of four years with at least a 24-week interval between the third and fourth doses, and the fourth and fifth doses AND
 - Children who received four doses of vaccine with the fourth dose administered after the age of four years and at least 24 weeks after the third dose AND
 - Children who received three doses of vaccine with the third dose at seven years and at least 24 weeks after the second dose.
- Up-to-date at 13 years:
 - a) to b) for seven years AND
 - Children who received five doses with the third dose administered before four years of age, the fourth dose at least 24 weeks after the third dose, and the fifth dose at or after 11 years of age and at least 24 weeks after the fourth dose AND
 - Children who received four doses with the fourth dose administered at or after 11 years of age and a 24-week interval between the second and third doses, and third and fourth doses AND
 - Children who received three doses with the third dose administered at or after 11 years of age and a 24-week interval between the second and third doses and a one-month-interval between first and second doses.
- Up-to-date at 15 and 17 years:
 - Children who received the four-dose primary series at two, four, six and 18 months and two booster doses at four to six years and Grade 8 (usually between 12 and 14 years of age) AND
 - b) to d) for 13 years.
- In 2018, eight former RHAs and AHA exceeded provincial coverage rates for four doses at 20 months and five years and up-to-date at 13 and 15 years while four RHAs were below.
- In 2018, three-dose coverage rate was higher among the 20-month age group compared to the eight-month age group: 90.7% vs. 80.4%. Likewise, the four-dose coverage rate for the 24-month age group was higher than the 20-month age group: 80.8% vs. 68.2%. A similar trend was observed for 2017 as well.
- In 2018, one former RHA was below the provincial rate in all thirteen age-dose categories and one was below in all but one category.
- In 2018, four former RHAs were at or above the provincial rate in all age-dose categories and one was at or above the provincial average in all but two categories.
- Coverage rates for former health regions in Peer Groups F and H should be interpreted with caution (see Data Notes).

SURVEILLANCE CASE DEFINITION: SASKATCHEWAN CDC MANUAL

Respiratory and Direct Contact Pertussis

Notification Timeline:

From Lab/Practitioner to Public Health: Immediate.

From Public Health to Ministry of Health: Within 2 weeks.

Public Health Follow-up Timeline: Immediate

Case Definition (adopted from Public Health Agency of Canada, 2008)

Confirmed Case	<p>Laboratory confirmation of infection:</p> <ul style="list-style-type: none"> isolation of <i>Bordetella pertussis</i> from an appropriate clinical specimen <p>OR</p> <ul style="list-style-type: none"> detection of <i>B. pertussis</i> DNA from an appropriate clinical specimen <p>AND one or more of the following:</p> <ul style="list-style-type: none"> cough lasting 2 weeks or longer paroxysmal cough of any duration cough with inspiratory "whoop" cough ending in vomiting or gagging, or associated with apnea <p>OR</p> <p>Epidemiologic link to a laboratory-confirmed case AND one or more of the following for which there is no other known cause:</p> <ul style="list-style-type: none"> paroxysmal cough of any duration cough with inspiratory "whoop" cough ending in vomiting or gagging, or associated with apnea
Probable Case	<p>Cough lasting 2 weeks or longer in the absence of appropriate laboratory tests and not epidemiologically linked to a laboratory-confirmed case</p> <p>AND one or more of the following, with no other known cause:</p> <ul style="list-style-type: none"> paroxysmal cough of any duration cough with inspiratory "whoop" cough ending in vomiting or gagging, or associated with apnea
Suspect Case	<p>One or more of the following, with no other known cause:</p> <ul style="list-style-type: none"> paroxysmal cough of any duration cough with inspiratory "whoop"




Photo Courtesy of Centers for Disease Control

DATA NOTES

Case Data Source: The Saskatchewan Integrated Public Health Information System (iPHIS) is a provincially mandated integrated client -centered case management information system that supports public health surveillance. Confirmed cases must meet the provincial surveillance case definition.

There are 10 peer groups used by Statistics Canada, each identified by a letter (A to J). A peer group consists of former health regions with similar socio-economic characteristics which facilitates comparisons within a peer group. The twelve former health regions and one health authority in Saskatchewan fall into four groups identified by letters A, D, F and H. The peer groups in this report are based on Statistics Canada 2011 peer groupings and should not be compared to current Statistics Canada peer groupings (2014).

Vaccine Coverage Data Source: Panorama is a comprehensive, integrated public health information system. Of the five modules in the system, four have been implemented: vaccine inventory, immunization, investigations and outbreaks management. When fully functional, it will help public health professionals work together to effectively manage vaccine inventories, immunizations, investigations, outbreaks and family health. Panorama's immunization module replaced the former Saskatchewan Immunization Management System (SIMS), on January 27, 2015. SIMS had been used province-wide since 2001. To learn more, please visit: www.ehealthsask.ca/services/panorama/Pages/default.aspx.

Most FNIHB-SK and NITHA communities, with the exception of those in Athabasca Health Authority (AHA), are not currently using Panorama. Therefore, immunization data for most First Nations children are missing or are incomplete. This report includes only those children with Saskatchewan health coverage and registered in Panorama under a former health region or AHA as of April 1, 2019. In other words, children with Saskatchewan health coverage and registered in Panorama under FNIHB-SK or NITHA jurisdiction are excluded (including those from FNIHB-SK and NITHA communities in AHA). This means this report does not include coverage statistics for the entire provincial or regional population.

The four-dose primary series pertussis containing vaccine is administered as diphtheria, tetanus, acellular pertussis, inactivated polio & *Haemophilus influenzae* type B (DTaP-IPV-Hib). The first booster at four to six years of age is DTaP-IPV vaccine and the second and final booster at Grade 8 is tetanus, diphtheria & acellular pertussis (Tdap) vaccine. Immunization coverage is based on those who turned three, five, eight, 12, 20 and 24 months, and five, seven, 13, 15 and 17 years by December 31 of 2017 and 2018. For example, the immunization coverage for seven-year-old children in 2018 is based on clients who were born in 2011 and the immunization doses they received by their seventh birthday.