

IMVAMUNE® Smallpox and Monkeypox Vaccine

What is monkeypox?

- Monkeypox is a rare viral illness that causes fever, headache, swollen lymph nodes and lethargy, followed by the development of a rash over a person's body.
- Monkeypox is spread through direct contact with monkey pox lesions or fluids from lesions of infected animals or people, through respiratory droplets from an infected person, from a pregnant woman to her baby, or indirect contact with items (e.g. bedding, towels).
- Monkeypox virus is not considered to be a sexually transmitted infection (STI) but it can spread through close contact during sexual activity.
- The virus may be spread to others even if the exposed person is not showing any symptoms.

What are the symptoms of monkeypox?

- The time between becoming infected and developing symptoms is 6 to 13 days (range is 5 to 21 days).
- Symptoms last from 2 to 4 weeks, and the disease usually resolves on its own without treatment.
- Initial symptoms may include fever, chills, muscle soreness, fatigue, headache, backache, swollen lymph nodes, sore throat, cough, vomiting or diarrhea.
- Sores may develop on or by the mouth or eyes.
- Within a few day of these symptoms, a rash characterized by a flat, red area on the skin that is covered with small bumps starts on the face, legs or arms, and can spread to other parts of the body including the palms of the hands, soles of the feet and genitals.
- Within 1 to 2 weeks, the rash becomes raised blisters that crust and scab over 2 to 3 weeks, and can be extremely itchy. Lesions are considered infectious until the scabs fall off and new skin can be seen.

Complications from monkeypox may include inflammation of the brain, blood poisoning, bacterial skin infections from scratching, dehydration, eye inflammation including the cornea, pneumonia and death.

How can monkeypox be prevented?

- Imvamune® protects against infection from the virus that causes monkeypox. The vaccine can prevent illness or make the illness less severe in those exposed to the virus.

- Avoid any close face-to-face contact with a person with monkeypox.
- Avoid touching the sores and lesions of a person with monkeypox.
- Avoid sharing or touching the bed linens, towels and clothing of a monkeypox case.
- Avoid sharing items such as toothbrushes, razors and hair brushes/combs with a person with monkeypox.
- Practice good hygiene (e.g. handwashing).

Who can get this vaccine for free?

- Those who are identified as a close contact to a probable or confirmed case of monkeypox, ideally within 4 days and up to 14 days from last exposure.
- Individuals who are transgender or those who self-identify as two spirit, bisexual, gay or men who have sex with men (MSM), or individuals who have sex with MSM; **AND one or more of the following:**
 - Having had a sexually transmitted infection in the past year.
 - Having two or more sexual partners or who are in a relationship where at least one of the partners has other sexual partners.
 - Engaging in sexual contact in sex-on-premises venues, events or social gatherings (sauna, bath house, club).
 - Have had or plan to have sexual contact with an anonymous partner (at an event or via a hook-up app).
 - Are planning to travel in the next three months to an area in Canada or internationally currently reporting monkeypox cases (<https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON396>)
- Individuals who self-identify as sex workers, regardless of their self-identified gender.
- Individuals who work or volunteer at places, events or social gatherings where there is sexual activity (sauna, bath house, club).
- Imvamune® may be offered to people who are immunocompromised, pregnant or breastfeeding, or under 18 years if they are at high risk of exposure, or following a known exposure to the virus.

Who should not get this vaccine?

- Those with a history of monkeypox infection do not need to be vaccinated.
- Persons who have a serious illness, with or without a fever, should delay immunization.
- Persons who have had a life-threatening reaction to a previous dose of a smallpox vaccine, or any components of the vaccine.
- Egg-allergic individuals may be immunized except if there is a known previous anaphylactic reaction to egg. Egg-allergic vaccine recipients should be kept under observation for 30 minutes following the administration of this vaccine.
- Talk to your healthcare provider if you:
 - Have a weakened immune system.
 - Have atopic dermatitis.
 - Are pregnant or breastfeeding.
- Certain people at increased risk of a condition called myocarditis (swelling of the heart muscle), including adolescent or young adult males, might consider waiting 4 weeks after IMVAMUNE® vaccination before getting an mRNA COVID-19 vaccine.

How is this vaccine given?

- The vaccine is given as an injection in the upper arm (subcutaneously) or under the skin of the arm (intradermally).

What are common reactions to this vaccine?

Vaccines are very safe. It is much safer to get this vaccine than to get monkeypox.

- The vaccine is made using weakened live vaccinia virus and cannot cause smallpox, monkeypox, or any other disease.
- Temporary pain, redness, swelling, warmth, hardness and itching at the injection site. Injection site reaction may be more common and last longer when the vaccine is administered under the skin (intradermally).
- Fatigue, headache, muscle and joint aches, headache, nausea, decreased appetite, fever or pain in the arm.
- These reactions may last for a few days.
- Persons who have atopic dermatitis may have more intense reactions or have a flare up after getting this vaccine.
- If you have signs and symptoms of a cardiac disorder (e.g. chest pain or discomfort, difficulty breathing or palpitations), seek medical attention immediately.

Other Considerations:

- If you need another live vaccine, get it at the same time or wait at least 4 weeks.
- If you need TB skin testing, the vaccine should be given on the same day or you should wait for 4 weeks before immunizing.

It is important to stay in the clinic for 15 minutes after getting any vaccine because there is an extremely rare possibility of a life-threatening allergic reaction called anaphylaxis.

This may include hives, difficulty breathing, or swelling of the throat, tongue or lips. **If this happens after you leave the clinic, call 911 or the local emergency number.** This reaction can be treated, and occurs in less than one in one million people who get the vaccine.

For more information, contact your local public health office, your physician, nurse practitioner, HealthLine online or by calling 811.

Who should you report reactions to?

- Report any adverse or unexpected reactions to your local public health nurse, your doctor, or nurse practitioner as soon as possible.

Talk to a public health nurse:

- If you have questions or concerns about your reaction to an immunization.
- If you had to go to a doctor, a hospital or to a health centre with a symptom that might be related to immunization.

What does this vaccine contain?

IMVAMUNE® is an attenuated live non-replicating viral vaccine that contains Tris buffer (Tris-hydroxymethyl-amino methane, sodium chloride, water for injection and hydrochloric acid), Trometamol (Tris-hydroxymethyl-amino methane), sodium chloride, water for injection. The vaccine contains trace amounts of host cell DNA and protein, benzonase, gentamicin and ciprofloxacin. Contains no preservatives or adjuvants.

Provincial immunization fact sheets are available at www.saskatchewan.ca/immunize.

References: Product monograph (IMVAMUNE® 2021).

Acetaminophen (Tylenol, Tempra) may be given for fever or soreness. ASA (Aspirin) must NOT be given to anyone younger than 18 years old because of the risk of Reye syndrome.

