

Crop Report

For the Period July 26 to August 1, 2022

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 14, August 4, 2022

Harvest has just begun in the province. These few fields are mainly in west central and southwest regions where crops are further ahead in development. Parts of these regions received rainfall and this has delayed further harvest activities such as combining and desiccating. In the eastern regions harvest is at least seven-10 days away, since crops are behind in development and in some areas are just now beginning to fill with seed, although some earlier seeded crops are close to being ready for desiccation. Rainfall varied significantly across the province last week with some areas getting nothing and others experiencing large, localized storms that resulted in flooding and crop damage. The Unity area received 53 mm, the Briercrest area 49 mm, the Avonlea area 40 mm, the Mayfair area 37 mm, the Lake Lenore area 24 mm, the Bulyea area 17 mm and the Swift Current area nine mm.

One year ago

Harvest had begun with three per cent of the crop now combined, ahead of the 5-year average of less than one per cent. Hot weather and little rain lead to further decline in topsoil moisture across the province, after harvest producers hope for a long soaking rain to replenish the soil.

Follow the 2022 Crop Report on Twitter @SKAgriculture

Even with the rainfall received this past week, topsoil moisture across the province continues to decline slightly. Cropland topsoil moisture is rated as four per cent surplus, 64 per cent adequate, 25 per cent short and seven per cent very short. Hay and pasture land topsoil moisture is rated as two per cent surplus, 65 per cent adequate, 20 per cent short and 13 per cent very short. After receiving much more rain than last year, pastures in some areas of the province have recovered from the 2021 drought and pasture condition ratings have improved tremendously. Pasture conditions are rated as 16 per cent excellent, 41 per cent good, 25 per cent fair, twelve per cent poor and six per cent very poor. Many pastures in the west still had cattle pulled off due to a lack in vegetation or dried up water sources.

The majority of crop damage this week was due to minor flooding, drought, disease, wind, grasshoppers and hail. Several storms crossed the province over the last week leaving behind some substantial crop damage. Some crops were laid down and lodged by strong winds and heavy rain while others were destroyed by hail. Hailstorms damaged crops from Marengo all the way to east of Lake Diefenbaker; buildings, machinery and vehicles were also damaged.

Follow the 2022 Crop Report on Twitter at @SKAgriculture.

For further information, contact Matthew Struthers, AAg,
Crops Extension Specialist, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3714, Email: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at saskatchewan.ca/crop-report.



Agriculture and
Agri-Food Canada

SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan

Saskatchewan Harvest Progress - August 1, 2022

*Other - crop that will not be harvested due to weather, insect or disease damage or will be greenfeed or silage

Winter Wheat	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	100	0	0	0	0
southwest	68	28	2	2	0
east central	100	0	0	0	0
west central	86	14	0	0	0
northeast	100	0	0	0	0
northwest	100	0	0	0	0
provincial	86	12	1	1	
Fall Rye	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	100	0	0	0	0
southwest	62	13	12	2	11
east central	100	0	0	0	0
west central	88	12	0	0	0
northeast	94	0	6	0	0
northwest	100	0	0	0	0
provincial	89	5	3	1	2
Spring Wheat	% Standing	% in swath	% ready to straight combine	% combined	
southeast	100	0	0	0	
southwest	100	0	0	0	
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	100	0	0	0	
Durum	% Standing	% in swath	% ready to straight combine	% combined	
southeast	100	0	0	0	
southwest	99	1	0	0	
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	n/a	n/a	n/a	n/a	
provincial	99	1	0	0	
Barley	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	99	0	0	0	1
southwest	99	0	0	0	1
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	99	0	0	0	1
Oats	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	100	0	0	0	0
southwest	92	0	0	0	8
east central	100	0	0	0	0
west central	98	0	0	0	2
northeast	100	0	0	0	0
northwest	100	0	0	0	0
provincial	98	0	0	0	2

Canaryseed	% Standing	% in swath	% ready to straight combine	% combined	
southeast	100	0	0	0	
southwest	100	0	0	0	
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	100	0	0	0	
Flax	% Standing	% in swath	% ready to straight combine	% combined	
southeast	100	0	0	0	
southwest	100	0	0	0	
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	100	0	0	0	
Canola	% Standing	% in swath	% ready to straight combine	% combined	
southeast	100	0	0	0	
southwest	100	0	0	0	
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	100	0	0	0	
Mustard	% Standing	% in swath	% ready to straight combine	% combined	
southeast	100	0	0	0	
southwest	100	0	0	0	
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	100	0	0	0	
Soybeans	% Standing	% in swath	% ready to straight combine	% combined	
southeast	100	0	0	0	
southwest	100	0	0	0	
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	100	0	0	0	
Field Peas	% Standing	% in swath	% ready to straight combine	% combined	
southeast	100	0	0	0	
southwest	87	0	10	3	
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	97	0	2	1	
Lentils	% Standing	% in swath	% ready to straight combine	% combined	
southeast	99	0	1	0	
southwest	93	0	3	4	
east central	100	0	0	0	

west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	97	0	2	1	
Chickpeas	% Standing	% in swath	% ready to straight combine	% combined	
southeast	100	0	0	0	
southwest	100	0	0	0	
east central	100	0	0	0	
west central	100	0	0	0	
northeast	100	0	0	0	
northwest	100	0	0	0	
provincial	100	0	0	0	

Southeastern Saskatchewan:

- Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE – Radville, Minton and Lake Alma areas

Harvest has not begun as crops are behind their normal development due to late seeding dates and cool rainy weather delaying growth. Any rain now will keep crops from drying up on hot windy days and will allow crops to fill their pods or heads with seed. Other crops that are still flowering need some warm sunny days to quicken their maturity. Earlier seeded crops are still not ready for harvest, but some are ready for desiccation and producers will be conducting those activities in the coming week. Yield estimates have not been finalized for the region, however many producers are expecting a far better crop in comparison to 2021.

The Briercrest area received 49 mm of rain last week, the Bienfait and Avonlea areas 40 mm and the Glenavon and Odessa areas 20 mm. While this rain has delayed haying operations in the region, it is very good for crops as they carry out seed filling before beginning to ripen for harvest. Cropland topsoil moisture is rated as twelve per cent surplus, 82 per cent adequate, five per cent short and one per cent very short. Hay and pasture land topsoil moisture is rated as five per cent surplus, 87 per cent adequate, seven per cent short and one per cent very short.

After receiving early season moisture in the form of snow and rain, along with regular rainfall since April 1, many pastures in the region have greatly improved and are now able to support cattle without issue. Pasture conditions are rated as 41 per cent excellent, 46 per cent good, eleven per cent fair, one per cent poor and one per cent very poor.

The majority of crop damage this week was due to heavy rain, wind, hail and grasshoppers. Strong winds and heavy rains resulted in some crop lodging, this will make harvest more difficult and possibly reduce quality of harvested seed. A hailstorm hit the northwest corner of the region and resulted in damage to crops, buildings and equipment.

Producers are busy finishing up haying and starting to gear up for harvest even if it is a few weeks away. Many producers will be starting their applications of desiccants in the coming week or so to prep crops for harvest.

Southwestern Saskatchewan:

- Crop District 3ASW – Coronach, Assiniboia and Ogema areas
- Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B – Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 – Consul, Maple Creek and Leader areas

Some producers in the region have begun to harvest crops, mainly winter cereals and pulse crops such as lentils and field peas. Rain will not be enough for the crops in the region. However, it would help pastures and hay land recover in hopes they would be able to sustain some fall grazing. Some cereal crops in the region are too poor to be harvested so they will likely be cut for feed as the hay yields in many parts of the region were very poor.

The southwest region did not receive widespread significant rainfall this week, but producers will take all they can. The Moose Jaw area received 30 mm, the Rockglen area 18 mm, the Goultdown area 14 mm and the Swift Current area nine mm. Cropland topsoil moisture is rated as 34 per cent adequate, 46 per cent short and 20 per cent very short. Hay and pasture land topsoil moisture is rated as 32 per cent adequate, 28 per cent short and 40 per cent very short.

Pasture conditions are rated as one per cent excellent, 19 per cent good, 29 per cent fair, 35 per cent poor and 17 per cent very poor. Producers in the region hope for more rain to help support pasture growth and recharge water sources for cattle. Some have noted concern for winter feed shortages due to reduced hay yields.

The majority of crop damage this week was due to drought, heat, wind and hail. A large hailstorm passed through the region destroying several fields and leaving many more severely damaged. Grasshoppers also continue to wreak havoc on crops in the region, some have lost a considerable amount of their fields to insect pests.

East-Central Saskatchewan:

- Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A – Lumsden, Craik, Watrous and Clavet areas

Harvest has not commenced in the region as crops are late due to delayed seeding in the spring and excess moisture slowing down development. Crops on lighter soils have begun to dry down and producers expect these crops to be ready for harvest in about one to two weeks. Producers will begin desiccating crops as soon as they are ready so that they may dry down faster and make harvest less difficult.

Rainfall this week kept crops on lighter sandier soils from drying down prematurely and helped pastures. The Rhein and Wadena areas received 43 mm, the Langenburg area 27 mm, the Calder area 19 mm and the Bulyea area 17 mm. While many parts of the region have not suffered from lack of rain, others have missed the rains over the past weeks and are becoming quite dry. Cropland topsoil moisture is rated as eight per cent surplus, 70 per cent adequate, 20 per cent short and two per cent very short. Hay and pasture land topsoil moisture is rated as three per cent surplus, 75 per cent adequate, 18 per cent short and four per cent very short.

Pasture conditions are rated as 20 per cent excellent, 51 per cent good, 23 per cent fair, three per cent poor and three per cent very poor. More adequate soil moisture this year has allowed pastures to recover from the stress of last year's drought and producers are very happy with how their grasslands look.

The majority of crop damage this week was due to thunderstorms that brought strong winds, heavy rains and hail which left many crops damaged in their wake. Localized flooding resulted where heavy rains occurred which also caused cereal crops to become lodged. Some producers who are struggling to keep pea aphids under economic thresholds in their pulse crops, will continue to spray for aphids until their crops are near to the point where they can be harvested.

West-Central Saskatchewan:

- Crop District 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A – Rosetown, Kindersley, Eston and Major areas
- Crop District 7B – Kerrobert, Macklin, Wilkie and Biggar areas

Harvest of winter cereals and pulse crops has just begun in the region. In the driest areas of the region, crops are struggling through the hot weather and have begun too rapidly mature. Producers can no longer wait and must begin to harvest the ripest of their crops. In areas that were lucky to receive some timely rains the crops do not appear to have suffered as much although they still are not expected to yield well. The recent rain may help later seeded crops, but most crops are so far along that any rain now will not improve their conditions.

The rain received this past week will hopefully give pastures and hay land some relief and allow for some regrowth. This would give producers the chance to fall graze some of their land. The Unity area received 53 mm, the Dinsmore area 46 mm, and the Smiley and Rosetown areas 11 mm. Cropland topsoil moisture is rated as 65 per cent adequate, 31 per cent short and four per cent very short. Hay and pasture land topsoil moisture is rated as 53 per cent adequate, 39 per cent short and eight per cent very short.

Warm, dry conditions are drying down pastures in the region and producers are hoping for more rain to help rejuvenate pasture growth. Pasture conditions are rated as 34 per cent good, 44 per cent fair, 18 per cent poor and four per cent very poor.

The majority of crop damage this week was due to drought, heat, wind and hail. Large swaths of the region were rocked by hail storms this past week leaving many fields cut right down to nothing. Grasshoppers are still large contributors to crop damage as well. Producers have sprayed multiple times to keep them under control but in some cases the grasshoppers are too many.

Northeastern Saskatchewan:

- Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas

Producers have noted that their crops look very good across the region but they could use a little more rain to help the crops fill with seed before they begin to ripen and dry down in time for harvest. Pulses and cereal crops appear to be the most mature with oilseeds being the most behind. With a few weeks of hot sunny weather crops in the region will begin to ripen quickly. Producers are likely 10 days or more from starting their harvest operations.

There were spotty rainstorms in the region this past week, the Prince Albert area received 29 mm, the Lake Lenore area 24 mm and the Melfort and Kinistino areas eight mm. Cropland topsoil moisture is rated as 80 per cent adequate, 19 per cent short and one per cent very short. Hay and pasture land topsoil moisture is rated as one per cent surplus, 77 per cent adequate and 22 per cent short.

Pastures in the region are doing much better than they were a year ago, with ample rainfall this growing season they have been able to recover from the drought stress of 2021. Pasture conditions are rated as twelve per cent excellent, 72 per cent good, 14 per cent fair and two per cent poor.

The majority of crop damage this week was due to strong winds that resulted in lodged crops, cereals crops such as barley, oats and wheat were mainly affected. There was a hailstorm in the northern half of the region that resulted in crop damage but the severity is not currently known.

Northwestern Saskatchewan:

- Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas


Harvest has not started in the northwest but crops have begun to turn and some will be ready for swathing or desiccating in the next seven-10 days. Rain still has the potential to help later seeded crops fill with seed and producers are hopeful they receive a little more before crops completely dry down. Producers are positive about their potential crop yields, stating they appear to be up from last year.

Rainfall in the northwest region ranged from nil to 51 mm in the Nielburg area, the Glaslyn area received 41 mm, the Duck Lake and Mayfair areas 37 mm and the Livelong area 24 mm. Cropland topsoil moisture is rated as one per cent surplus, 77 per cent adequate, 21 per cent short and one per cent very short. Hay and pasture land topsoil moisture is rated as one per cent surplus, 81 per cent adequate, 17 per cent short and one per cent very short.

Pastures in the region benefited greatly from early season moisture and regular rainfall through the spring and their conditions have improved since last year. They are rated as 19 per cent excellent, 40 per cent good, 38 per cent fair and three per cent poor.


The majority of crop damage this week was due to heavy rains and strong winds causing some crops to lodge, there was some minor hail damage as well. Producers are worried that either a severe hailstorm or an early frost will get to their crops before they are able to harvest.

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

from July 26 to August 1, 2022



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period from July 26 to August 1, 2022


Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	N/A	369	4B	139	Gull Lake	N/A	153	7A	259	Snipe Lake	4	158
	3	Enniskillen	14	416		229	Miry Creek	12	179.9		287	St. Andrews	16	132
	4	Coal Fields	40	606		231	Happyland	1	116		288	Pleasant Valley	14	132
	6	Cambria	25	297	5A	181	Langenburg	trace	380		290	Kindersley	N/A	145.7
	32	Reciprocity	N/A	256		183 A	Fertile Belt	N/A	549		292	Milton	18	272.5
	34	Browning	N/A	0		211	Churchbridge	27	284		317 A	Marriott	15	143
	61	Antler	N/A	386		213	Saltcoats	1	286		317 B	Marriott	11	210.3
	64	Brock	19	364		215	Stanley	5	379		319	Winslow	N/A	75.1
	65	Tecumseh	11	332		216	Tullymet	NIL	156.7		320 A	Oakdale	24.5	244.7
1B	95	Golden West	N/A	0		217	Lipton	5	379		320 B	Oakdale	12	235
	122	Martin	22	495		241	Calder	19	301		321	Prairiedale	11	296.5
	123	Silverwood	24	460		243	Wallace	N/A	240	7B	347	Biggar	13	172
	124	Kingsley	6	373		245 A	Garry	5	338		350	Mariposa	20	138
	125 A	Chester	20	329		245 B	Garry	8	310		351	Progress	12	138
	125 B	Chester	28	285		246 A	Ituna Bon Accord	trace	221		377	Glenside	24	129
	151	Rocanville	18	436		246 B	Ituna Bon Accord	1	292.3		378	Rosemount	30	177
	154 A	Elcapo	5	285		247	Kellross	trace	245.5		382	Eye Hill	N/A	146.5
	155	Wolseley	7	300		248	Touchwood	NIL	196		409 A	Buffalo	N/A	0
2A	67	Weyburn	N/A	311	5B	273	Sliding Hills	43	250		409 B	Buffalo	26.5	103.3
	68	Brokenshell	25	396		275	Insinger	N/A	331	8A	410	Round Valley	52.5	120.5
	96	Fillmore	N/A	0		276	Foam Lake	25	258		394	Hudson Bay	5	191
	97	Wellington	18.5	274.5		277 A	Emerald	2	321		395	Porcupine	N/A	269
	100	Elmsthorpe	40	230		277 B	Emerald	2	253		397 A	Barrier Valley	N/A	251.6
2B	127 A	Francis	20	377.5		301	St.Philips	21.1	289.1		397 B	Barrier Valley	N/A	133
	127 B	Francis	20	159.1		305	Invermay	22.5	304.5		427	Tisdale	N/A	0
	129	Bratt's Lake	42	360		307	Elfros	7	239		428	Star City	N/A	202
	131	Baildon	49	220		308 A	Big Quill	5	203		456	Arborfield	11	206
	156 A	Indian Head	7.1	255.4		308 B	Big Quill	N/A	0		457 A	Connaught	N/A	0
	156 B	Indian Head	18	360		331	Livingston	12	265		457 B	Moose Range	N/A	118.8
	159 A	Sherwood	N/A	242.5		334	Preeceville	trace	18		486	Moose Range	28	203.2
	159 B	Sherwood	NIL	67		336	Sasman	N/A	117		487	Nipawin	N/A	215
	160	Pense	9	118		337	Lakeview	42	267	8B	369	St. Peter	NIL	178
	161 A	Moose Jaw	33	209		366	Kelvington	2	212.5		370 A	Humboldt	NIL	213
	162 A	Caron	28.1	218.5		367	Ponass Lake	N/A	102.5		370 B	Humboldt	N/A	0
	162 B	Caron	44	276.5	6A	189	Lumsden	N/A	70		371	Bayne	NIL	190
	191	Marquis	N/A	0		190 A	Dufferin	N/A	220		372	Grant	7	185.4
3ASE	38 A	Laurier	13.8	305.2		190 B	Dufferin	8	208		400	Three Lakes	24	248.5
	38 B	Laurier	18	344		190 C	Dufferin	18	171		429	Flett's Springs	6	242
3ASW	10	Happy Valley	N/A	292		190 D	Dufferin	N/A	95		430	Invergordon	8	223
	40	Bengough	N/A	0		219 A	Longlaketon	N/A	203		459	Kinistino	8	305
	43	Old Post	18	227.5		219 B	Longlaketon	20	189	9AE	460	Birch Hills	3.1	195.7
	73 A	Stonehenge	NIL	203.4		220	McKillop	17	223		461	Prince Albert	12.5	155
	73 B	Stonehenge	N/A	150.8		221	Sarnia	11.7	210.9		488	Torch River	7	222
	74	Wood River	NIL	176.4		222	Craik	13	269		491	Buckland	19	261
3AN	102	Lake Johnston	N/A	211		251	Big Arm	NIL	166		520	Paddockwood	29	143
	103	Sutton	N/A	72		252	Arm River	N/A	115		521	Lakeland	29	143
	132 A	Hillsborough	N/A	0		279	Mount Hope	0.8	83.4	9AW	405	Great Bend	N/A	29
	132 B	Hillsborough	30	277		282	McCraney	10	169		406	Mayfield	27	63.8
	193	Eyebrook	7	192		310	Usborne	1.4	222		435	Redberry	6	159
3BS	75	Pinto Creek	trace	172		312	Morris	1.2	197.8		436	Douglas	19	118
	77	Wise Creek	NIL	152		339	Leroy	trace	218.6		437	North Battleford	N/A	173.5
	78	Grassy Creek	NIL	155		340	Wolverine	N/A	0		463	Duck Lake	13	340
	106 A	Whiska Creek	NIL	164		341	Viscount	N/A	0		466	Meeting Lake	37	231
	106 B	Whiska Creek	NIL	228.5		343	Blucher	12	251		467 A	Round Hill	27	180
	107	Lac Pelletier	N/A	0	6B	223 A	Huron	9	147		467 B	Round Hill	36	185
	108	Bone Creek	N/A	184		223 B	Huron	15	202		493	Shellbrook	14	146.5
3BN	138	Webb	trace	233.8		223 C	Huron	N/A	0		496	Spiritwood	N/A	310
	165	Morse	14	212.5		284 A	Rudy	10	126		497	Medstead	21	289.5
	168 A	Riverside	9	142		285	Fertile Valley	5	107	9B	440	Hillsdale	51	231
	168 B	Riverside	N/A	136.6		286	Milden	46	223		442	Manitou Lake	42.7	227.1
	226	Victory	N/A	0		314	Dundurn	7	163		498	Parkdale	41	351
	228	Lacadena	9	135.5		344	Corman Park	4	232		499 A	Mervin	24.6	340.6
	257	Monet	1.5	64.5		376	Eagle Creek	33	124		499 B	Mervin	24	237.5
4A	51	Reno	NIL	179		403 A	Rosthern	5	161		501 A	Frenchman Butte	11	217
	79 A	Arlington	NIL	158		403 B	Rosthern	N/A	91		501 B	Frenchman Butte	N/A	60
	79 B	Arlington	N/A	237							501 C	Frenchman Butte	9	238
	109	Carmichael	N/A	0							502	Britannia	N/A	96
	110	Piapot	N/A	181.9							561	Loon Lake	22	218
	111	Maple Creek	N/A	173							588	Meadow Lake	N/A	227
											622	Beaver River	6	205.4

Municipality No: A, B, C and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.


Cumulative Rainfall

from April 1 to August 1, 2022



Cropland Topsoil Moisture Conditions

August 1, 2022



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan

1:3,500,000
0 25 50 100 150 200
Kilometers

Projection: UTM Zone 13 Datum: NAD83




Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture

August 3, 2022

Hay and Pasture Topsoil Moisture Conditions


August 1, 2022



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Minimum Temperature


from July 26 to August 1, 2022



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Maximum Temperature

from July 26 to August 1, 2022



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.