

Community Respiratory Illness Surveillance Program

(CRISP) Situation Report: March 16, 2023 (Reporting Period February 26 – March 11, 2023)

Summary

- COVID-19 activity continues at stable levels, with an increase in the most recent reporting period. Respiratory Syncytial Virus (RSV) cases have dropped by over 50% (from 85 to 41 cases).
 - The number of positive lab tests for COVID-19 increased from 202 in the week Feb 12- Feb 18 to 237 in the most recent week. Test positivity increased from 6.5% to 8.1%.
 - RSV cases trended downward over the past four weeks. The proportion of cases is now almost equal among preschool aged children and seniors.
- Influenza activity in Saskatchewan increased this past week with the emergence of type B activity, typically seen in the late influenza season.
- COVID-19, RSV and other respiratory viruses account for 56%, 14% and 24% of hospitalizations due to respiratory illnesses across the province, respectively.
- Sentinel indicators of respiratory transmission in the community fluctuates, as the season wanes;
 - Weekly visits to the Saskatchewan emergency departments for respiratory-like illness (RLI) have remained stable at 15.5 in the previous week to 15.1 per 1,000 this week.
 - Calls to 811 HealthLine for RLIs decreased from 65.9 to 54.0 per 1,000 calls.
 - School-illness absenteeism remained stable over the last four reporting weeks. The data indicated that 11.3% of Saskatchewan students were absent due to illness in the week of Feb 12 - 18, and 11.2% in the most recent surveillance week.
 - The most commonly detected organism in the sentinel provider program this week was Rhinovirus (common cold).
 - Regina wastewater viral load increased from moderate to moderate-high and the rest of Saskatchewan's viral load remained as moderate. The trajectory is decreasing in Saskatoon and North Battleford in the current week compared to the previous week.

COVID-19

- COVID-19 test positivity was 8.1% in the most recent week, an increase from 6.5% in the Feb 12- Feb 18 reporting week. Cases are largely in the 20 to 64 years (39%) and 65+ (45%) age groups.
- COVID-19 outbreaks in high-risk settings averaged five per week over the past four weeks – this has increased to thirteen outbreaks reported in the past two weeks.
- XBB.1.5 and its sublineages (denoted as XBB.1.5*) are the most commonly detected variants (41.2% of current reporting period), followed by BQ.1.1* (38.7%) and BQ.1* (9.1%).
- COVID-19 hospitalizations have increased from 126 for the previous two weeks, to 140 for the most recent two weeks.
- COVID-19 ICU admissions increased from 11 in the previous two weeks, to 15 in the most recent two weeks.
- The proportion of staffed inpatient beds occupied by COVID-19 patients slightly increased in recent two weeks to 5.9%, compared to the previous two weeks at 5.4% and 5.7%, respectively.

- In the past two weeks, Feb 26 to Mar 11, there have been 14 deaths among COVID-19 cases, one in 0 to 19 years age-group, two in 20 to 59 years age-group and 11 in the 60 years or older group.
- Having a COVID-19 booster in the last six months reduces the risk of a COVID-19 death eight times compared to unvaccinated individuals and over four times compared to those without a recent booster dose. Of those aged five years and older, 20% have received their latest booster dose in the last six months. With the exception of Regina, all areas of the province have less than 50% of their population up-to-date¹ for COVID-19 vaccines.
- Only 21% of individuals aged 12+ years have received a bivalent booster dose (n = 215,058 doses).
- Less than half of individuals aged 50+ have had more than one booster dose (47%).

Influenza

- Influenza cases were decreasing until type B activity emerged, typically seen in the latter part of an influenza season. Type B influenza generally has a mild clinical manifestation. Influenza activity was low in the past two weeks with sporadic cases distributed in communities throughout the province.
- Influenza test positivity fluctuated between 0.3% and 1.0% over the past four weeks. The emergence of type B activity is responsible for this increase.
- Influenza cases affected both preschool (3 cases) and working-aged people 20-64 years (3 cases) equally in the past week.
- No influenza outbreaks in high-risk settings were reported in the past month.
- There were two influenza hospitalizations and no ICU admissions reported during the weeks of February 26 to March 11, 2023.
- One death due to Influenza was reported in the past six weeks.
- The influenza immunization campaign launched October 11, 2022. As of February 28, 2023, 27% of the Saskatchewan population have received an influenza vaccine. This is a 10% decrease in doses administered compared to the same time last year and no change from the previous month.

Other Respiratory Viruses²

- RSV cases decreased from 164 in mid-January to 36 detections in the past week. Lab test positivity for RSV dropped from 12.4% to 5.2% over the past four weeks and has the lowest positivity of the respiratory viruses this week.
- RSV cases equally affected the pediatric age group (16 cases) and seniors group (14 cases).
- RSV hospitalizations have decreased from 75 for the previous two weeks, to 35 for the most recent two weeks. RSV ICU admissions have decreased from 14 for the previous two weeks, to 8 in the most recent two weeks. Children aged 0 – 19 years account for 72% of RSV hospitalizations and ICU admissions over the past two weeks.
- ‘Other’ respiratory viruses remained stable between 104 and 82 detections over the past month. The positivity rate remained stable week over week and is 5.6% this past week.
- Outbreaks from ‘other’ viruses in high-risk settings averaged four per week over the past four weeks.

* Represents all sublineages of Omicron

¹ Up-to-date = completed a primary series and at least one additional booster, age 5+ years

² Parainfluenza viruses 1 – 4 (PIV 1 – 4); Adenovirus (ADV); Rhinovirus (RV); Human Metapneumovirus (HMPV); NOTE: RSV test positivity now exceeds 1% and is reported separately.

Table 1: Viral indicators by surveillance period, February 26 – March 11, 2023

Report date	COVID-19 positive laboratory test	COVID-19 test positivity	COVID-19 outbreaks	Influenza positive laboratory test	Influenza test positivity	Influenza outbreaks	RSV positive laboratory test	RSV test positivity	'Other' ¹ positive laboratory test	'Other' ¹ test positivity	'Other' ¹ outbreaks
Mar 05 – 11, 2023	237	8.1%	6	7	0.8%	0	41	5.2%	93	5.6%	4
Feb 26 – Mar 04, 2023	217	7.1%	7	4	0.4%	0	85	9.1%	104	6.1%	7
Feb 19 – 25, 2023	194	6.6%	5	9	1.0%	0	96	12.1%	82	5.1%	2
Feb 12 – 18, 2023	202	6.5%	2	3	0.3%	0	118	12.4%	101	6.0%	4

Notes: ¹Parainfluenza viruses 1 – 4; Adenovirus; Human Metapneumovirus, seasonal Coronavirus. See Technical Notes for details.

Table 2: Patient-confirmed respiratory illness by age group, March 05 - 11, 2023

Age group (Years)	COVID-19 case count	Influenza case count	RSV case count	Others case count ¹
0-4	16 (8%)	3 (42.8%)	16 (44%)	N/A
5-19	15 (8%)	-	-	
20-64	78 (39%)	3 (42.8%)	6 (17%)	
≥65	90 (45%)	1 (14.4%)	14 (39%)	
Total	199 (100%)	7 (100%)	36 (100%)	93 (100%)

Notes: ¹Parainfluenza viruses 1 – 4; Adenovirus; Human Metapneumovirus, seasonal Coronavirus; age-specific data is unavailable (N/A) for other respiratory pathogens. Due to the rounding, total percentage may not add to 100%. See Technical Notes for further details.

Table 3: Sentinel* indicators by surveillance period, February 26 – March 11, 2023

Report date	School illness absenteeism >=10% ¹	RLI** ED visits per 1,000 ²	RLI** 811 calls per 1,000	COVID-19 Wastewater indicator ³	Sentinel provider test positivity	Most commonly detected virus: Sentinel providers
Mar 05 – Mar 11, 2023	11.2%	15.1	54.0	Moderate (n=6), Moderate-high (n=1)	33.3	Rhinovirus
Feb 26 – Mar 04, 2023	10.2%	15.5	65.9	Low (n=1), Moderate (n=5), High (n=1)	20.0	Parainfluenza virus
Feb 19 – 25, 2023	N/A [‡]	13.3	72.2	Low (n=1), Moderate (n=4), Moderate-high (n=1), High (n=1)	N/A [‡]	N/A [‡]
Feb 12 – 18, 2023	11.3%	21.2	52.4	Moderate (n=3), Moderate-high (n=3), High (n=1)	16.7	Human Metapneumovirus

Notes: *Sentinel surveillance are sampling programs representative of the population; ¹School absenteeism is the proportion of scheduled children who were absent from the class due to illness. The type of illness is not specified. [‡]Not Available (N/A). Schools were not in session and therefore there was no data available for the week of Feb 19-25. **Respiratory-like illness (RLI) ² Feb 26–Mar 04 based on reports from five of thirteen reporting areas; Mar 05–11 based on reports from six of thirteen reporting areas. ³Count of wastewater treatment facilities reporting low, moderate or high levels of viral load causing COVID-19 infection (see Technical Notes for details).

Table 4: Outcome, health care capacity and immunization coverage indicators by surveillance period, February 26 – March 11, 2023[‡]

Report date	Hospital admissions – COVID-19 ¹	ICU admissions – COVID-19	Hospital admissions – Influenza	ICU admissions – Influenza	Hospital admissions – RSV	ICU admissions – RSV	% of staffed inpatient beds occupied by COVID-19 patients ²	Deaths – COVID-19 ³	Deaths – Influenza ³	Proportion of population up-to-date – COVID-19 vaccine ⁴	Proportion of population immunized for Influenza vaccine ⁵
Mar 05 – Mar 11, 2023	74	4	2	0	8	5	5.9%	5	0	46.2%	27%
Feb 26 – Mar 04, 2023	66	11	0	0	27	3	5.9%	9	0	46.2%	
Feb 19 – 25, 2023	54	6	0	0	39	6	5.4%	7	1	46.2%	
Feb 12 – 18, 2023	58	5	0	0	36	8	5.7%	3	0	46.2%	

[‡]Additional information on hospital admission stratified by respiratory organism and age group is provided below in **Figure 5** and **6** respectively. Viral infection may not be the main reason for the admission.

³Fourteen COVID-19 deaths were reported in the past two weeks. One influenza deaths reported in the last four weeks.

Cases by respiratory organisms across the age groups

- From February 26 to March 11, 2023, there were 247 respiratory illness cases hospitalized with lab-positive Covid19 (140), influenza (2), RSV (35), other respiratory illnesses (60), and co-infected cases (10). Among the 247 cases, Covid19 lab positives (140), were among age groups of 0-19 (8), 20-59 (27), and ≥60 (105). Influenza lab positives (2) cases were among the age groups of 0-19 (1), 20-59 (1). For RSV lab positives (35) cases were among the age groups of 0-19 (25), 20-59 (3) and ≥60 (7). For other respiratory lab positives (60) cases were in the age group of 0-19 (37), 20-59 (7), and ≥60 (16). For Co-infections lab positives (10) cases were in the age group of 0-19 (8), 20-59 (1), and ≥60 (1).
- From February 26 to March 11, 2023, there were 34 respiratory illness cases admitted to the ICU with lab-positive Covid19 (15), influenza (0), RSV (8), other respiratory illnesses (8), and co-infected cases (3). Among the 34 cases, Covid19 lab positives (15) cases were among the age groups of 20-59 (6) and ≥60 (9). Influenza lab positives (0). For RSV lab positives (8) cases were among the age groups of 0-19 (6) and 20-59 (2). For other respiratory lab positives (8) cases were in the age group of 0-19 (6), 20-59 (1) and ≥60 (1). For Co-infections lab positives (3) cases were among the age groups of 0-19 (2) and 20-59 (1).

Notes:

¹ Because of the delay in date tested result, it affects the total number of Influenza (A/B), RSV and other respiratory virus admissions for a particular day. This lag in data impacts mostly the last couple of days from the day the report is updated. The counts for influenza, RSV, and other respiratory virus-associated hospital and ICU admissions refer to individuals with laboratory or point of care tests positive for influenza, RSV, and other respiratory viruses, respectively, occurring within four days before the admission date AND/OR at any point during the hospital stay. The counts for COVID-19 hospital and ICU admissions refer to individuals with laboratory tests positive for COVID-19 virus, occurring within 21 days before the admission date and/or at any point during the hospital stay or 7 days from the discharge. Episodes of care considers patients total movement within the health system related to their condition. It combines 2 or more admission from 2 or more different facilities, if they are transfers (No break in care). Transfer: Admission to any other hospital within 24 hours of discharge from previous hospital. Co-infected cases: positive for influenza and RSV or, positive for influenza and other respiratory virus or positive for RSV and other respiratory viruses or, positive for COVID-19 and influenza or, positive for COVID-19 and RSV or, Positive for COVID-19 and other resp viruses. Other includes: Parainfluenza 1-4, Adenovirus, Enterovirus, Human Metapneumovirus, Rhinovirus, Seasonal Coronavirus (O43, NL63, 229E, HKU1f.)


² 7-day average of percentage of acute inpatient beds staffed and in operation that are occupied by COVID-positive patients as of 8AM census. NA: Recent two weeks information not available

³ Includes deaths entered into Panorama IOM among lab-confirmed cases. Deaths reported based on the actual date of death. Deaths reported in previous periods subject to change due death reporting data lags

⁴ Up-to-date = completed a primary series and at least one additional booster, age 5+ years.


⁵ Up-to-date = received a vaccination within the current influenza season, age 6 months+. The most recent rate is as of February 28, 2023. This column will be updated at month end. See Technical Notes for details.

Figure 1: Epidemic curve, respiratory illness by respiratory organism and test positivity, August 29, 2021 – March 11, 2023




Data sources: Panorama IOM extracted on March 13, 2023 (COVID-19 cases)
 Respiratory Virus Detections Surveillance System (influenza and other respiratory) (RRPL extracted March 13, 2023)
 As of September 4, 2022, COVID-19 cases include new and reinfections.
 For the two weeks of Feb 26 to Mar 11, 2023, there were 390 COVID-19 cases (48 were 0 to 19 years; 135 were 20 to 59 years; and 207 were 60 years and older).
 For the two weeks of Feb 26 to Mar 11, 2023, there were 11 influenza lab detections
 For the two weeks of Feb 26 to Mar 11, 2023, there were 126 RSV detections
 For the two weeks of Feb 26 to Mar 11, 2023, there were 197 other viral lab detections (parainfluenza, adenovirus, human metapneumovirus, rhinovirus, coronavirus)

Figure 2: Percentage of SARS-CoV-2 variants by surveillance week starting*, January 22 – February 26, 2023



Data Source: Roy Romanow Provincial Laboratory, Saskatchewan Health Authority, as of March 13, 2023
 The most recent VOC data available from the Provincial database is as of surveillance week ending March 4, 2023
 * Surveillance week correspond to specimen collection date.

Figure 3: Proportion of residents by age group and COVID-19 vaccination status to March 11, 2023



Data sources: Panorama March 13, 2023; Saskatchewan Covered Population, 12-Nov-2022, Ministry of Health version (2022 Version 2)


Note:

Of those five years and older: 46.2% have completed their series and received a booster dose; 19.7% have received their latest dose in the last six months.

Of those 12 years and older, 20.7% (215,058) were administered a Moderna or a Pfizer bivalent.

As of March 11, 2023, cumulatively doses administered are as follows: dose 1, n=994,750; dose 2, n=929,464; dose 3, n=533,754; completed plus 2 or more additional doses, n=381,483; total doses = 2,839,451.

Figure 4: Booster dose within last 6 months reduces the risk of Hospitalization, ICU admission or Death (age-adjusted rates), June 24, 2022 – December 24, 2022




Data Source: Digital Health Analytics, Saskatchewan Health Authority

Definitions: Unvaccinated - Individuals with no record of vaccine received or vaccinated with first dose but less than 21 days from receiving the first dose.

Vaccinated with 2 doses - Individuals who have received their second dose for more than 14 days or their third dose is less than 14 days. Booster in the past 6 months - Individuals who have received any booster dose (3rd, 4th, 5th and so on) within the last 6 months. Details on methodology is provided in the Technical Notes.


Figure 5: The number of COVID-19, influenza, RSV, other respiratory viruses and co-infected cases admitted to hospital by week of the admission, February 12 – March 11, 2023*



Data sources: Digital Health Analytics, Saskatchewan Health Authority, Episode of Care methodology (Admission, Discharge, Transfer Database (ADT, RPPL, Panorama); data extracted on March 13, 2023. * Viral infection may not be the main reason for the admission.

Note: Because of the delay in date tested result, it affects the total number of COVID 19 admissions for a particular day. This lag in data impacts mostly the last couple of days from the day the report is updated. Includes lab or point of care positive for influenza, RSV, other respiratory viruses, four days prior to date of admission AND/OR at any point during admission. Episode of Care considers patients total movement within the health system related to their condition. It combines 2 or more admissions from 2 or more different facilities, if they are transferred (no break in care). Transfer = admission to any other hospital within 24 hours of discharge from previous hospital. Co-infected cases = positive for Influenza and RSV, or, Influenza and ‘other’, or, RSV and ‘other’, or, COVID-19 and Influenza, or, COVID-19 and RSV, or, COVID-19 and ‘other’.


Figure 6: The number of COVID-19, influenza, RSV, other respiratory viruses and co-infected cases admitted to hospital by age group, February 26 – March 11, 2023*



Data source: Digital Health Analytics, Saskatchewan Health Authority, Episode of Care methodology (Admission, Discharge, Transfer Database (ADT, RPPL, Panorama); data extracted on March 13, 2023. * Viral infection may not be the main reason for the admission

Note: Because of the delay in date tested result, it affects the total number of COVID 19 admissions for a particular day. This lag in data impacts mostly the last couple of days from the day the report is updated. Includes lab or point of care positive for influenza, RSV, other respiratory viruses, four days prior to date of admission AND/OR at any point during admission. Episode of Care considers patients total movement within the health system related to their condition. It combines 2 or more admissions from 2 or more different facilities, if they are transferred (no break in care). Transfer = admission to any other hospital within 24 hours of discharge from previous hospital. Co-infected cases = positive for Influenza and RSV, or, Influenza and ‘other’, or, RSV and ‘other’, or, COVID-19 and Influenza, or, COVID-19 and RSV, or, COVID-19 and ‘other’.

Figure 7: COVID-19 deaths by age group and week, September 4, 2021 – March 11, 2023*



Data Source: Panorama March 13, 2023

In the past two weeks, Feb 26 to Mar 11, there have been 14 deaths in COVID-19 cases, one in 0 to 19 years age-group, two in 20 to 59 years age-group and 11 in the 60 years or older group.

* Total COVID-19 deaths from Mar 2020 to date; n=1,913

Table 5: Community Respiratory Infection Surveillance Program Indicators by zone, March 05 - 11, 2023

Location	Test positivity – COVID-19 ¹ (positive lab tests)	Test positivity – Influenza (positive lab tests)	RLI* visits to EDs per 1,000 ²	RLI* 811 calls per 1,000 ³	School illness absenteeism >=10% ⁴	Wastewater indicator [†]	Proportion of population up-to-date vaccination – COVID-19 ⁵	Proportion of population immunized for Influenza (February 28, 2023) ⁶
Far North West (Meadow Lake and area)	9.8% (5)	6.7% (2)	No data	-	2.5%	-	24.3%	11%
Far North Central	16.7% (1)	0.0%	No data	-	0.0%	-	14.5%	4%
Far North East (La Ronge and area)	12.2% (5)	4.2% (1)	No data	-	8.7%	-	25.1%	9%
North West (North Battleford/ Lloydminster and area)	9.8% (13)	1.3% (1)	1.4	53.1	12.5%	Moderate	35.6%	19%
North Central (Prince Albert and area)	6.7% (8)	0.0%	No data	-	11.2%	Moderate	40.7%	22%
North East (Melfort and area)	5.4% (4)	0.0%	47.6	29.8	10.4%	-	44.0%	26%
Saskatoon	8.9% (51)	0.6% (1)	9.0	64.4	11.7%	Moderate	49.3%	29%
Central West (Kindersley and area)	8.5% (5)	4.0% (1)	132.8	-	9.2%	-	44.0%	29%
Central East (Yorkton/Melville and area)	16.5% (30)	0.0%	No data	-	10.3%	Moderate	45.9%	25%
Regina	11.0% (32)	0.5% (1)	No data	49.5	11.5%	Moderate-high	52.0%	28%
South West (Swift Current/Maple Creek and area)	23.1% (12)	0.0%	55.6	61.0	10.6%	Moderate	41.0%	27%
South Central (Moose Jaw and area)	9.1% (7)	0.0%	No data	-	12.4%	Moderate	44.4%	27%
South East (Weyburn/Estevan and area)	14.6% (14)	0.0%	75.9	56.2	12.7%	-	39.5%	24%
Unknown/Out of Province	4.3% (50)	0.0%	No data	-	7.3%	-	-	-
SASKATCHEWAN	8.1% (237)	0.9% (7)	15.1	54.0	11.2%	-	46.2%	27%

Notes: ¹by week of lab detection; effective Oct 30, 2022 includes cases who tested positive more than once >= 90 days apart; ²For COVID-19 test positivity, all tests reported were performed within the province. ³Based on reports from six of thirteen reporting areas; *Respiratory-like illness; ⁴811 data available at the six Integrated Service Areas geographical level; ⁵Unknown represents the number of students who were absent from the class due to illness with no known geography for the school. School absenteeism is the proportion of scheduled children who were absent from the class due to illness. The type of illness is not specified.; [†]SK overall estimate is currently unavailable as this metric tends to overestimate and underestimate WW level due to varied patterns across regions, which is difficult to synchronize with the population size of each region.; ⁶Up-to-date = completed a primary series and at least one additional booster, age 5+ years; ⁷received a vaccination within the current influenza season, age 6 months+; Does not include doses administered through NITHA or FNIHB, therefore some zones may underestimate coverage. This column will be updated at month end. ⁸Includes positive tests with pending locations. Numbers in parenthesis represents positive lab tests.

Technical Notes

1. **Laboratory surveillance:** Conducted through epidemiological analyses and Laboratory surveillance: Conducted through epidemiological analyses and positivity rate monitoring (counts and proportion of positive specimens, week of specimen collection, age category, geographical area, etiological type where applicable) of selected respiratory specimens submitted to the provincial laboratory in SK. Whole genome sequencing is conducted to detect changes (emergence of sub-lineages, variant proportion, etc) of clinical and public health importance among circulating respiratory organisms.
2. **Sentinel health providers:** Comprise a geographical-based network in practices across the province (n = 13 zones) who submit one to two specimens weekly to the Virology Section of the Roy Romanow Provincial Laboratory (RRPL), Saskatchewan Health Authority, from patients presenting with respiratory-like symptoms. Specimens are tested for a wider complement of respiratory organisms to monitor respiratory illness activity in the community. Assessment of co-infection (infected by more than one respiratory virus organism concurrently) occurs through sentinel provider submissions.
3. **Wastewater data:** Provided by the University of Saskatchewan and University of Regina Wastewater Team. Viral load for each zone was used to determine risk level (Low, Medium, Medium-High, High) using a four-bin system based on 100% of early Omicron peak reported. Locations sampled, includes: Saskatoon, Regina, Lumsden, North Battleford, Prince Albert, Yorkton, Swift Current, Moose Jaw, and Weyburn
4. **Data collection from Emergency Departments (ED):** Monitoring is done for a twenty-four hour period on at least one-week day (the exact time vary with the ED schedule). The ED reports to local public health services in their area on Wednesday afternoon and public health report to the Ministry of Health on Thursday each week. The count of Respiratory Like Illness (RLI) patients as a proportion of total ED admissions is captured.
5. **Reporting ED surveillance information:** Because there is no centralized data capture source for ED admissions in the province, each health area sets up a mechanism for EDs to report to public health services. Public health aggregates raw data from their EDs on the prescribed data collection form and sends it to the Ministry of Health for overall provincial monitoring. FNIHB and NITHA will report to the local zone which the ED or health centre is located. This does not preclude monitoring in First Nations health care facilities.
6. **HealthLine 811 callers with Respiratory Symptoms (RLI):** This count of response protocols collected by HealthLine nurses specific to callers reporting respiratory-like symptoms. HealthLine data is collected for a seven day week, Monday to Sunday. Data is transformed into the rate of callers with respiratory symptoms from each Integrated Service Area (ISA) per 1000 calls from that ISA concerning any type of symptom.
7. **A confirmed outbreak:** Defined as two or more lab confirmed respiratory virus cases in high-risk settings where transmission is evident or there is a high level of suspicion of transmission. Outbreaks are reported by the week they were reported to the local public health office and not necessarily in the week that the outbreak began. For this report outbreaks in high risk settings comprise long term care facilities, personal care homes and group homes
8. **COVID hospitalized admissions** is the number of C-19 positive cases that during the surveillance week were admitted as an inpatient to an acute care facility. This includes patients with C-19 related illness, incidental COVID infection, and patients under investigation. COVID ICU admissions is the number of C-19 positive cases that during the surveillance week were admitted to an ICU location in SK. This includes both infectious and non-infectious cases.
9. **Influenza, RSV and other respiratory virus admissions:** Delays in testing results affect the total number of Influenza, RSV and other respiratory virus admissions for a particular day. This lag in data has the greatest impact on the two days prior to when the report is updated. Counts include individuals who are laboratory positive for influenza, RSV, and other respiratory viruses, within four days prior to date of admission AND/OR at any point during the hospital stay. Episode of Care considers patients' total movement within the health system related to their condition. It combines 2 or more admissions from 2 or more different facilities, if they are transfers (i.e., no break in care). Transfer: Admission to any other hospital within 24 hours of discharge from previous hospital. Co-infected Cases = if positive for Influenza and RSV or, positive for Influenza and Other Respiratory viruses or, positive for RSV and Other respiratory viruses or, positive for Covid-19 and Influenza or, positive for Covid-19 and RSV or, positive for Covid-19 and Other Respiratory viruses.
10. **Variant of concern (VOC):** VOCs are SARS-CoV-2 viruses that have undergone genetic modification or mutation causing altered virus infectivity, replication and pathogenicity. As a result it can alter host immune response. The Roy Romanow Provincial Laboratory (RRPL) tests for and monitors COVID-19 variants of concern (VOCs) in Saskatchewan. Confirmation of VOC lineages is done by conducting whole genome sequencing (WGS) at RRPL or the National Microbiology Laboratory, one to two weeks to complete WGS. Data sources for VOCs analysis include testing data from the RRPL, and epidemiological information from Panorama. Where geographical zone is missing in RRPL or Panorama data, the Saskatchewan postal code file is used to identify cases' geographical information.
11. **VOC Lineages BA.2, BA.2.75, CH.1.1, BA.5, BF.7, BQ.1, BQ.1.1, XBB, XBB.1.5, and XBF** are all classified under the WHO Label of "Omicron". Omicron lineages BA.2.75, CH.1.1, BF.7, BQ.1, BQ.1.1, XBB, XBB.1.5, and XBF are emerging global variants that RRPL continues to monitor. Lineages that are not explicitly listed are aggregated under their corresponding parent lineage (e.g., BA.1, BA.2, BA.4, or BA.5). Previously, CH.1.1 was aggregated with BA.2.75; BA.2.75 was aggregated with BA.2; BF.7, BQ.1, and BQ.1.1 were aggregated with BA.5. Percentages are shown when a lineage constitutes 5% or more of total specimens evaluated for a given surveillance week.
12. **COVID-19 cases:** Effective September 4, 2022, COVID-19 cases are based on lab detection and include cases who tested positive more than once 90 days, or further, apart. Prior to this, cases include, only, first time cases reported and entered into Panorama.
13. **COVID-19 Deaths:** Includes deaths entered into Panorama IOM among laboratory confirmed cases. Deaths are reported based on the actual date of death. Deaths in previous periods may be adjusted from previous reports due to data lag
14. **COVID-19 Immunizations:** Up-to-date (UTD) COVID-19 vaccination is the proportion of people having completed a primary series and one booster for ages five and older divided by the eligible population found in the Saskatchewan Covered Population, 12-Nov-2022 Ministry of Health version (2022 Version 2). Though vaccinated children six months to four years of age may be technically UTD, this specific definition does not apply to them. In addition, UTD in last six months is calculated by the proportion of people having received one or more boosters within the previous six months.
15. **Influenza immunizations:** UTD Influenza vaccination is the proportion of people, six months and older, having one influenza dose this season divided by the eligible population found in the Saskatchewan Covered Population, 12-Nov-2022 Ministry of Health version (2022 Version 2). Vaccination for the current influenza season officially began October 11, 2022. Some doses were administered prior to the start date.
16. **Staffed Inpatient beds:** Weekly average COVID Occupancy is a 7-Day average percentage of acute inpatient beds staffed and in operation COVID positive patients occupy. The full calculation of this metric is: $\text{Average COVID occupancy} = \frac{\sum(8\text{am covid census}) + \sum(8\text{am beds staffed and in operation})}{\sum(\text{Planned beds} + \text{Surge Beds} - \text{Closed})} \times 100\%$. Where "bed staffed and in operation" = "Planned beds" + "Surge Beds" - "Closed" and $\sum(\dots)$ indicates summation over 7-day period from Sunday to Saturday. 8am COVID census is taken from the ADT patient registration, which is fed to the provincial data-mart and archived hourly. 8am planned bed, surge beds, and closed beds is compiled via data feeds from APF (Saskatoon & Regina) and the provincial bed edits interface (INH & IRH).
17. **Rate of COVID-19 hospitalization (ICU or Death)** were calculated by summing the daily number of hospitalizations (ICUs or Deaths) for the period by vaccine status (numerator) divided by the mid period population by respective vaccine status (denominator), multiplied by 100,000. This estimate is further divided by the number of days to obtain the daily rate. Denominator for individuals in the Booster in the past 6-months group are all Saskatchewan residents who have had their booster dose within the last 6 months. To eliminate bias of age all rates are adjusted by age. Direct standardization method is employed using the Saskatchewan population as the standard population. Age at first dose used in the rate calculation. Individuals with unknown age are excluded from age-specific analyses. Estimates of relative risk (i.e. rate ratios) is obtained by comparing vaccinated with 2 doses (Any Booster dose) and unvaccinated. Risk estimates may differ from other reports due to differing methodologies.