

Notice of Immunization

Important: Keep this form with the student's immunization records or other health documents.

| _____ was immunized _____ with the vaccine(s) noted below: Date | | Arm | |
|---|---|------|-------|
| <input type="checkbox"/> | Diphtheria, Tetanus, Pertussis, Polio, <i>Haemophilus influenzae</i> type b | Left | Right |
| <input type="checkbox"/> | <i>Haemophilus influenzae</i> type b | Left | Right |
| <input type="checkbox"/> | Hepatitis A | Left | Right |
| <input type="checkbox"/> | Hepatitis B | Left | Right |
| <input type="checkbox"/> | Human Papillomavirus (9 HPV types) | Left | Right |
| <input type="checkbox"/> | Measles, Mumps, Rubella | Left | Right |
| <input type="checkbox"/> | Measles, Mumps, Rubella, Varicella | Left | Right |
| <input type="checkbox"/> | Meningococcal B | Left | Right |
| <input type="checkbox"/> | Meningococcal Conjugate C | Left | Right |
| <input type="checkbox"/> | Meningococcal Conjugate ACYW-135 | Left | Right |
| <input type="checkbox"/> | Pneumococcal Conjugate 13 | Left | Right |
| <input type="checkbox"/> | Pneumococcal Polysaccharide 23 | Left | Right |
| <input type="checkbox"/> | Polio | Left | Right |
| <input type="checkbox"/> | Tetanus, Diphtheria, Pertussis | Left | Right |
| <input type="checkbox"/> | Tetanus, Diphtheria, Pertussis, Polio | Left | Right |
| <input type="checkbox"/> | Varicella | Left | Right |
| <input type="checkbox"/> | Other: | | |

Care after immunization:

- Children can use their arm normally afterwards.
- Common side effects like redness, pain and tenderness at the injection site, a fever, or feeling tired are generally mild and usually last for 1 – 2 days. For redness or swelling at the injection site, apply a cold compress.
- For specific vaccine side effects, refer to the vaccine fact sheets that were included with the consent form or go to: www.saskatchewan.ca/immunize
- For information about immunization programs, call HealthLine at 811 or go to: www.saskatchewan.ca/immunize
- **Please report unexpected side effects to a public health nurse, a physician or nurse practitioner.**

Acetaminophen (Tylenol, Tempra) may be given for fever or soreness. ASA (Aspirin) must NOT be given to anyone younger than 18 years old because of the risk of Reye syndrome.

July 2017 PHNF 401