

Guide to Crop Protection 2019

For Reference Until December 31, 2019
This publication is updated annually and replaces
the 2018 and previous issues.

Introduction	7
How to Use This Book	7
Product Labels and PCP Numbers.....	7
Safe Use of Herbicides, Fungicides and Insecticides	7
Protecting Yourself from Exposure to Herbicides, Fungicides and Insecticides.....	8
Protective Clothing.....	9
Protecting Your Eyes, Face and Lungs.....	9
Understanding Maximum Residue Limit Statements in the Guide	10
Avoiding Spray Drift	10
Herbicide Efficacy with Low-drift Nozzles.....	11
Handling a Drift Complaint.....	11
Mixing Pesticides	11
Container Disposal.....	12
Sprayer Cleaning	12
How to Identify Crop and Weed Leaf Stages	13
Trade Names, Active Ingredients and Formulations.....	16
Key to Product Pages	34
Weed Control.....	37
Integrated Weed Management	37
Making Spray Decisions.....	37
Weed Resistance to Herbicides	45
Adjuvants and Your Herbicide	52
Crop and Herbicide Recommendation Tables	54
Special Weed Problems	72
Soil Residual Herbicides.....	78
Effect of Rainfall on Herbicide Efficacy	81
Products Available as Prepackaged Tank Mixes	82
Herbicide Product Pages	83
Plant Disease Control.....	423
Integrated Plant Disease Management.....	423
Effects of Weather	426
Resistance Management	426
Fungicide Modes of Action.....	426
Foliar Fungicide Tables	429
Foliar Fungicide Product Pages	436
Seed Treatment Tables	524
Seed Treatment Product Pages	531

Insect Control	596
Additional Resources	596
Insect Management Decisions	596
Preharvest Interval	596
Field Scouting	596
Economic Thresholds	596
Estimating Percent Defoliation	596
Hazard Ratings of Insecticides to Bees	597
Reducing Bee Losses from Insecticides	598
Insecticide Poisoning in Humans	598
Resistance of Insects to Insecticides	598
Insecticide Groups	599
Field Scouting and Insect Management Charts (for individual crops)	600
Insect Control in Stored Grain	636
Insecticide Product Pages	639
Pesticide Index	3
Herbicide Directory	3
Foliar Fungicide Directory	5
Seed Treatment Directory	6
Insecticide Directory	6
Crop Protection Companies	Inside Cover
Emergency Numbers	Inside Cover
Agricultural Offices	Inside Cover

Table 1: Metric Conversion Factors* (Approximate)

Metric Unit	Metric to Imperial	Imperial Unit	Imperial to Metric	Metric Unit
Linear centimetre (cm)	x 0.39	Linear inch	x 2.54	Linear centimetre (cm)
Area square metre (m ²) hectare (ha)	x 1.2 x 2.5	Area square yard acres	x 0.84 x 0.4	Area square metre (m ²) hectare (ha)
Volume litre (L)	x 0.22	Volume gallon	x 4.55	Volume litre (L)
Pressure kilopascals (kPa)	x 0.14	Pressure psi	x 6.9	Pressure kilopascals (kPa)
Weight gram (g) kilogram (kg)	x 0.04 x 2.2	Weight ounce (oz) pound (lb)	x 28.35 x 0.454	Weight gram (g) kilogram (kg)
Agricultural litres per hectare (L/ha) litres per hectare (L/ha) litres per hectare (L/ha) millilitres per hectare (mL/ha) kilograms per hectare (kg/ha) grams per hectare (g/ha)	x 0.089 x 0.357 x 0.71 x 0.014 x 0.89 x 0.014	Agricultural gallons/acre quarts/acre pints/acre fluid ounces (fl. oz)/acre pounds (lb)/acre ounces (oz)/acre	x 11.23 x 2.81 x 1.41 x 70.22 x 1.12 x 70	Agricultural litres per hectare (L/ha) litres per hectare (L/ha) litres per hectare (L/ha) millilitres per hectare (mL/ha) kilograms per hectare (kg/ha) grams per hectare (g/ha)

*EXAMPLE: To convert centimetres to inches, multiply by 0.39; conversely, to convert inches to centimetres, multiply by 2.54.

CAUTION: Herbicide labels are in metric units only. Conversion between the Metric and Imperial system may result in confusion. It is recommended to use metric units only.

Pesticide Index

Herbicide Directory

2,4-D.....	83	Cobutox 625 (see 2,4-DB).....	86
2,4-DB.....	86	Command 360 ME.....	160
AAtrex Liquid.....	89	Conquer.....	161
Accent 75DF/Accent Grande.....	91	Contender (see Quizalofop).....	340
Accurate (see Metsulfuron).....	295	Cordon (see Fenoxaprop).....	195
Aim EC.....	93	Craven (see Diquat).....	173
Ally (see Metsulfuron).....	295	Credit 45 (see Glyphosate).....	233
Altitude FX2.....	95	Credit Xtreme (see Glyphosate).....	233
Amitrol 240.....	97	Crush'R 540 (see Glyphosate).....	233
Antler (see Clethodim).....	144	Crush'R Plus (see Glyphosate).....	233
Approve (see Bromoxynil/2,4-D Ester).....	130	Curtail M (see Clopyralid/MCPA).....	154
Ares.....	99	Davai 80SL (see Imazamox).....	260
Armezon (see Topramezone).....	378	Deploy WDG (see Thifensulfuron/tribenuron).....	373
Armory (see Diquat).....	173	Desica (see Diquat).....	173
Arrow (see Clethodim).....	144	Dicamba.....	163
Arrow-All-In (see Clethodim).....	144	Dicamba/Mecoprop/MCPA.....	168
Assert 300SC (see Imazamethabenz).....	257	Dichlorprop/2,4-D.....	170
Assure II (see Quizalofop).....	340	Dichlorprop-DX (see Dichlorprop/2,4-D).....	170
Attain XC (see Fluroxypyr + 2,4-D).....	220	Diquat.....	173
Aurora (see Clodinafop).....	148	Diquat 240 (see Diquat).....	173
Authority 480.....	101	Distinct.....	176
Authority Charge.....	103	DriFast (see Diquat).....	173
Authority Supreme.....	103	Dual II Magnum.....	178
Avadex Brands.....	106	Duet (see Imazamox/Imazethapyr).....	262
Avert (see Imazamethabenz).....	257	DyVel.....	179
Axial.....	109	DyVel DSp.....	181
Axial iPak.....	111	Eclipse Brands (see Glyphosate + Clopyralid).....	242
Axial Xtreme.....	112	Edge Granular.....	184
Axial Xtreme iPak.....	114	Embutox (see 2,4-DB).....	86
Badge II (see Bromoxynil/MCPA ester).....	132	Enforcer D.....	186
Banvel II (see Dicamba).....	163	Enforcer M (see Bromoxynil + MCPA + Fluroxypyr).....	136
Banvel VM (see Dicamba).....	163	Enforcer MSU.....	188
Barricade II.....	114	Engenia (see Dicamba).....	163
Basagran Brands (see Bentazon).....	117	Eptam Liquid EC.....	189
Bengal WB (see Fenoxaprop).....	195	Escort.....	191
Bentazon.....	117	Etaprop XT (see Dichlorprop/2,4-D).....	170
Benta Super (see Bentazon).....	117	Esteem (see Clopyralid/MCPA + Fluroxypyr).....	156
Bison (see Tralkoxydim).....	382	Everest 2.0/Everest 3.0 (see Flucarbazone).....	214
BlackHawk (with carfentrazone).....	121	Express FX.....	193
BlackHawk (with pyraflufen-ethyl).....	122	Express FX (co-pack).....	195
Blitz (see Florasulam).....	203	Express Pro (see Tribenuron/Metsulfuron).....	394
Bolster (see Diquat).....	173	Express SG (see Tribenuron).....	390
Bonanza (see Trifluralin).....	396	Facet L (see Quinclorac).....	337
Boost (see Thifensulfuron/tribenuron).....	373	Factor 540 (see Glyphosate).....	233
Broadband.....	124	Fenoxaprop.....	195
BroadSide (see Thifensulfuron/tribenuron+MCPA ester).....	377	FeXapan (see Dicamba).....	163
Bromotril II (see Bromoxynil).....	126	Fierce.....	198
Bromoxynil.....	126	FirstPass (see Florasulam).....	203
Bromoxynil/2,4-D Ester.....	130	Flexstar GT.....	201
Bromoxynil/MCPA ester.....	132	Florasulam.....	203
Bromoxynil + MCPA + Fluroxypyr.....	136	Florasulam + 2,4-D.....	205
Brotex 240/Brotex 480 (see Bromoxynil).....	126	Florasulam + Clopyralid + MCPA.....	207
Buctril M (see Bromoxynil/MCPA ester).....	132	Florasulam + glyphosate.....	208
Buzzin (see Metribuzin).....	291	Florasulam + MCPA Ester.....	210
Cadillac (see Clodinafop).....	148	Florasulam/fluroxypyr + MCPA ester.....	212
Cadillac One (see Clodinafop).....	148	Flucarbazone.....	214
Caliber 625 (see 2,4-DB).....	86	Flurox-24 (see Fluroxypyr + 2,4-D).....	220
Casoron.....	139	Fluroxypyr.....	217
Centurion (see Clethodim).....	144	Fluroxypyr + 2,4-D.....	220
Chateau.....	410	Fluroxypyr + MCPA.....	222
Cirpreme/Cirpreme XC.....	140	Foax (see Clodinafop).....	148
CleanStart.....	142	Focus.....	224
ClearOut 41 Plus (see Glyphosate).....	233	Foothills NG (see Clodinafop).....	148
Clethodim.....	144	ForceFighter M (see Bromoxynil + MCPA + Fluroxypyr).....	136
Clethodim 250 (see Clethodim).....	144	Fortress MicroActiv.....	226
Clever (see Quinclorac).....	317	Frontier Max.....	228
Clodinafop.....	148	Frontline 2,4-D (see Florasulam + 2,4-D).....	205
Clopyralid.....	151	Frontline XL (see Florasulam + MCPA Ester).....	210
Clopyralid/MCPA.....	154	Gladiator (see Imazethapyr).....	265
Clopyralid/MCPA + fluroxypyr.....	156	Glykamba.....	243
Clovitox Plus (see MCPB / MCPA).....	288	Glufosinate 150SN.....	230

Glyphosate.....	233	MultiStar (see Imazethapyr).....	265
Glyphosate + Clopyralid	242	Muster Toss-N-Go	301
GoldWing	247	Navius.....	303
Gramoxone	228	NextStep NG (see Clodinafop).....	148
Grazon XC.....	247	Nimble (see Thifensulfuron/tribenuron).....	373
Guardsman Diquat (see Diquat)	173	Nuance (see Tribenuron).....	390
Guardsman Glyphosate (see Glyphosate)	233	Nuance Pro (see Tribenuron/Metsulfuron).....	394
Heat Brands	249	Nufarm Tralkoxydim Liquid (see Tralkoxydim).....	382
Heat Complete	253	OctTain XL	305
MPower HellCat (see Fenoxaprop).....	195	Odyssey/Odyssey NXT (see Imazamox/Imazethapyr).....	262
Horizon NG (see Clodinafop).....	148	Odyssey Ultra/Odyssey Ultra NXT	308
Hotshot	255	Olympus	309
Ikwin (see Fluroxypyr).....	217	Optica Trio.....	310
Imazamethabenz.....	257	Option 2.25 OD	312
Imazamox.....	260	Oracle (see Dicamba)	163
Imazamox/Imazethapyr	262	Outlook	314
Imazethapyr	265	Outshine (see Florasulam/fluroxypyr + MCPA ester).....	212
Impact (see Topramezone)	378	Overdrive.....	316
Inferno Duo	267	Paradigm	317
Inferno WDG (see Tribenuron).....	390	Pardner (see Bromoxynil)	126
Infinity	269	Patron 240 EC (see Clethodim).....	144
Infinity FX	271	Permit WG	320
Ingenious (see Quinclorac)	337	Phantom (see Imazethapyr)	265
Kerb	272	Pinnacle SG Toss-N-Go.....	323
Koril 235 (see Bromoxynil)	126	Pixxaro.....	325
Korrex II	274	Poast Ultra.....	327
Ladder (see Clodinafop).....	148	Predicade.....	330
Ladder All-In-One (see Clodinafop)	148	PrePass Flex (see Florasulam).....	203
Leader (see Bromoxynil/2,4-D Ester)	130	PrePass XC (see Florasulam + glyphosate).....	208
Liberty 150SN (see Glufosinate 150SN).....	230	Prestige Brands (see Clopyralid/MCPA + Fluroxypyr).....	156
Liberty 200SN	277	Primextra II Magnum	331
Linuron.....	279	Princep Nine-T (see Simazine)	363
Linuron 400 (see Linuron)	279	Priority (see Florasulam + glyphosate)	208
Liquid Achieve (see Tralkoxydim).....	382	Prism SG.....	333
Logic M (see Bromoxynil/MCPA ester).....	132	Pulsar	335
Lontrel 360/Lontrel XC (see Clopyralid).....	151	Puma Advance (see Fenoxaprop).....	195
Lorox L (see Linuron).....	279	Pursuit 240 (see Imazethapyr)	265
Loveland Bromax (see Bromoxynil).....	126	Pyralid (see Clopyralid).....	151
Luxxur	281	Quinclorac.....	337
Manipulator 620	283	Quizalofop	340
Marengo (see Tralkoxydim).....	382	Reclaim II	343
MasterLine Quinclorac (see Quinclorac)	337	Refine M (see Thifensulfuron/tribenuron + MCPA ester)	377
Matrix (see Glyphosate)	233	Refine SG (see Thifensulfuron/tribenuron)	373
MCPA.....	284	Reflex	345
MCPB / MCPA	288	Reglone (see Diquat).....	173
Mecoprop-p	290	Reglone Ion (see Diquat)	173
Metribuzin.....	291	Restore II	347
Metrix SC (see Metribuzin).....	291	Retain SG	350
Metsulfuron	295	Reward	351
Mextrol 450 (see Bromoxynil/MCPA ester).....	132	Rexade.....	352
Mizuna (see Imazamox)	260	Rezuvant.....	354
Momentum.....	297	Rival (see Trifluralin).....	396
MPower 2,4-D (see 2,4-D)	83	Roundup brand products (Transorb HC; WeatherMax) (see Glyphosate)	233
MPower Battlefront (see Florasulam).....	203	Roundup Xtend	355
MPower Battlefront 2,4-D (see Florasulam + 2,4-D).....	205	R/T 540 (see Glyphosate)	233
MPower Battlefront CM (see Florasulam + Clopyralid + MCPA)	207	Rush 24 (see Fluroxypyr + 2,4-D).....	220
MPower Battlefront M (see Florasulam + MCPA)	210	Rush M (see Fluroxypyr + MCPA)	222
MPower Boa (see Bentazon).....	117	Salute	358
MPower Bromoxynil (see Bromoxynil).....	126	Select (see Clethodim).....	144
MPower Buck M (see Bromoxynil/MCPA ester).....	132	Sencor 75 DF (see Metribuzin).....	291
MPower Clobber (see Clopyralid)	151	Shadow RTM (see Clethodim).....	144
MPower Clobber G (see Glyphosate + Clopyralid)	242	Sharda Glyphosate (see Glyphosate)	233
MPower Disruptor 360/540 (see Glyphosate)	233	Sharpshooter (see Glyphosate).....	233
MPower Foxy (see Fluroxypyr).....	217	Sharpshooter Plus (see Glyphosate).....	233
MPower Foxy CM (see Clopyralid/MCPA + Fluroxypyr)	156	Shieldex 400SC.....	259
MPower Foxy M (see Fluroxypyr + MCPA).....	222	Sierra 2.0/Sierra 3.0 (see Flucarbazone).....	214
MPower Foxy Pro (see Fluroxypyr + 2,4-D)	220	Signal (see Clodinafop)	148
MPower Good Harvest	299	Signal FSU	361
MPower Kamikaze (see Imazethapyr).....	265	Simazine	363
MPower Kickoff (see Florasulam + Glyphosate)	208	Simazine 480 (see Simazine)	363
MPower Independence (see Clethodim).....	144	Simplicity.....	365
MPower Ninja (see Imazamox/Imazethapyr)	262	Slam'R (see Clodinafop)	148
MPower R (see Thifensulfuron/tribenuron)	373	Smoke (see Glyphosate)	233
MPower Vigor (see Glufosinate 150SN).....	230	Solo/Solo ADV (see Imazamox).....	260
MPower X (see Tribenuron).....	390		

Solo Ultra.....	368	Treflan (see Trifluralin).....	396
Sortan IS.....	369	Tribenuron	390
Spike (see Tribenuron).....	390	Tribenuron + 2,4-D.....	393
Spectrum (see Florasulam + Clopyralid + MCPA)	207	Tribenuron/Metsulfuron	394
Squadron (see Metribuzin)	291	TriCor 75 DF (see Metribuzin).....	291
Stage (see Diquat).....	173	Trifluralin.....	396
StartUp (see Glyphosate).....	233	Triton C.....	401
Stellar/Stellar XL (see Florasulam/fluroxypyr + MCPA ester)	212	Triton K.....	403
Stonewall (see Glyphosate)	233	Trophy (see fluroxypyr+MCPA).....	222
Sword (see Dicamba/Mecoprop/MCPA)	168	Tropotox Plus (see MCPB / MCPA)	288
Tandem	371	Tundra	405
Target (see Dicamba/Mecoprop/MCPA)	168	Turboprop (see Dichlorprop/2,4-D)	170
Tensile.....	372	Ultim 75DF/Ultim Grande	408
Thifensulfuron/tribenuron.....	373	Ultra Blazer.....	410
Thifensulfuron/tribenuron + MCPA ester	377	Valtera.....	411
Thrasher II (see Bromoxynil/2,4-D Ester)	130	Varro.....	414
Thumper (see Bromoxynil/2,4-D Ester)	130	Vector/Vector 540 (see Glyphosate)	233
Topline (see Florasulam + MCPA Ester).....	210	Velocity m3.....	417
Topramezone.....	378	Velpar DF CU	419
Topside (see MCPB / MCPA)	288	Vigil WB (see Fenoxaprop).....	195
Tordon 22K.....	380	Viper ADV.....	421
Tracker XP (see Dicamba/Mecoprop/MCPA)	168	VP-480 (see Glyphosate)	233
Tralkoxydim	382	WildCat/WildCat Enhanced (see Fenoxaprop).....	195
Travallas	385	Xtendimax (see Dicamba)	163
Traxos.....	387	Yuma GL (see Quizalofop).....	340
TraxosTwo.....	389		

Foliar Fungicide Directory

Acapela.....	436	Manzate Pro-Stick (see Mancozeb)	477
Allegro 500F	439	Nexicor	478
Aprovia Top.....	440	Nufarm Propiconazole Fungicide (see Propiconazole).....	489
Azoshy (see Azoxystrobin).....	441	Orondis Ultra.....	479
Azoxystrobin	441	Overall 240 SC (see Iprodione)	469
Bravo 500 (see Chlorothalonil)	447	Palliser (see Tebuconazole)	515
Bravo Zn(see Chlorothalonil)	447	Parasol WG (see Copper).....	451
Bumper 432 EC (see Propiconazole).....	489	Penncozeb 75 DF (see Mancozeb)	477
Cabrio Plus.....	443	Phosphorous acid.....	480
Cantus WDG Fungicide.....	444	Phostrol.....	481
Caramba	445	Pivot 418 EC (see Propiconazole).....	489
Chlorothalonil.....	447	Polyram DF.....	483
Confine Extra (see Phosphorous acid)	480	Priaxor.....	484
Contans WG.....	449	Prodex SC (see Iprodione).....	469
Copper.....	451	Proline 480 SC.....	486
Copper 53W (see Copper)	451	Propel (see Propiconazole)	489
Copper Spray (see Copper).....	451	Propi Super 25 EC (see Propiconazole).....	489
Cosavet DF Edge.....	452	Propiconazole	489
Cotegra.....	453	Propulse.....	492
Cueva (see Copper).....	451	Prosaro 250 EC/Prosaro XTR	493
Curzate 60 DF.....	454	Pyraclostrobin.....	495
Delaro 325 SC	455	Quadris (see Azoxystrobin).....	441
Dithane Rainshield (see Mancozeb)	477	Quadris Top.....	498
Double Nickel 55/Double Nickel LC.....	457	Quash.....	499
Dyax	459	Quilt.....	501
Echo 720 (see Chlorothalonil).....	447	Ranman 400SC	504
Echo 90DF (see Chlorothalonil).....	447	Rampart (see Phosphorous acid).....	480
Elatus.....	460	Reason 500SC	505
Elixir.....	462	Regalia Maxx	506
Evito 480	463	Revus.....	507
Fitness (see Propiconazole).....	489	Ridomil Gold Products.....	508
Folicur 250EW (see Tebuconazole).....	515	Rovral Flo (see Iprodione).....	469
Fontelis.....	465	Scala SC.....	509
Forum	466	Sercadis.....	510
Fullback 125SC	467	Serenade OPTI	511
Gavel 75 DF.....	468	Serenade Soil	513
Headline EC (see Lance AG).....	471	Tanos 50 DF	514
Hornet 432 F (see Tebuconazole)	515	Tebuconazole.....	515
Iprodione.....	469	Tilt 250E (see Propiconazole)	489
Kenja 400SC.....	470	Topnotch	517
Lance AG.....	471	Trivapro	518
Lance WDG	473	Twinline.....	520
LifeGard WG.....	474	Vertisan	521
Luna Tranquility	475	Zampro.....	523
Mancozeb.....	477		

Seed Treatment Directory

Agrox FL.....	531	NipsIt SUITE Cereals OF Seed Protectant.....	568
Alias 240 SC (see Imidacloprid).....	551	Phosphorous acid.....	569
Allegiance FL (see Metalaxyl).....	565	Poncho 600 FS (see Clothianidin).....	534
Apron Advance/Apron Maxx RTA + Vibrance500 FS.....	532	Potato ST 16 (see Mancozeb).....	561
Belmont 2.7 FS (see Metalaxyl).....	565	Prosper EverGol.....	570
Clothianidin.....	534	Rampart (see Phosphorous acid).....	569
Confine Extra (see Phosphorous acid).....	569	Rancona Pinnacle (see Ipconazole + Metalaxyl).....	558
Cruiser 5FS.....	535	Rancona V RS.....	572
Cruiser Maxx Corn.....	537	Rancona Trio.....	572
Cruiser Maxx Potato Extreme.....	538	Rascendo (see Visivio).....	559
Cruiser Maxx Vibrance Beans.....	539	Raxil MD.....	574
Cruiser Maxx Vibrance Pulses.....	540	Raxil PRO.....	575
Cruiser Vibrance Quattro.....	541	Raxil PRO Shield.....	576
Cyantraniliprole.....	543	Reason 500SC.....	577
Emesto Silver.....	545	Revus.....	578
EverGol Energy.....	546	Senator PSPT.....	579
Fortenza (see Cyantraniliprole).....	543	Serenade OPTI.....	580
Gaucho CS FL.....	547	Solan MZ (see Mancozeb).....	561
General Storage Disinfectant.....	548	Sombrero 600 FS (see Imidacloprid).....	551
Heads Up Plant Protectant.....	549	Stadium.....	581
Helix Vibrance.....	550	StorOx.....	582
Imidacloprid.....	551	Stress Shield 600 (see Imidacloprid).....	551
Insure Cereal.....	553	Titan (see Clothianidin).....	534
Insure Cereal FX4.....	554	Titan Emesto.....	583
Insure Pulse.....	555	Trilex EverGol.....	583
INTEGO Solo Fungicide.....	557	Trilex EverGol Shield.....	585
Ipconazole + Metalaxyl.....	558	Tuberseal (see Mancozeb).....	561
Lumiderm (see Cyantraniliprole).....	543	Verimark (see Cyantraniliprole).....	543
Lumisena.....	559	Vibrance 500FS.....	586
Mancozeb.....	561	Vibrance Maxx RFC/RTA.....	587
Maxim D/Maxim PSP/Maxim MZ PSP.....	562	Vibrance Maxx RFC with INTEGO Seed Treatment.....	589
Maxim Quattro.....	563	Vibrance Quattro.....	590
Mertect SC.....	564	Vibrance Ultra Potato.....	591
Metalaxyl.....	565	Visivio.....	592
Metlock CT.....	567	VitaFlo Brands.....	594
NipsIt INSIDE 600 Insecticide (see Clothianidin).....	534		

Insecticide Directory

Actara.....	639	Mako Insecticide (see Cypermethrin).....	650
Admire 240/Admire SPT (see Imidacloprid).....	663	Malathion.....	668
Agri-Mek SC.....	640	Matador (see Lambda-cyhalothrin).....	665
Alias 240 SC (see Imidacloprid).....	663	Minecto Duo 40WG.....	670
Ambush (see Permethrin).....	676	Minecto Pro.....	671
Assail.....	641	Movento 240 SC.....	672
Beleaf 50 SG.....	642	MPOWER Krypton (see Chlorpyrifos).....	644
Capture 240 EC.....	643	Nolo Bait.....	673
Chlorpyrifos.....	644	Nufos 4E (see Chlorpyrifos).....	644
Citadel 480EC (see Chlorpyrifos).....	644	Oberon.....	674
Closer.....	645	Orthene.....	675
Clutch.....	646	Perm-UP (see Permethrin).....	676
Concept.....	647	Permethrin.....	676
Coragen.....	649	Poleci 2.5 EC Western (see Deltamethrin).....	653
Cygon 480 EC/Cygon 480-Ag (see Dimethoate).....	656	Pounce (see Permethrin).....	676
Cypermethrin.....	650	Pyrifos 15G (see Chlorpyrifos).....	644
Decis 5 EC (see Deltamethrin).....	653	Pyrinex 480EC (see Chlorpyrifos).....	644
Delegate.....	652	Rimon 10 EC.....	677
Deltamethrin.....	653	Sefina.....	678
Dibrom.....	655	Sevin XLR.....	680
Dimethoate.....	656	Sharda chlorpyrifos 480 EC (see Chlorpyrifos).....	644
Dipel 2X DF.....	657	Ship 250 EC. (see Cypermethrin).....	650
Eco Bran.....	658	Silencer 120 EC (see Lambda-cyhalothrin).....	665
Exirel.....	659	Sivanto Prime.....	681
Entrust 80 W.....	660	Sluggo Professional.....	682
Fulfill.....	661	Success 480 SC.....	683
Imidacloprid.....	663	Superior 70 Oil.....	684
Imidan.....	664	Thimet 20G.....	685
Lagon 480E (see Dimethoate).....	656	UP-Cyde 2.5 EC (see Cypermethrin).....	650
Lambda-cyhalothrin.....	665	Voliam Xpress.....	687
Lannate.....	667	Warhawk 480EC (see Chlorpyrifos).....	644
Lorsban 4E/Lorsban NT (see Chlorpyrifos).....	644		

Introduction

How to Use This Book

This publication is only a guide. Always refer to the product label for application details and precautions. If the information in this publication differs from the label information, follow label instructions.

The *Guide to Crop Protection* is divided into five chapters: (1) Introduction; (2) Weed Control; (3) Foliar Fungicides; (4) Seed Treatments; and (5) Insect Control.

To use the information in each of these sections, use the following process:

1. Turn to the charts at the beginning of each section. There is a set of charts for weeds, plant diseases and insect control. Select the chart for the crop you want or plan to grow. Use the chart to match your weed, disease or insect problems with the products available for that crop.
2. Once you have narrowed your product choices down to a few candidates, go to the recommendation section for that product. Products are listed alphabetically. Read the recommendations thoroughly for each product you are considering.
3. Read the product label attached to the container for detailed instructions on application.

This publication is intended to be used as a guide only. Information contained herein is that available at time of printing.

While every effort has been made to ensure accuracy, the provincial government does not accept responsibility for label changes. When more than one trade name is listed, not all weeds or tank mixes may appear on all labels. Consult product labels attached to pesticide containers for final detailed instructions.

Certain recommendations in this publication are given in quantity of commercial product per acre (mL, L, g or kg/acre). Product labels are given in quantity of product per hectare (mL, L, g or kg/ha). To avoid application errors be sure to read and understand label recommendations.

The *Guide to Crop Protection* includes the most recent recommendations for weed, plant disease and insect control in field and forage crops. These recommendations are based on the uses registered under the Pest Management Regulatory Agency's *Pest Control Products Act*. It is an offence under *The Pest Control Products Act* to apply any chemical in a manner not consistent with the product label. If you have any doubts regarding the instructions in this publication, or on the product label, contact the company representative, your local agricultural office or the Pest Management Regulatory Agency for further advice.

Product Labels and PCP Numbers

On each Product Page you will see a Registration or PCP number, so named because it is mandated by the *Pest Control Products Act*. Under the Act, every pesticide requires a unique identifier – the product's Registration or PCP number. That number must also appear on the product's label.

The pesticide label packaged with the product is the authoritative source of information on use of the product and will contain more detailed information than is included in this Guide. Some products have a number of trade names for the same active ingredient. However, each product will have its own Registration (PCP) number and these appear next to the registrants' names. Users who are seeking more detailed information than is provided in this guide, prior to purchase, can use the Registration (PCP) number to access a sample product label online through the Pest Management Regulatory Agency's (PMRA) website or they can contact the PMRA Hotline by phone at 1-800-267-6315.

Visit <http://pr-rp.hc-sc.gc.ca/lr-re/index-eng.php> to access the Electronic Label Search Tool. The PMRA Product Information database can be searched by a product's trade name, active ingredient, company name or Registration (PCP) number. Since several products can contain the same active ingredient and there are often several versions of the same or similar labels on this database, using the PCP number is the most direct route to finding the label that links to the product page in this Guide. **There may be some differences between a label found on the package and the sample labels found on the PMRA-Label Search web site so always refer to the packaged product label when applying the product.**

Once the product is located, you may click on its number to view an Adobe Acrobat (PDF) document containing the label and any supplemental registrations. Some of these documents run to many pages but you can use the 'Find' capabilities of the Acrobat Reader plug-in for your browser to jump to specific areas of interest. If you do not have Adobe Acrobat Reader installed on your computer, you can download a free version from www.adobe.com.


Safe Use of Herbicides, Fungicides and Insecticides

Herbicides, fungicides and insecticides are classified according to the use hazard and risk involved. The categories of hazard are:

- toxicity
- flammability
- explosive potential
- corrosivity
- other




The degree of risk is represented by symbols taken from common traffic sign shapes represented by the stop, caution and yield signs. The signal word for each of the signs is danger (high risk), warning (moderate risk) and caution (low risk). Where the risk is minimal, no designation is required. The label on the container will carry the appropriate signs for the protection of the user. Degree of risk symbols for herbicides, insecticides and fungicides used in field and forage crops are included in the product directory. The symbols are illustrated in Figure 1.

Figure 1. Degree of Risk and Hazard Symbols



LD₅₀ values are used to rate the toxicity of pesticides. The LD₅₀ refers to the dose of pesticide (in mg per kg of the test animal's body weight) that is lethal to 50 percent of the group of test animals. For example, if a pesticide has an LD₅₀ value of 10 mg/kg, and the test animals each weigh 1 kg, then 50 percent of the animals would die if they each ate 10 mg of the pesticide.

Table 2. Oral LD₅₀ Values as they relate to the Risk/Hazard Symbols

 DANGER POISON LD ₅₀ less than 500 mg/kg Indicates high toxicity	 WARNING POISON LD ₅₀ 500-1,000 mg/kg Indicates moderate toxicity
 CAUTION POISON LD ₅₀ 1,000-2,500 mg/kg Indicates low toxicity	SYMBOL ABSENT LD ₅₀ greater than 2,500 mg/kg Indicates very low toxicity

Different types of protective equipment are required for pesticides that differ in toxicity. Special equipment requirements are described on the product label, but in general the following precautions must be taken when using pesticides of different hazard ratings.

- **Danger Poison** – requires goggles, respirator, gloves and skin protection, avoid fumes and spray mist.
- **Warning Poison** – requires goggles, gloves and skin protection, avoid fumes and spray mist.
- **Caution Poison** – requires gloves and skin protection, avoid fumes and spray mist.

The absence of a hazard symbol on a pesticide label indicates low toxicity to mammals. Nevertheless, protective clothing should be worn when using pesticides that do not have a hazard symbol.

Protecting Yourself from Exposure to Herbicides, Fungicides and Insecticides

The use of protective equipment and sound safety procedures will help minimize your exposure to herbicides, fungicides and insecticides. Follow the 10 rules for safe application listed below, and wear the safety equipment recommended.

10 Rules for Safe Application

1. Never smoke or eat while applying pesticides.
2. Avoid inhaling sprays or dusts. Wear protective clothing and a respirator.
3. Sprayer lines carrying chemicals should not enter the operator's cab.
4. Have soap, water and a towel available. Should concentrated product spill on skin, hands, face or eyes, wash immediately.
5. Wash hands and face when leaving the treated area, before break periods, lunch or urination.
6. Bathe or shower and change into clean clothing after working with pesticides. Wash clothing each day before re-use.
7. Call a physician or get the patient to a hospital immediately if symptoms of illness occur during or shortly after pesticide application. Be sure to take along the product label or container.
8. Store pesticides out of reach of children and where there is no chance of contact with human food or livestock feeds. Do not store herbicides with insecticides and avoid cross-contamination. Storage areas should be locked.
9. Keep chemicals in their original containers, never in unmarked containers or bottles used for food or drink.
10. Follow proper container disposal methods. All containers should be triple rinsed or pressure rinsed, punctured to render the container non-reusable, and delivered to designated disposal sites.

Protective Clothing

Wear protective equipment as described in the chart to reduce exposure.

EQUIPMENT	PROTECTION	HOW TO WEAR IT
Coveralls	<p>There are two types of coveralls: disposable and reusable.</p> <p>Disposable coveralls are lightweight and comfortable on warm days. They can be worn for mixing and applying pesticides, then discarded at the day's end. If they become contaminated, they should be discarded at once.</p> <p>The second type of coverall is made of washable fabric and may be reused many times. These fabric coveralls are adequate for use with all but the most highly toxic and concentrated pesticides.</p>	<p>Button (or zip) right up to the neck. Loose coveralls around the neck will suck and blow pesticide in and out of the interior of the coveralls as you bend and move.</p> <p>Wear coveralls over a long-sleeved shirt and pants.</p>
Aprons	<p>When pouring or otherwise handling concentrated pesticides, it makes good sense to wear protection in the form of an apron. The apron protects the front of your body from spills or splashes of the concentrate.</p> <p>The apron should be made of rubber or synthetic liquid-proof material that will resist the solvents.</p>	<p>Make sure the apron covers your body from your chest to your boots.</p>
Gloves	<p>Protect your hands by wearing chemical-resistant gloves. Neoprene gloves provide the best protection. Natural rubber gloves may be used when handling organo-phosphorus or carbamate pesticides. Be sure that they are designed for use with solvents and pesticides. Never use lined gloves, gloves with wristbands or leather gloves.</p>	<p>Put gloves on and roll up the first inch or two of the cuff. That way when you lift your hands, any liquid on the gloves won't drip down your arms.</p>
Hats	<p>Use a chemical-resistant hat, preferably made of washable plastic. The hat may be a hard hat or made of flexible plastic. In either case, it should have a plastic sweatband. Wash and dry entire hat after each use and before storing. Ordinary baseball caps with cloth sweatbands are dangerous as they absorb the pesticide and recontaminate the forehead each time you wear them. Even small amounts of moderately or slightly toxic pesticides may cause severe skin irritation or other illness if exposure continues for several days.</p>	
Boots	<p>Wear chemical-resistant, unlined boots. These boots are available in a variety of styles and materials.</p> <p>Neoprene boots are the best. Knee-length boots offer greater protection because they extend above the lower end of the apron. Avoid leather or fabric boots and shoes because these will absorb pesticides and cannot be cleaned effectively.</p>	<p>Wear your pant legs outside the top of your boots. This will prevent spills and splashes from running into the boot and onto your leg.</p>

Protecting Your Eyes, Face and Lungs

Wear the following equipment to protect your facial area from exposure.

EQUIPMENT	PROTECTION	HOW TO WEAR IT
Goggles	<p>Chemical-resistant goggles keep your eyes safe from both splashing and, if using dry formulations, dusts or granules. Don't use goggles with cloth or elastic headbands as these will absorb pesticides.</p>	<p>Wear goggles snugly on your face so that the sides of your head are protected from splashes. If you wear glasses, make sure you purchase goggles that fit snugly over them. Never wear contact lenses when working around pesticides.</p>

Respirators	Only NIOSH-approved respirators should be used. Do not exchange parts of different respirators. (For example, do not use a cartridge produced by Company "A" with a respirator produced by Company "B" as the combination may not provide adequate protection to the user). Dust masks are ineffective in protecting against herbicide vapours. Similarly, the filters on tractor cabs are intended to remove dust and are not designed to protect against herbicide vapours or mists. Chemical cartridge respirators are recommended for outdoor use when mixing and applying herbicides.	When carrying out operations, change filters each day. The cartridge should be replaced when chemical odour becomes apparent or when breathing becomes difficult. New cartridges should always be installed at the beginning of the spray season. Prior to commencing work, check the face seal while the respirator is on the wearer's face. Regardless of design, respirators cannot be worn securely by people wearing beards, moustaches or sideburns.
Face Shields	Goggles offer some protection, but frequently full-face protection is advised or required according to the pesticide label. It is especially important to protect your eyes and face when pouring or mixing liquid concentrates. Effective face shields are made of clear plastic.	Since the shield attaches to the hard hat, you can raise or lower it as needed.

Understanding Maximum Residue Limit Statements in the Guide

To ensure the safety of Canadian food, maximum residue limits (MRLs) set the maximum allowable amount of a pesticide residue on a crop or in a processed crop product (e.g. oil or flour). Residue levels are typically assessed for pesticides registered on crops grown for food. MRLs even exist on imported food for pesticides or pesticide uses not registered in Canada.

Health Canada's Pest Management Regulatory Agency (PMRA) is responsible for setting MRLs in Canada. Similarly, importing countries set their own MRLs (also referred to as 'import tolerances') that Canadian crop exports are subject to. Trade issues between importing and exporting countries can arise due to variability in MRLs or a lack of established MRLs.

Crop pesticide uses that may contribute to trade irritations have been flagged on product pages in the Guide to Crop Protection with the statement: **'Note: As of January 1, 2019 <http://keepingitclean.ca> indicates that grain from crops treated with this product may have market access concerns. Please see pg 10 for more information AND consult potential grain buyer(s) before using this product.'** Manitoba Agriculture and Saskatchewan Ministry of Agriculture have included such statements on products uses with known or potential MRL issues. However, this may not be a complete list of product uses with potential trade issues.

Producers can follow these practices to help prevent exceeding MRLs:

- Read and follow product labels, especially with respect to registered crops, maximum application rates, maximum number of applications per season, crop stage and pre-harvest intervals.
- Talk to your commodity buyer before applying a pesticide, especially for new pesticide chemistries, new products and products registered on new crops.

More information on MRLs and 'flagged' products is available at <http://keepingitclean.ca/>.

Avoiding Spray Drift

To minimize the risk of drift, follow these guidelines:

1. Do not spray in winds above 16 km/hr (10 m/hr).
2. Do not spray under dead calm conditions in early morning, night, or late evening. These are often associated with temperature inversions, and the combination of these factors can result in long-distance spray drift (2 km or more). Fog or dust that seems to hang in the air is a good indicator of an inversion.
3. Avoid nozzle pressures above 45 psi (310 kPa) for conventional flat fan tips.
4. Use a minimum of 45 L/acre water for all pesticides unless otherwise specified for the product.
5. Take note of buffer zones identified in the "Restrictions" section of this guide. Do not spray when the wind is blowing towards a nearby sensitive crop, shelterbelt, garden, or water body.
6. Use amine formulations of 2,4-D or MCPA where possible. Use special care when applying volatile herbicides (most herbicides in Group 3 and Group 4, particularly ester formulations). Avoid spraying these products on or immediately before hot days.
7. Ensure that air flow from air assisted sprayers is properly set to minimize airblast rebound and drift for different crop canopies.
8. Operate nozzles at their minimum recommended height. For 80° tips, this is 18" (45 cm), and for 110° tips, this is 12" (35 cm). Orienting nozzles forward allows further height reductions.
9. Special nozzles are now available that create coarse, low-drift sprays. Pre-orifice, Turbo-TeeJet, or venturi-type nozzles are available from a number of manufacturers, and these reduce drift by 50 to 95 percent. (Refer to the section entitled **Herbicide Efficacy with Low-Drift Nozzles**).
10. Consider equipping your sprayer with protective shrouds. A number of different designs are available that can reduce drift between 35 and 75 percent.
11. Reduce travel speeds. Rapid air movement over nozzle tips increases the risk of fine droplets prone to drift and turbulence from the sprayer itself can increase the uncertainty of spray deposition.

For more information on reducing drift, see the website: www.Sprayers101.com.

Herbicide Efficacy with Low-drift Nozzles

A number of low-drift nozzles are now available from different suppliers. Well established nozzles, such as the Turbo TeeJet, reduce drift by about 50 percent and provide equivalent efficacy to a standard flat fan nozzle. Newer nozzles (“venturi” types) are best known for their dramatic ability to reduce drift (50 to 95 percent). Research suggests that these nozzles perform well at conventional carrier volumes, travel speeds, and product rates. Some aspects require special attention:

Pressure: Some venturi-type nozzles require higher pressures to operate properly. Below 40 psi (275 kPa), patterns for these designs may deteriorate rapidly resulting in poor overlaps and erratic control. Design improvements have resulted in venturi nozzles that require less pressure to operate effectively. When using automatic rate controllers, make sure your pressures match the recommended pressure ranges for good nozzle performance.

Water Volume: Droplet size becomes more important at lower water volumes. Little is known about low-drift nozzle performance at or below 5 gal/acre (23 L/acre). Since low-drift nozzles generate fewer droplets than conventional nozzles, ensure that water volumes are high enough for coverage when using coarse sprays.

Weed Type: Difficult-to-wet weeds, such as wild oats, green foxtail, lamb’s-quarters, and cleavers, typically require finer sprays for effective coverage. When using venturi nozzles on these weeds, make sure your pressure is high enough to achieve good coverage. Larger weeds and reduced product rates typically make chemical control more difficult, and these conditions may also reveal some performance differences between nozzles.

Herbicide Type: Herbicides that belong to herbicide Groups 2, 4, and 9 perform well with venturi nozzles, even at normal pressures (40 psi). Application of herbicides in Groups 1, 6, 8, 10 and 14 may require higher pressures with venturi nozzles to maintain good performance, especially under challenging conditions. Wild oat control may be reduced with the coarsest sprays, even when applied at high pressure.

Check with your chemical representative to see if the manufacturer supports the use of low-drift nozzles with their products.

More information is available in the factsheet “Pesticide Application and Choosing the Right Nozzles,” available from your local extension office or at the Saskatchewan Ministry of Agriculture Website: Saskatchewan.ca/agriculture.

Handling a Drift Complaint

When spray drift occurs, it is important to take the right steps to resolve the complaint. If you suspect that your crop or property has been damaged because of spray drift, use the following guidelines for resolving the situation.

1. Contact the suspected applicator as soon as possible. View the damage with the suspected applicator and determine if that person did, in fact, cause the damage.
2. Are you sure that the symptoms or damage you see has been caused by spray drift? Contact your local agriculture office or agronomist to discuss the injury symptoms.
3. If the damage was caused by the applicator, determine the extent of the damage and the level of compensation (if any) with the applicator.

4. If the situation cannot be resolved quickly because of disagreements on the extent of damage, cause of the damage, or level of compensation, contact your local agricultural office to discuss options on how to proceed. Documentation will be required, particularly if insurance companies are involved.
5. The involvement of a private consultant is recommended if documentation is required. Required documentation often includes samples of the damaged plants, photographs, and yield comparisons to determine losses. Your agricultural office can provide you with a list of private consultants in your area.
6. The best approach is to start an open and honest line of communication with the suspected applicator. The majority of drift complaints are resolved quickly and efficiently by communicating with the applicator, without the involvement of outside parties.

Mixing Pesticides

The ability to control a broad range of weeds or other pests in one pass is the advantage that a mix of two or more products allows. If tank mixing is not done in the correct order, the result could be a tank-load of material that may not control the target pests, cause injury to the crop, plug nozzles, or leave an undesirable residue in the tank that will require extensive cleaning. Mistakes like these are costly, could put the user at unnecessary risk of exposure to the products, or create an environmental disposal problem.

To avoid mixing that may result in incompatibilities, **always consult the label of the products that are being used** to learn the correct order. **Remember to add all like components at the same stage of mixing.** The list below is a general rule-of-thumb for mixing pesticides:

1. Fill the spray tank with 1/4 to 3/4 the amount of water required for the application and turn on the sprayer agitation. Check the products that are being used for the correct amount to add. Once agitation has begun, maintain until the tank is emptied.
2. Add any water conditioner (fertilizer or pH adjuster) additives to the tank.
3. Add any wettable powders, or water dispersible granules (DF, DG, or WDG). Add dry products slowly to prevent clogged return lines. Allow sprayer to agitate for a few minutes, allowing the product to become completely suspended in the tank, before adding the next component.
4. Shake any containers of liquid pesticide thoroughly before adding to ensure they are well mixed.
5. Add any oil dispersions (OD) or flowable liquid suspensions (F, SC) to the tank. Allow to mix.
6. Add emulsifiable concentrates (EC) or emulsions (ME, SE) to the tank and allow to mix.
7. Add any pesticides that are solutions (SN) (i.e. amines and salts)
8. Add any surfactants or other adjuvants.

Remember to always consult the label for compatible mixes and recommended mixing order.

Many pesticides will break down if left in the tank for an extended period. Try not to mix any more than you can spray at one time.

If you need to stop spraying for a short time, leave the sprayer agitation running to keep products from settling or separating in the tank.

Container Disposal

Proper disposal of used containers and unused pesticides is important to protect the environment and prevent contamination of soil and water resources. Rinse all containers prior to disposal to reduce environmental contamination caused by open dumping of unwanted containers. Only mix as much pesticide solution as is needed to treat the desired area.

Triple Rinsing

Triple rinsing renders used pesticide containers (metal, plastic, glass) more than 99.9 percent free of residues, in most cases. Here are the steps that should be followed:

1. Empty contents of the container into the spray tank and drain in a vertical position for 30 seconds.
2. Add a measured amount of rinse water or other diluent until container is about one-fifth full.
3. Rinse the container thoroughly and pour the rinsate into the spray tank.
4. Repeat the procedure twice (it should take only about 5 minutes in total).
5. Puncture or break triple rinsed containers to render them non-reusable. Paper bags should be rinsed once prior to disposal.

Pressure Rinsing

Pressure rinsers can be used to rinse any size of empty pesticide container that can be lifted into position over the spray tank. A 30 second rinse with a pressure rinser is convenient and just as effective as triple rinsing. Pressure rinsers are constructed to be thrust into the bottom of a metal can or plastic jug. Holes, situated laterally in the rinser tip, direct water from a pressurized source against the inner sides of the container and effectively wash the residual pesticide into the spray tank. Some farmers have found it convenient to attach a rinser to the pump on their large water storage tank to minimize container handling. Pressure rinsers have the added advantage of rendering containers useless by automatically puncturing them.

Disposal of Containers

Properly rinsed containers should be delivered to a designated pesticide container disposal site. Contact your ag Provincial Agriculture Office, municipal office or weed supervisor for the locations of pesticide container disposal sites in your municipality for more information on pesticide container recycling see www.cleanfarms.ca.

Sprayer Cleaning

When pesticide application is completed each day it is important to empty and clean the sprayer thoroughly to prevent the breakdown of certain pesticides, prevent adhesion of the pesticide to the sprayer, and to maintain the sprayer parts in good condition. Certain pesticides break down very quickly when left in solution, and several pesticide solutions can be corrosive to sprayer parts. Sprayer cleaning is especially important when changing from

one crop to another or from one pesticide to another. Each year several reports are logged of herbicide damage cause by carryover of product residue in the tank. To avoid the risk of contamination, sprayers should be cleaned as soon as possible after application is completed.

Do not clean sprayers where rinsate can run off into ditches or other water bodies, near sensitive plants or shelterbelts, or where other people or animals are likely to walk, to avoid unnecessary exposure to people, animals and the environment.

There are three basic types of rinse solution for cleaning sprayer tanks. Their recipes and basic procedures are outlined below:

- **The Ammonia Rinse** – Fill spray tank and add 1 L of household ammonia (3%) for every 100 L of clean water needed for the rinse and begin agitation. Allow solution to flush through the booms until the boom is completely filled with ammonia solution and top up the tank with water. Circulate the ammonia solution through the tank and pump system for 15 minutes. Flush hoses and booms with ammonia rinse solution again (minimum 5 minutes) before emptying. Remove nozzles and screens and scrub with 0.1 L household ammonia per 10 L clean water and an old toothbrush. Perform clean water rinse to remove ammonia solution prior to next spray load. Some herbicides recommend leaving the ammonia rinse in the tank over night to improve cleaning potential.
- **The Fresh Water Rinse** – The spray tank cleaning should begin and end with a fresh water rinse to remove the majority of potential contaminants prior to the cleansing process or prior to the next round of spraying. Drain the tank of its previous contents and fill the tank with clean water. Open nozzle valves and pump clean water through the booms and hoses. Top up the tank with more clean water and circulate/agitate for at least 10 minutes and empty the tank of waste water. If this is the first rinse after spraying, a high pressure hose could be used to clean residue from all surfaces in the tank. Do not enter the tank during the cleaning process.
- **The Detergent Rinse** – After rinsing with clean water, fill spray tank and add a heavy-duty detergent at 0.25 L per 100 L of water (some suggest a non-ionic surfactant such as Agral 90 or Agsurf at 0.6 L per 100 L of water). Circulate the mixture for a minimum of 5 minutes and spray out through sprayer nozzles. Nozzles and screens are removed and cleaned individually with the same detergent solution in a small container. Soaking in this solution for several hours also helps to loosen any deposits.

The above solutions are just components of the overall sprayer cleaning process. Typical rinse instructions will repeat a combination of one or two or all of these basic rinses. Below we will give some generic rinse instructions utilizing the basic rinses as components of the larger cleaning procedure. Never enter the tank during the cleaning process as some cleansers may release dangerous gases.

- **Method A** – Drain contents of tank – 1 to 2X Water Rinse – 2X Ammonia Rinse – 2X Water Rinse (one just prior to the next spraying event)
- **Method B** – Drain contents of tank – 2X Water Rinse – 2X Detergent Rinse – 2X Water Rinse

- **Method C** – Drain contents of tank – Several repetitions of the Water Rinse with nozzles and screens removed and checked for debris. Products: Adrenalin, Altitude, Amitrol 240, Ares.

The above directions are general processes based on the similarities of tank cleaning recommendations between products in each of the herbicide groupings. Always follow the specific instructions on the product label.

Several products in the guide do not have label instructions regarding tank cleaning. In the case of products that have no cleaning recommendations on the label, there are some basic principals that can be applied. Products that are water based formulations can usually be cleaned from spray tanks using **Method C** above. Products that are formulated as an EC, SC or F (flowable) or use a petroleum based adjuvant should at least use **Method B**. The detergent breaks down the oil that may be sticking to the side of the tank. Products in Group 2 (most will already have a recommendation), with the exception of the 'IMI' products (see Table 8 on page 45), will require the use of **Method A**. The ammonia in **Method A** either increases the solubility of the product allowing it to be easily removed from the tank surfaces or speeds the breakdown of these products in water. If the product that is to be cleaned out of the tank is a combination of these elements, use a combination of Methods to clean the tank. In these cases, use a good commercial tank-cleaning product from a recognized source, with both ammonia and detergent as components.

Group 2 compounds are highly active on sensitive plants so even a small amount remaining in the sprayer can present a risk of injury. They can also occasionally be trapped on the tank walls and plumbing by petroleum based formulations or adjuvants when tank mixed with other products, resulting in tank residues that may be tougher to remove. A way to reduce the chance of this occurring is to add detergent at 0.25 L per 100 L to the Ammonia

Rinse portion to assist with the breakdown of the petroleum coating so that the ammonia may rid the tank of Group 2 product.

It is very important to clean sprayers immediately after every use. With a more diverse rotation, the likelihood of damage from lack of care increases dramatically.

How to Identify Crop and Weed Leaf Stages

Recognition of plant growth stages is essential for effective weed and disease control. Many herbicides and fungicides are safe on a crop only when applied at a specific growth stage. Similarly, weeds are controlled only when they are at certain growth stages.


For most post-emergent products, growth stages are described by the number of leaves. The following is a description of how to count leaves for staging.

Cereals and Annual Grass Weeds

Manufacturers generally use two different systems of staging for grasses. The minimum stages of application are similar, while the later stages may differ.

Some manufacturers use "Total Leaf Count" stages based on the number of leaves on the entire plant, including tillers or secondary shoots. Most recommendations are based on the number of main stem leaves and tillers. Tillers or stools are the secondary shoots or stems of a grass plant. Similar to the branches of a broadleaf plant, tillers will emerge from the axils between the leaf and main shoot. Tillers usually begin to appear at the 3 or 4 leaf stage. When staging a plant in this manner, be sure to identify the tillers first, then count only leaves that originate from the main shoot.

Figure 2. Leaf Stages of Cereals and Annual Grass Weeds



Broadleaf Weeds

Cotyledons – These are the seed leaves that usually emerge above ground. On some plants, such as faba beans, lentils and peas, they stay below the soil surface. Cotyledons are not true leaves and are not counted when determining leaf number. They are a different shape than the true leaves and may dry up and disappear at an early stage.

Alternate leaves – Some plants have one leaf at each node on the stem. The next leaf emerges at the next higher node and extends away from the stem in the opposite direction. These plants (lamb's quarters and wild mustard are good examples) are said to have alternate leaves. To determine the leaf stage, simply count the number of leaves present (Figure 3).

Opposite leaves – Plants with two leaves at each node, one on each side of the stem, are said to have opposite leaves. The next pair of leaves on the next node are rotated about 45° so that they are not directly over the previous pair. Plants with opposite leaves have even-leaf numbers only. When counting, the leaf number progresses from cotyledons to 2 leaf, 4 leaf, etc. These plants generally appear shorter than plants with alternate leaves at a similar leaf stage. **Be sure to count each pair as two leaves.** Hemp nettle is a weed that has opposite leaves (Figure 3).

Whorled leaves – More complex plants like cleavers may have whorled leaves. These plants have three or more leaves at each node on the stem. The leaf number in each whorl may vary, so be sure to count each individual leaf unless the Guide or label recommendation refers to the number of leaf whorls (Figure 3).

Figure 3. Leaf Stages of Broadleaf Weeds and Crops








Figure 4. Leaf Stages of Certain Special Crops and Forages



Stages of alfalfa, red clover and alsike clover leaf development



Trade Names, Active Ingredients and Formulations

Legend: (F) = formulated component; (B) = blended granules; (DC) = divided container

WP wettable powder EC emulsifiable concentrate SP water soluble powder SL soluble concentrate A amine
 WG water dispersible granule G granule EW emulsion, oil in water OD oil dispersion E ester
 TB Tablet SC suspension concentrate DS powder/dust for dry seed treatment FS flowable concentrate for seed treatment LS solution for seed
 SG water soluble granule AS aqueous suspension treatment

Herbicides

(Component 1) Active Ingredient* - Formulation	(Component 2) Active Ingredient* - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
2,4-D Amine - 200 g/L SL (F)	aminopyralid - 40 g/L SL (F)			Restore II	Corteva Agriscience
2,4-D Amine - 295 g/L SL (F)	dicamba - 110 g/L SL (F)	mecoprop-P - 80 g/L SL (F)		DyVe/DSp	BASF Canada
2,4-D Amine - 600, 700 g/L SL				2,4-D Amine or Ester	Various
2,4-D Ester - 225 g/L EC (F)	bromoxynil - 225 g/L EC (F)			Thrasher II	ADAMA Canada
2,4-D Ester - 225 g/L EC (F)	bromoxynil - 225 g/L EC (F)			Leader	IPCO
2,4-D Ester - 225 g/L EC (F)	bromoxynil - 225 g/L EC (F)			Approve	Nufarm Agriculture
2,4-D Ester - 240 g/L EC (F)	bromoxynil - 90 g/L EC (F)			Enforcer D	Nufarm Agriculture
2,4-D Ester - 280 g/L EC (F)	bromoxynil - 280 g/L EC (F)	fluroxypyr - 80 g/L EC (F)		Thumper	Bayer CropScience
2,4-D Ester - 282 g/L EC (F)	dichlorprop - 300 g/L EC (F)			Dichlorprop-D	IPCO
2,4-D Ester - 282 g/L EC (F)	dichlorprop - 300 g/L EC (F)			Turboprop	UAP
2,4-D Ester - 360 g/L EC (F)	fluroxypyr - 90 g/L EC (F)			OcTrain XL	Corteva Agriscience
2,4-D Ester - 360 g/L EC (F2)	fluroxypyr - 90 g/L EC (F2)	clodinafop - 25 g/L EC (F1)	pinoxaden - 25 g/L EC (F1)	TraxosTwo	Syngenta Canada
2,4-D Ester - 360 g/L SL (F)	picloram - 97.5 g/L SL (F)			Grazon XC	Corteva Agriscience
2,4-D Ester - 400 g/L EC (F)	dichlorprop-P - 210 g/L EC (F)			Dichlorprop-DX	IPCO
2,4-D Ester - 400 g/L EC (F)	dichlorprop-P - 210 g/L EC (F)			Estaprop XT	Nufarm Agriculture
2,4-D ester - 473 g/L (F)	pyraflufen - 6.1 g/L (F)			BlackHawk (new)	Nufarm Agriculture
2,4-D Ester - 564 g/L EC	thifensulfuron:tribenuron - 33.3%:16.7% SG	fluroxypyr - 333 g/L EC		Retain SG (new)	Loveland Products
2,4-D Ester - 564 g/L EC	thifensulfuron:tribenuron - 33.3%:16.7% SG	fluroxypyr - 180 g/L EC		Retain SG (old)	Loveland Products
2,4-D Ester - 660 g/L EC	fluroxypyr - 180 g/L EC			Rush 24	ADAMA Canada
2,4-D Ester - 660 g/L EC	florasulam - 50 g/L SC			MPower Battlefront 2,4-D	Agracity
2,4-D Ester - 660 g/L EC	fluroxypyr - 180 g/L EC			MPower Foxy Pro	Agracity
2,4-D Ester - 660 g/L EC	tribenuron - 75% WG			MPower X-Ko	Agracity
2,4-D Ester - 660 g/L EC	fluroxypyr - 333 g/L EC			Attain XC	Corteva Agriscience
2,4-D Ester - 660 g/L EC	florasulam - 50 g/L SC			Frontline 2,4-D XC	Corteva Agriscience
2,4-D Ester - 660 g/L EC	aminopyralid - 52.5% WG (F)			Reclaim II	Corteva Agriscience
2,4-D Ester - 660 g/L EC	pyroxulam - 15% WG (F)	metsulfuron - 9.45% WG (F)		Rexade	Corteva Agriscience
2,4-D Ester - 660 g/L EC	carfentrazone - 240 g/L EC	hauloxifen - 5% WG (F)		BlackHawk (old)	Nufarm Agriculture
2,4-D Ester - 660 g/L EC	fluroxypyr - 180 g/L EC			Flurox-24	Nufarm Agriculture
2,4-D Ester - 660 g/L EC	tribenuron - 75% WG			Ko-Act	Nufarm Agriculture
2,4-D Ester - 660 g/L EC	tribenuron - 8.25% WG (F)	dicamba - 58.45% WG (F)		Salvo	UAP
2,4-D Ester - 660g/L EC				Triton K	FMC
2,4-DB Ester - 625 g/L EC				Cobutox 625	IPCO
2,4-DB Ester - 625 g/L EC				Caliber	Loveland Products
2,4-DB Ester - 625 g/L EC				Embutox 625	Nufarm Agriculture
aminocyclopyrachlor - 39.5% WG (F)	metsulfuron - 12.6% WG (F)			Navius	Bayer Environmental Sciences
acifluorfen - 240 g/L SL				Ultra Blazer	United Phosphorus Inc.
aminopyralid - 40 g/L SL (F)				Restore II	Corteva Agriscience
aminopyralid - 52.5% WG (F)		2,4-D Ester - 660 g/L EC		Reclaim II	Corteva Agriscience

(Component 1) Active Ingredient** - Formulation	(Component 2) Active Ingredient** - Formulation	(Component 3) Active Ingredient** - Formulation	(Component 4) Active Ingredient** - Formulation	Product	Company
amitrole - 231 g/L SL	metolachlor - 400 g/L SC (F)			Amitrol 240	Nufarm Agriculture
atrazine - 320 g/L SC (F)	imazamox - 20 g/L SL (F)			Primextra II Magnum	Syngenta Canada
atrazine - 480 g/L SC				AAtrex Liquid	Syngenta Canada
bentazon - 429 g/L SL (F)				Viper ADV	BASF Canada
bentazon - 480 g/L SL				MPower Boa	Agracity
bentazon - 480 g/L SL				Basagran	BASF Canada
bentazon - 480 g/L SL				Basagran Forté	BASF Canada
bentazon - 480 g/L SL				Benta Super	Sharda CropChem
bromoxynil - 175 g/L EC (F)	pyrasulfotole - 31.3 g/L EC (F)	thiencarbazone - 5 g/L SC (F)		Velocity m3	Bayer CropScience
bromoxynil - 200 g/L EC (F)	fluroxypyr - 80 g/L EC (F)	MCPA Ester - 225 g/L EC (F)		Enforcer M	Nufarm Agriculture
bromoxynil - 200 g/L EC (F)	fluroxypyr - 80 g/L EC (F)	MCPA ester - 200 g/L EC (F)	thifensulfuron:tribenuron - 50%:25% SG	Enforcer MSU	Nufarm Agriculture
bromoxynil - 210 g/L EC (F)	pyrasulfotole - 37.5 g/L EC (F)			Infinity	Bayer CropScience
bromoxynil - 210 g/L EC (F)	pyrasulfotole - 37.5 g/L EC (F)	fluroxypyr - 180 g/L EC		Infinity FX	Bayer CropScience
bromoxynil - 210 g/L EC (F)	pyrasulfotole - 37.5 g/L EC (F)	pinoxaden - 50 g/L EC		Axial iPak	Syngenta Canada
bromoxynil - 210 g/L EC (F)	pyrasulfotole - 37.5 g/L EC (F)	fluroxypyr - 87.5 g/L EC (F)	pinoxaden - 50 g/L EC (F)	Axial Xtreme iPak	Syngenta Canada
bromoxynil - 225 g/L EC (F)	MCPA Ester - 225 g/L EC (F)			Badge	ADAMA Canada
bromoxynil - 225 g/L EC (F)	2,4-D Ester - 225 g/L EC (F)			Thrasher II	ADAMA Canada
bromoxynil - 225 g/L EC (F)	MCPA Ester - 225 g/L EC (F)			MPower Buck M	Agracity
bromoxynil - 225 g/L EC (F)	2,4-D Ester - 225 g/L EC (F)			Leader	IPCO
bromoxynil - 225 g/L EC (F)	2,4-D Ester - 225 g/L EC (F)			Logic M	IPCO
bromoxynil - 225 g/L EC (F)	MCPA Ester - 225 g/L EC (F)			Approve	Nufarm Agriculture
bromoxynil - 225 g/L EC (F)	2,4-D Ester - 225 g/L EC (F)			Mextrol 450	Nufarm Agriculture
bromoxynil - 225 g/L EC (F)	MCPA Ester - 225 g/L EC (F)			ForceFighter	ADAMA Canada
bromoxynil - 225g/L EC (F)	carfentrazone - 240 g/L EC	fluroxypyr - 180 g/L EC		Conquer	Nufarm Agriculture
bromoxynil - 235g/L EC				Koril 235	Nufarm Agriculture
bromoxynil - 235g/L EC				Bromotril II	ADAMA Canada
bromoxynil - 240 g/L EC	florasulam - 50 g/L SC			Hot Shot	ADAMA Canada
bromoxynil - 240 g/L EC				MPower Bromoxynil	Agracity
bromoxynil - 240 g/L EC				Brotex	IPCO
bromoxynil - 240 g/L EC				Pardner	Bayer CropScience
bromoxynil - 240 g/L EC	MCPA Ester - 280 g/L EC (F)			Buctril M	Bayer CropScience
bromoxynil - 240 g/L EC	2,4-D Ester - 280 g/L EC (F)			Thumper	Bayer CropScience
bromoxynil - 280 g/L EC				Brotex 480	IPCO
bromoxynil - 280 g/L EC (F)				Loveland Bromax	Loveland Products
bromoxynil - 280 g/L EC (F)				Tundra	Bayer CropScience
bromoxynil - 280 g/L EC (F)	fenoxaprop-p - 46 g/L EC (F)	pyrasulfotole - 15.5 g/L EC (F)		Enforcer D	Nufarm Agriculture
bromoxynil - 480 g/L SL	fluroxypyr - 80 g/L EC (F)	2,4-D Ester - 240 g/L EC (F)		Aim EC	FMC
bromoxynil - 480 g/L SL				Authority Charge	FMC
bromoxynil - 87.5 g/L EC (F)	sulfentrazone - 480 g/L SC			Focus (co-pack)	FMC
bromoxynil - 90 g/L EC (F)	pyroxasulfone - 85% WG			BlackHawk (old)	Nufarm Agriculture
carfentrazone - 240 g/L EC	2,4-D ester - 660 g/L EC			Cleanstart	Nufarm Agriculture
carfentrazone - 240 g/L EC	glyphosate - 356 g/L SL			Conquer	Nufarm Agriculture
carfentrazone - 240 g/L EC	bromoxynil - 235g/L EC			Focus	FMC
carfentrazone - 53 g/L SE (F)	pyroxasulfone - 447 g/L SE (F)			Manipulator	Taminco/Engage Agro
chloromequat chloride - 620 g/L SL				Arrow-All-In	ADAMA Canada
clethodim - 120 g/L EC				Arrow	ADAMA Canada
clethodim - 240 g/L EC				MPower Independence	Agracity
clethodim - 240 g/L EC				Select	Arysta LifeScience
clethodim - 240 g/L EC				Centurion	Bayer CropScience
clethodim - 240 g/L EC				Patron 240EC	IPCO
clethodim - 240 g/L EC				Shadow RTM	Loveland Products
clethodim - 240 g/L EC				Statue	Nufarm Agriculture
clethodim - 240 g/L EC				Antler	Winfield United

(Component 1) Active Ingredient** - Formulation	(Component 2) Active Ingredient** - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
clethodim - 250 g/L EC clodinafop - 112 g/L (F)	fluroxypyr - 217 g/L EC (F)	thifensulfuron:tribenuron - 50%:25% SG		Clethodim 250 Signal/FSU	AgriStar Nufarm Agriculture
clodinafop - 240 g/L EC clodinafop - 240 g/L EC clodinafop - 240 g/L EC clodinafop - 240 g/L EC clodinafop - 240 g/L EC clodinafop - 25 g/L EC (F) clodinafop - 25 g/L EC (F1) clodinafop - 60 g/L EC clodinafop - 60 g/L EC clodinafop - 80 g/L EC clomazone - 360 g/L ME cloprralid - 360 g/L cloprralid - 360 g/L SL cloprralid - 360 g/L SL cloprralid - 360 g/L SL cloprralid - 360 g/L SL cloprralid - 50 g/L EC cloprralid - 50 g/L EC cloprralid - 50 g/L EC (F) cloprralid - 50 g/L EC (F) cloprralid - 50 g/L EC (F) cloprralid - 600 g/L SL cloprralid - 600 g/L SL cloprralid - 75% WG cloprralid - 75% WG cloprralid - 90g/L SL (F) dicamba - 110 g/L SL (F) dicamba - 120 g/L SL (F) dicamba - 350 g/L SL dicamba - 350 g/L SL dicamba - 46 g/L (F) dicamba - 480 g/L SL dicamba - 480 g/L SL dicamba - 480 g/L SL dicamba - 480 g/L SL dicamba - 480 g/L SL dicamba - 50% WG (F) dicamba - 50% WG (F) dicamba - 58.45% WG (F) dicamba - 600 g/L SL dicamba - 62.5 g/L SL (F) dicamba - 62.5 g/L SL (F) dicamba - 62.5 g/L SL (F) dicamba - 84 g/L SL (F) dicamba - 87 g/L EC (F) dichlobenil - 4% G	pinoxaden - 25 g/L EC (F) pinoxaden - 25 g/L EC (F1) hauloxifen - 16.2 g/L EC (F) glyphosate - 480 g/L SL glyphosate - 480 g/L SL florasulam - 50 g/L SC fluroxypyr - 333 g/L EC fluroxypyr - 333 g/L EC florasulam - 50 g/L SC MCPA Ester - 280 g/L EC (F) hauloxifen - 16.2 g/L EC (F) imazamox - 70% WG imazapyr - 15 g/L SL (F) fluroxypyr - 90 g/L EC (F) mecoprop-P - 80 g/L SL (F) glyphosate - 240 g/L SL (F) glyphosate - 194 g/L SL (F) florasulam - 25% WG tribenuron - 50% SG diflufenzopyr - 20% WG (F) diflufenzopyr - 20% WG (F) tribenuron - 8.25% WG (F) mecoprop-P - 62.5 g/L SL (F) mecoprop-P - 62.5 g/L SL (F) mecoprop-P - 62.5 g/L SL (F) MCPA K+ - 336 g/L SL (F) fluroxypyr - 113 g/L EC (F) 2,4-D Ester - 400 g/L EC (F) 2,4-D Ester - 400 g/L EC (F) MCPA Amine - 160 g/L SL (F)	2,4-D Ester - 360 g/L EC (F2) florasulam - 20% WG (F) MCPA Ester - 280 g/L EC MCPA Ester - 280 g/L EC MCPA Ester - 280 g/L EC (F) MCPA Ester - 280 g/L EC (F) florasulam - 20% WG (F) imazamox - 33 g/L SL (F) 2,4-D Amine - 295 g/L SL (F) 2,4-D ester - 660g/L EC MCPA amine - 275 g/L SL (F) MCPA amine - 275 g/L SL (F) MCPA amine - 275 g/L SL (F)	fluroxypyr - 90 g/L EC (F2)	Ladder MPower Aurora Slam'R Foax Signal Cadillac Traxos TraxosTwo FootHills NG Horizon NG Ladder All In Cadillac One Command 360 ME Cipreme MPower Clobber MPower Clobber G Eclipse III Lontriel 360 Pyralid MPower Battlefront CM MPower Foxy CM Prestige XC Spectrum Curtail M Cipreme XC Lontriel XC Tensile Salute Momentum DyVel DSp Roundup Xtend FeXapan Xtendimax Glykamba Banvel II Banvel VM Korrex II Express FX Oracle Distinct Overdrive Triton K Engenia Tracker XP Sword Target DyVel Pulsar Casoron Dichlorprop-DX Estaprop XT Optica Trio	ADAMA Canada Agracity AgriStar Great Northern Growers Nufarm Agriculture Winfield United Syngenta Canada Syngenta Canada Loveland Products Syngenta Canada ADAMA Canada Winfield United FMC Corteva Agriscience Agracity Agracity Corteva Agriscience Corteva Agriscience Sharda CropChem Agracity Agracity Corteva Agriscience Corteva Agriscience Nufarm Agriculture Corteva Agriscience BASF Canada Corteva Agriscience BASF Canada Corteva Agriscience Loveland Products BASF Canada Bayer CropScience Corteva Agriscience Nufarm Agriculture BASF Canada BASF Canada Corteva Agriscience FMC Gharda (UAP) BASF Canada BASF Canada FMC BASF Canada IPCO Loveland Products Syngenta Canada BASF Canada Syngenta Canada Arysta LifeScience IPCO Nufarm Agriculture UAP

(Component 1) Active Ingredient* - Formulation	(Component 2) Active Ingredient* - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
flumioxazin - 33.5% WG (F)	pyroxasulfone - 44.5% WG (F)			Fierce	Valent Canada
flumioxazin - 51% WG				Chateau	Valent Canada
flumioxazin - 51% WG				Vaiteira	Valent Canada
fluroxypyr - 100 g/L SC (F)	florasulam - 2.5 g/L SC (F)	MCPA ester - 600 g ae/L		Outshine	ADAMA Canada
fluroxypyr - 100 g/L SC (F)	florasulam - 2.5 g/L SC (F)	MCPA ester - 600 g/L		Stellar	Corteva Agriscience
fluroxypyr - 100 g/L SC (F)	florasulam - 2.5 g/L SC (F)	MCPA ester - 350 g/L (F)		Stellar XL	Corteva Agriscience
fluroxypyr - 113 g/L EC (F)	dicamba - 87 g/L EC (F)			Pulsar	Syngenta Canada
fluroxypyr - 150 g/L SC (F)	thifensulfuron - 30 g/L SC (F)			Travallas	FMC
fluroxypyr - 180 g/L	MCPA ester - 600 g/L EC			Rush M	ADAMA Canada
fluroxypyr - 180 g/L	MCPA ester - 600 g/LEC			MPower Foxy M	Agricity
fluroxypyr - 180 g/L	imazamox - 120 g/L SL			Altitude FX2	BASF Canada
fluroxypyr - 180 g/L	MCPA ester - 600 g/LEC			Trophy	Nufarm Agriculture
fluroxypyr - 180 g/LEC	MCPA ester - 225 g/LEC (F)	bromoxynil - 225g/L EC (F)		ForceFighter	ADAMA Canada
fluroxypyr - 180 g/LEC	2,4-D Ester - 660 g/LEC			Rush 24	ADAMA Canada
fluroxypyr - 180 g/LEC	2,4-D Ester - 660 g/LEC			MPower Foxy Pro	Agricity
fluroxypyr - 180 g/LEC				MPower Foxy	Agricity
fluroxypyr - 180 g/LEC	pyrasulfotole - 37.5 g/L EC (F)	bromoxynil - 210 g/L EC (F)		Infinity FX	Bayer CropScience
fluroxypyr - 180 g/LEC	thifensulfuron:tribenuron - 33.3%:16.7% SG			Ikwin	Great Northern Growers
fluroxypyr - 180 g/LEC	2,4-D Ester - 660 g/L EC	2,4-D Ester - 564 g/L EC		Retain SG (old)	Loveland Products
fluroxypyr - 217 g/LEC (F)	clodinafop - 112 g/L (F)				
fluroxypyr - 250 g/LEC (F)	MCPA ester - 600 g/LEC	thifensulfuron:tribenuron - 50%:25% SG		Flurox-24	Nufarm Agriculture
fluroxypyr - 250 g/LEC (F)	pinoxaden - 50 g/LEC (F)	hauloxifen - 16.2 g/L EC (F)		Signal FSU	Nufarm Agriculture
fluroxypyr - 333 g/LEC	clopyralid - 50 g/L EC	hauloxifen - 16.2 g/L EC (F)		Pixxaro	Corteva Agriscience
fluroxypyr - 333 g/LEC	2,4-D E - 660 g/LEC	MCPA Ester - 280 g/L EC		Rezuvant	Corteva Agriscience
fluroxypyr - 333 g/LEC	clopyralid - 50 g/LEC (F)			MPower Foxy CM	Agricity
fluroxypyr - 333 g/LEC	thifensulfuron:tribenuron - 50%:50% SG	MCPA Ester - 280 g/L EC (F)		Attain XC	Corteva Agriscience
fluroxypyr - 333 g/LEC	thifensulfuron:tribenuron - 50%:50% SG			Prestige XC	Corteva Agriscience
fluroxypyr - 333 g/LEC	thifensulfuron:tribenuron - 50%:50% SG			Tandem	Corteva Agriscience
fluroxypyr - 333 g/LEC	thifensulfuron:tribenuron - 50%:50% SG			Barricade II	FMC
fluroxypyr - 333 g/LEC	thifensulfuron:tribenuron - 50%:50% SG	thiencarbazone - 10 g/L SC	MCPA ester - 600 g/L EC	Predicade	FMC
fluroxypyr - 333 g/LEC	thifensulfuron:tribenuron - 33.3%:16.7% SG	2,4-D Ester - 564 g/L EC		Retain SG (new)	Loveland Products
fluroxypyr - 80 g/L EC (F)	2,4-D Ester - 240 g/L EC (F)	bromoxynil - 90 g/L EC (F)		Enforcer D	Nufarm Agriculture
fluroxypyr - 80 g/L EC (F)	MCPA - 200 g/L EC (F)	bromoxynil - 200g/L EC (F)		Enforcer M	Nufarm Agriculture
fluroxypyr - 80 g/L EC (F)	bromoxynil - 200 g/L EC (F)	MCPA ester - 200 g/L EC (F)	thifensulfuron: tribenuron - 50%:25% SG	Enforcer MSU	Nufarm Agriculture
fluroxypyr - 87.5 g/L EC (F)	pinoxaden - 50 g/L EC (F)			Axial Xtreme	Syngenta Canada
fluroxypyr - 87.5 g/L EC (F)	pinoxaden - 50 g/L EC (F)		pyrasulfotole-37.5 g/L EC (F)	Axial Xtreme iPak	Syngenta Canada
fluroxypyr - 90 g/L EC (F)	2,4-D Ester - 360 g/LEC (F)			OcTtain XL	Corteva Agriscience
fluroxypyr - 90 g/L EC (F)	clopyralid - 90g/L SL (F)			Momentum	Loveland Products
fluroxypyr - 90 g/L EC (F2)	2,4-D Ester - 360 g/LEC (F2)	pinoxaden - 25 g/L EC (F1)		TraxosTwo	Syngenta Canada
fomesafen - 240 g/L SL	glyphosate - 271 g/L SL (F)			Reflex	Syngenta Canada
fomesafen - 67 g/L SL (F)				FlexStar GT	Syngenta Canada
foramsulfuron - 22.5 g/L OD				Option 2.25 OD	Bayer CropScience
glufosinate - 150 g/L SL				MPower GoodHarvest	Agricity
glufosinate - 150 g/L SL				MPower Vigor	Agricity
glufosinate - 150 g/L SL				Liberty 150 SN	BASF Canada
glufosinate - 150 g/L SL				Liberty 200 SN	BASF Canada
glufosinate - 200 g/L SL				Glykamba	Nufarm Agriculture
glyphosate - 194 g/L SL (F)	dicamba - 46 g/L (F)			Roundup Xtend	Bayer CropScience
glyphosate - 240 g/L SL (F)	dicamba - 120 g/L SL (F)			FlexStar GT	Syngenta Canada
glyphosate - 271 g/L SL (F)	fomesafen - 67 g/L SL (F)				

(Component 1) Active Ingredient* - Formulation	(Component 2) Active Ingredient* - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
imazethapyr - 240 g/L SL				MPower Kamikaze	Agracity
imazethapyr - 240 g/L SL				Pursuit	BASF Canada
imazethapyr - 240 g/L SL				MulfiStar	Loveland Products
imazethapyr - 240 g/L SL				Gladiator	Univar Canada
imazethapyr - 35% WG (F)	imazamox - 35% WG (F)			MPower Ninja	Agracity
imazethapyr - 35% WG (F)	imazamox - 35% WG (F)			Odyssey	BASF Canada
imazethapyr - 35% WG (F)	imazamox - 35% WG (F)			Odyssey NXT	BASF Canada
imazethapyr - 35% WG (F)	imazamox - 35% WG (F)	sethoxydim - 450 g/L EC		Odyssey Ultra	BASF Canada
imazethapyr - 35% WG (F)	imazamox - 35% WG (F)	sethoxydim - 450 g/L EC		Odyssey Ultra NXT	BASF Canada
linuron - 400 g/L SC				Duet	Loveland Products
linuron - 480 g/L SC				Linuron 400	UAP
MCPA Amine - 160 g/L SL (F)				Lorox L	Tessenderlo Kerley Inc.
MCPA amine - 275 g/L SL (F)	mecoprop-P - 130 g/L SL (F)			Optica Trio	UAP
MCPA amine - 275 g/L SL (F)	dicamba - 62.5 g/L SL (F)			Tracker XP	IPCO
MCPA amine - 275 g/L SL (F)	dicamba - 62.5 g/L SL (F)			Sword	Loveland Products
MCPA amine - 275 g/L SL (F)	dicamba - 62.5 g/L SL (F)			Target	Syngenta Canada
MCPA Amine - 500, 600 g/L SL				MCPA Amine	Various
MCPA Ester - 200 g/L EC (F)	bromoxynil - 200g/L EC (F)			Enforcer M	Nufarm Agriculture
MCPA ester - 200 g/L EC (F)	bromoxynil - 200 g/L EC (F)			Enforcer MSU	Nufarm Agriculture
			thifensulfuron; tribenuron - 50%:25% SG		
MCPA Ester - 225 g/L EC (F)	bromoxynil - 225 g/L EC (F)			Badge	ADAMA Canada
MCPA Ester - 225 g/L EC (F)	bromoxynil - 225g/L EC (F)			ForceFighter	ADAMA Canada
MCPA Ester - 225 g/L EC (F)	bromoxynil - 225 g/L EC (F)			Logic M	IPCO
MCPA Ester - 225 g/L EC (F)	bromoxynil - 225 g/L EC (F)			Mextrol 450	Nufarm Agriculture
MCPA Ester - 280 g/L EC	florasulam - 50 g/L SC			MPower Battleground CM	Agracity
MCPA Ester - 280 g/L EC	fluroxypyr - 333 g/L EC			MPower Foxyy CM	Agracity
MCPA Ester - 280 g/L EC (F)	bromoxynil - 280 g/L EC (F)			Buctril M	Bayer CropScience
MCPA ester - 280 g/L EC (F)	florasulam - 4 g/L EC (F)			Frontline XL	Corteva Agriscience
MCPA ester - 280 g/L EC (F)	fluroxypyr - 333 g/L EC			Prestige XC	Corteva Agriscience
MCPA Ester - 280 g/L EC (F)	florasulam - 50 g/L SC			Spectrum	Corteva Agriscience
MCPA Ester - 280 g/L EC (F)	clopyralid - 50 g/L EC (F)			Curtail M	Nufarm Agriculture
MCPA Ester - 350 g/L (F)	fluroxypyr - 100 g/L SC (F)			Stellar XL	Corteva Agriscience
MCPA ester - 420 g/L (F)	pyraflufen - 13.5 g/L (F)			Goldwing	Nufarm Agriculture
MCPA ester - 500 or 600 g/L EC	thifensulfuron; tribenuron - 33.3%:16.7% SG			Refine M	FMC
	thifensulfuron; tribenuron - 33.3%:16.7% SG			BroadSide	Loveland Products
MCPA Ester - 500, 600 g/L EC	florasulam - 2.5 g/L SC (F)			MCPA Ester	Various
MCPA ester - 600 g ae/L	florasulam - 2.5 g/L SC (F)			Outshine	ADAMA Canada
MCPA ester - 600 g/L	fluroxypyr - 180 g/L			Stellar	Corteva Agriscience
MCPA ester - 600 g/L EC	florasulam - 50 g/L SC			Rush M	ADAMA Canada
MCPA Ester - 600 g/L EC	florasulam - 50 g/L SC			Topline	ADAMA Canada
MCPA Ester - 600 g/L EC	fluroxypyr - 180 g/L			MPower Battleground M	Agracity
MCPA ester - 600 g/L EC	hauloxifen - 16.2 g/L EC (F)			MPower Foxyy M	Agracity
MCPA ester - 600 g/L EC	thifensulfuron; tribenuron - 50%:50% SG			Pixxaro	Corteva Agriscience
MCPA ester - 600 g/L EC	fluroxypyr - 180 g/L			Predicade	FMC
MCPA ester - 600 g/L EC	MCPB - 375 g/L SL (F)				
MCPA K+ - 25 g/L SL (F)	MCPB - 375 g/L SL (F)			Trophy	Nufarm Agriculture
MCPA K+ - 25 g/L SL (F)	MCPB - 375 g/L SL (F)			Topside	Loveland Products
MCPA K+ - 336 g/L SL (F)	dicamba - 84 g/L SL (F)			Tropotox Plus	Nufarm Agriculture
MCPA K+ - 400 g/L SL				DyVel	BASF Canada
MCPA Na+ - 25 g/L SL (F)	MCPB - 375 g/L SL (F)			MCPA K+	Various
MCPA Na+ - 300 g/L SL				Glovitox Plus	IPCO
				MCPA Na	Various

(Component 1) Active Ingredient** - Formulation	(Component 2) Active Ingredient** - Formulation	(Component 3) Active Ingredient** - Formulation	(Component 4) Active Ingredient** - Formulation	Product	Company
MCPB - 375 g/L SL (F)	MCPA Na+ - 25 g/L SL (F)			<i>Clovitox Plus</i>	IPCO
MCPB - 375 g/L SL (F)	MCPA K+ - 25 g/L SL (F)			<i>Topside</i>	Loveland Products
MCPB - 375 g/L SL (F)	MCPA K+ - 25 g/L SL (F)			<i>Tropotox Plus</i>	Nufarm Agriculture
mecoprop-P - 130 g/L SL (F)	MCPA Amine - 160 g/L SL (F)	dichlorprop-P - 310 g/L SL (F)		<i>Optica Trio</i>	UAP
mecoprop-P - 150 g/L SL				<i>Mecoprop-P</i>	UAP
mecoprop-P - 62.5 g/L SL (F)	dicamba - 62.5 g/L SL (F)	MCPA amine - 275 g/L SL (F)		<i>Tracker XP</i>	IPCO
mecoprop-P - 62.5 g/L SL (F)	dicamba - 62.5 g/L SL (F)	MCPA amine - 275 g/L SL (F)		<i>Sword</i>	Loveland Products
mecoprop-P - 62.5 g/L SL (F)	dicamba - 62.5 g/L SL (F)	MCPA amine - 275 g/L SL (F)		<i>Target</i>	Syngenta Canada
mecoprop-P - 80 g/L SL (F)	dicamba - 110 g/L SL (F)	2,4-D Amine - 295 g/L SL (F)		<i>DyVelD5p</i>	BASF Canada
metolachlor - 400 g/L SC (F)	atrazine - 320 g/L SC (F)			<i>Primextra II Magnum</i>	Syngenta Canada
metolachlor - 915 g/L EC				<i>Dual II Magnum</i>	Syngenta Canada
metribuzin - 480 g/L SC				<i>Matrix SC</i>	Sharda CropChem
metribuzin - 70% WG				<i>Buzzin</i>	Sharda CropChem
metribuzin - 75% WG				<i>Squadron</i>	ADAMA Canada
metribuzin - 75% WG				<i>Sencor</i>	Bayer CropScience
metribuzin - 75% WG				<i>TriCor</i>	United Phosphorus Inc.
metribuzin - 75% WG				<i>Navius</i>	Bayer Environmental Sciences
metribuzin - 12.6% WG (F)	aminocyclopyrachlor - 39.5% WG (F)			<i>Travallas</i>	FMC
	thifensulfuron - 30 g/L SC (F)	fluroxypyr - 150 g/L SC (F)		<i>Ally Toss-N-Go</i>	FMC
metulfuron - 3 g/L SC (F)				<i>Escort</i>	Bayer Environmental Sciences
metulfuron - 60% SG					
metulfuron - 60% WG					
metulfuron - 8.6% SG (B)	tribenuron - 42.9% SG (B)			<i>MPower X-Pro</i>	Agracity
metulfuron - 8.6% SG (B)	tribenuron - 42.9% SG (B)			<i>Express Pro</i>	FMC
metulfuron - 9.45% WG (F)	aminopyralid - 52.5% WG (F)	2,4-D Ester - 660 g/L EC		<i>Reclaim II</i>	Corteva Agriscience
nicosulfuron - 37.5% WG (F)	rimsulfuron - 37.5% WG (F)			<i>Ultim</i>	Corteva Agriscience
nicosulfuron - 75% WG				<i>Accent</i>	Syngenta Canada
paraquat - 200 g/L SL				<i>Gramoxone</i>	Syngenta Canada
picloram - 240 g/L SL				<i>Tordon 22K</i>	Corteva Agriscience
picloram - 97.5 g/L SL (F)				<i>Grazon XC</i>	Syngenta Canada
pinoxaden - 25 g/L EC (F)				<i>Traxos</i>	Syngenta Canada
pinoxaden - 25 g/L EC (F1)				<i>TraxosTwo</i>	Corteva Agriscience
pinoxaden - 50 g/L EC				<i>Rezuvant</i>	Syngenta Canada
pinoxaden - 50 g/L EC				<i>Axial iPak</i>	Syngenta Canada
pinoxaden - 50 g/L EC (F)				<i>Axial Xtreme</i>	Syngenta Canada
pinoxaden - 50 g/L EC (F1)				<i>Axial Xtreme iPak</i>	Syngenta Canada
pinoxaden - 50 g/L EC (F)				<i>Axial</i>	Syngenta Canada
pinoxaden - 92.7 g/L EC (F)				<i>Broadband</i>	Syngenta Canada
propoxycarbazono - 70% WG				<i>Olympus</i>	Bayer CropScience
propyzamide - 400 g/L SC				<i>Kerb SC</i>	Corteva Agriscience
propyzamide - 50% WSP				<i>Kerb 50WSP</i>	Corteva Agriscience
pyraflufen - 13.5 g/L (F)				<i>Goldwing</i>	Nufarm Agriculture
pyraflufen - 6.1 g/L (F)				<i>BlackHawk (new)</i>	Nufarm Agriculture
pyrasulfotole - 15.5 g/L EC (F)				<i>Tundra</i>	Bayer CropScience
pyrasulfotole - 31.3 g/L EC (F)				<i>Velocity m3</i>	Bayer CropScience
pyrasulfotole - 37.5 g/L EC (F)				<i>Infinity</i>	Bayer CropScience
pyrasulfotole - 37.5 g/L EC (F)				<i>Infinity FX</i>	Bayer CropScience
pyrasulfotole - 37.5 g/L EC (F)				<i>Axial iPak</i>	Syngenta Canada
pyrasulfotole - 37.5 g/L EC (F)				<i>Axial Xtreme iPak</i>	Syngenta Canada
pyroxasulfone - 250 g/L SC (F)				<i>Authority Supreme</i>	FMC
pyroxasulfone - 44.5% WG (F)				<i>Fierce</i>	Valent Canada
pyroxasulfone - 447 g/L SE (F)				<i>Focus</i>	FMC
pyroxasulfone - 85% WG				<i>Focus (co-pack)</i>	FMC
pyroxasulfone - 15% WG (F)				<i>Resxade</i>	Corteva Agriscience

(Component 1) Active Ingredient** - Formulation	(Component 2) Active Ingredient** - Formulation	(Component 3) Active Ingredient** - Formulation	(Component 4) Active Ingredient** - Formulation	Product	Company
pyroxsulam - 21.5% WG pyroxsulam - 30 g/L OD pyroxsulam - 30 g/L SC quinclorac - 51.55% SG (B)	fluroxypyr - 333 g/L EC thifensulfuron:tribenuron - 10.3%:5.2% SG (B)			Simplicity GoDRI Tandem Simplicity OD Triton C	Corteva Agriscience Corteva Agriscience Corteva Agriscience FMC
quinclorac - 75% WDG quinclorac - 75% WDG quinclorac - 75% WDG quinclorac 180 g/L SN quizalofop-p - 96 g/L EC quizalofop-p - 96 g/L EC quizalofop-p - 96 g/L EC rimsulfuron - 20% WG rimsulfuron - 25% WG rimsulfuron - 37.5% WG (F) saflufenacil - 342 g/L SC saflufenacil - 70% SG sethoxydim - 450 g/L EC sethoxydim - 450 g/L EC sethoxydim - 450 g/L EC sethoxydim - 450 g/L EC simazine - 480 g/L SC simazine - 90% WG sulfentrazone - 250 g/L SC (F) sulfentrazone - 480 g/L SC sulfentrazone - 480 g/L SC thiencarbazone - 10 g/L SC thiencarbazone - 10 g/L SC thiencarbazone - 10 g/L SC	nicosulfuron - 37.5% WG (F) imazamox - 35% WG (F) imazamox - 35% WG (F) imazamox - 25 g/L SL pyroxa sulfone - 250 g/L SC (F) carfentrazone - 240 g/L EC tribenuron - 50% SG fluroxypyr - 333 g/L EC	imazethapyr - 35% WG (F) imazethapyr - 35% WG (F)		Clever Ingenious Masterline Quinclorac Facet L Assure II Yuma GL Contender Sortan IS Prism SG Ultim 75DF/Grande Heat LQ Heat WG Odyssey Ultra Odyssey Ultra NXT Poast Ultra Solo Ultra Simazine 480 Princep Nine T Authority Supreme Authority Authority Charge Luxxur Varro Predicade	Great Northern Growers UAP Univar Canada BASF Canada Corteva Agriscience Gowan Canada IPCO Corteva Agriscience Corteva Agriscience Corteva Agriscience BASF Canada BASF Canada BASF Canada BASF Canada BASF Canada Loveland Products Univar ProVMWeb FMC FMC FMC Bayer CropScience Bayer CropScience FMC
thiencarbazone - 5 g/L SC (F) thifensulfuron - 30 g/L SC (F) thifensulfuron - 50% SG thifensulfuron:tribenuron - 10.3%:5.2% SG (B) thifensulfuron:tribenuron - 33.3%:16.7% SG thifensulfuron:tribenuron - 33.3%:16.7% SG thifensulfuron:tribenuron - 33.3%:16.7% SG thifensulfuron:tribenuron - 33.3%:16.7% SG thifensulfuron:tribenuron - 33.3%:16.7% SG thifensulfuron:tribenuron - 50%:25% SG thifensulfuron:tribenuron - 50%:25% SG thifensulfuron:tribenuron - 50%:25% SG (F) thifensulfuron:tribenuron - 50%:25% SG (F)	pyrasulfotole - 31.3 g/L EC (F) fluroxypyr - 150 g/L SC (F) quinclorac - 51.55% SG (B) MCPA Ester - 500 or 600 g/L EC MCPA ester - 500 or 600 g/L EC fluroxypyr - 333 g/L EC fluroxypyr - 180 g/L EC bromoxynil - 200 g/L EC (F) clodinafop - 112 g/L (F)	thifensulfuron:tribenuron - 50%:50% SG bromoxynil - 175 g/L EC (F) metsulfuron - 3 g/L SC (F)	MCPA ester - 600 g/L EC	Velocity m3 Travallas Pinnacle SG Triton C Refine M Refine SG BroadSide Retain SG (new) Retain SG (old) Enforcer MSU Signal FSU MPower R Deploy	Bayer CropScience FMC FMC FMC FMC FMC Loveland Products Loveland Products Loveland Products Nufarm Agriculture Nufarm Agriculture Agricity Anysta LifeScience

(Component 1) Active Ingredient** - Formulation	(Component 2) Active Ingredient** - Formulation	(Component 3) Active Ingredient** - Formulation	(Component 4) Active Ingredient** - Formulation	Product	Company
thiencarbazono - 10 g/L SC	thiencarbazono - 10 g/L SC	MCPA ester - 600 g/L EC	thiencarbazono - 10 g/L SC	Predicade	FMC
thiencarbazono - 10 g/L SC	thiencarbazono - 10 g/L SC	MCPA ester - 600 g/L EC	thiencarbazono - 10 g/L SC	Barricade II	FMC
thiencarbazono - 10 g/L SC	thiencarbazono - 10 g/L SC	MCPA ester - 600 g/L EC	thiencarbazono - 10 g/L SC	Armezon Impact Shieldex 400SC	BASF Canada UAP ISK BioSciences/Engage
thiencarbazono - 10 g/L SC	thiencarbazono - 10 g/L SC	MCPA ester - 600 g/L EC	thiencarbazono - 10 g/L SC	Bison Achieve (Liquid Achieve) Marengo Nufarm Trailkoxydim Avadex MicroActiv Fortress MicroActiv Avadex Liquid EC Inferno Duo MPower X-Pro Express Pro Luxxur Express FX Express SG MPower Extra MPower X-Ko Inferno Ko-Act Spike Triton K Treflan MicroActiv Rival 10G Bonanza 10G Treflan Liquid EC Bonanza 480 Liquid Rival EC	Agro ADAMA Canada Corteva Agriscience Loveland Products Nufarm Agriculture Gowan Canada Gowan Canada Gowan Canada Arysta LifeScience Agracity FMC Bayer CropScience FMC FMC Agracity Agracity Arysta LifeScience Nufarm Agriculture Nufarm Agriculture FMC Gowan Canada Nufarm Agriculture UAP Gowan Canada UAP Nufarm Agriculture
thiencarbazono - 10 g/L SC	thiencarbazono - 10 g/L SC	MCPA ester - 600 g/L EC	thiencarbazono - 10 g/L SC	Zampro Trivapro Quilt Topnotch Quadris Top Elatus Azoshy 250SC Quadris Double Nickel/LC	BASF Canada Syngenta Canada Syngenta Canada ADAMA Canada Syngenta Canada Syngenta Canada Sharda Crop Chem Syngenta Canada UAP
thiencarbazono - 10 g/L SC	thiencarbazono - 10 g/L SC	MCPA ester - 600 g/L EC	thiencarbazono - 10 g/L SC	Double Nickel 155	UAP

Foliar Fungicides

(Component 1) Active Ingredient** - Formulation	(Component 2) Active Ingredient** - Formulation	(Component 3) Active Ingredient** - Formulation	(Component 4) Active Ingredient** - Formulation	Product	Company
ametoctradin - 300 g/L SC (F)	ametoctradin - 300 g/L SC (F)	ametoctradin - 300 g/L SC (F)	ametoctradin - 300 g/L SC (F)	Zampro	BASF Canada
azoxystrobin - 75 g/L SC (F)	azoxystrobin - 75 g/L SC (F)	azoxystrobin - 75 g/L SC (F)	azoxystrobin - 75 g/L SC (F)	Trivapro	Syngenta Canada
azoxystrobin - 75 g/L SC (F)	azoxystrobin - 75 g/L SC (F)	azoxystrobin - 75 g/L SC (F)	azoxystrobin - 75 g/L SC (F)	Quilt	Syngenta Canada
azoxystrobin - 143 g/L SC (F)	azoxystrobin - 143 g/L SC (F)	azoxystrobin - 143 g/L SC (F)	azoxystrobin - 143 g/L SC (F)	Topnotch	ADAMA Canada
azoxystrobin - 200 g/L SC (F)	azoxystrobin - 200 g/L SC (F)	azoxystrobin - 200 g/L SC (F)	azoxystrobin - 200 g/L SC (F)	Quadris Top	Syngenta Canada
azoxystrobin - 250 g/L SC	azoxystrobin - 250 g/L SC	azoxystrobin - 250 g/L SC	azoxystrobin - 250 g/L SC	Elatus	Syngenta Canada
azoxystrobin - 250 g/L SC	azoxystrobin - 250 g/L SC	azoxystrobin - 250 g/L SC	azoxystrobin - 250 g/L SC	Azoshy 250SC	Sharda Crop Chem
azoxystrobin - 250 g/L SC	azoxystrobin - 250 g/L SC	azoxystrobin - 250 g/L SC	azoxystrobin - 250 g/L SC	Quadris	Syngenta Canada
Bacillus amyloliquefaciens - 5x10 ¹⁰ spores/mL AS	Bacillus amyloliquefaciens - 5x10 ¹⁰ spores/mL AS	Bacillus amyloliquefaciens - 5x10 ¹⁰ spores/mL AS	Bacillus amyloliquefaciens - 5x10 ¹⁰ spores/mL AS	Double Nickel/LC	UAP
Bacillus amyloliquefaciens - 1x10 ¹⁰ spores/mL AS	Bacillus amyloliquefaciens - 1x10 ¹⁰ spores/mL AS	Bacillus amyloliquefaciens - 1x10 ¹⁰ spores/mL AS	Bacillus amyloliquefaciens - 1x10 ¹⁰ spores/mL AS	Double Nickel 155	UAP

(Component 1) Active Ingredient** - Formulation	(Component 2) Active Ingredient** - Formulation	(Component 3) Active Ingredient** - Formulation	(Component 4) Active Ingredient** - Formulation	Product	Company
mancozeb - 75% WG				Manzate Pro-Stick	United Phosphorus
mancozeb - 75% WG				Penncozeb 75DF	United Phosphorus
mandipropamid - 250 g/L SC				Revus	Syngenta Canada
Mandipropamid - 250 g/L SC	oxathiopiprolin - 30 g/L SC			Orondis Ultra	Syngenta Canada
metalaxyl-M - 480 g/L EC				Ridomil Gold 480 EC	Syngenta Canada
metalaxyl-M - 480 g/L SL				Ridomil Gold 480 SL	Syngenta Canada
metalaxyl-M - 480 g/L EC (DC)	chlorothalonil - 500 g/L SC (DC)			Ridomil Gold / Bravo	Syngenta Canada
metalaxyl-M - 480 g/L SL (DC)	chlorothalonil - 500 g/L SC (DC)			Ridomil Gold SL / Bravo	Syngenta Canada
metconazole - 90 g/L EC				Caramba	BASF Canada
metconazole - 50% WG				Quash	Valent Canada
metconazole - 80 g/L EC (F)	pyraclostrobin - 130 g/L EC (F)			Twinline	BASF Canada
metiram - 55% WG (F)	pyraclostrobin - 5% WG (F)			Cabrio Plus	BASF Canada
metiram - 80% WG				Polyram DF	BASF Canada
mono/di-potassium salts of phosphorous acid - 53% SL				Confine Extra	Winfield Solutions
mono/di-potassium salts of phosphorous acid - 53% SL					
oxathiopiprolin - 30 g/L SC	Mandipropamid - 250 g/L SC			Rampart	Loveland Products
penthiopyrad - 200 g/L SC					Canada
penthiopyrad - 200 g/L EC				Orondis Ultra	Syngenta Canada
picoxystrobin - 250 g/L SC				Fontelis	Corteva Agriscience
propiconazole - 125 g/L EC (F)				Vertisan	Division of DowDuPont
propiconazole - 250 g/L EC				Acapela	Corteva Agriscience
propiconazole - 250 g/L EC				Nexicor	Division of DowDuPont
propiconazole - 250 g/L EC	fluxapyroxad - 30 g/L EC (F)			Tilt 250E	BASF Canada
propiconazole - 250 g/L EC				Propel	Syngenta Canada
propiconazole - 250 g/L EC				Propi Super 25 EC	Syngenta Canada
propiconazole - 432 g/L EC				Bumper 432 EC	Sharda CropChem
propiconazole - 418 g/L EC				Co-Op Pivot	ADAMA Canada
propiconazole - 418 g/L EC				Nufarm Propiconazole	Federated Co-operatives Limited
propiconazole - 418 g/L EC				Fitness	Nufarm Agriculture
propiconazole - 418 g/L EC					Loveland Products
propiconazole - 418 g/L EC				Pivot 418EC	Canada
propiconazole - 125 g/L SC (F)				Quilt	IPC0
propiconazole - 124 g/L SC (F)				Topnotch	Syngenta Canada
propiconazole - 125g/L SC (F)				Trivapro	ADAMA Canada
prothioconazole - 480 g/L SC	benzovindiflupyr - 100 g/L EC			Proline 480SC	Syngenta Canada
prothioconazole - 150 g/L SC (F)				Cotegra	Bayer
prothioconazole - 125 g/L EC (F)				Prosaro 250EC	BASF Canada
prothioconazole - 125 g/L EC (F)				Prosaro XTR	Bayer
prothioconazole - 175 g/L SC (F)				Delaro 325 SC	Bayer
prothioconazole - 200 g/L SC (F)				Propulse	Bayer
pyraclostrobin - 5% WG (F)				Cabrio Plus	BASF Canada
pyraclostrobin - 130 g/L EC (F)				Twinline	BASF Canada
pyraclostrobin - 200 g/L EC (F)	fluxapyroxad - 30 g/L EC (F)			Nexicor	BASF Canada
pyraclostrobin - 250 g/L EC				Headline EC	BASF Canada
pyraclostrobin - 250 g/L EC				MPower Spade	BASF Canada
pyraclostrobin - 250 g/L EC				Lance AG	BASF Canada
pyraclostrobin - 250 g/L SC (F)				Dyax	BASF Canada
pyraclostrobin - 333 g/L SC (F)				Priaxor	BASF Canada
pyrimethanil - 400 g/L SC				Scala SC	BASF Canada
pyrimethanil - 375 g/L SC (F)	fluopyram - 125 g/L SC (F)			Luna Tranquility	Bayer



(Component 1) Active Ingredient* - Formulation	(Component 2) Active Ingredient* - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
<i>Reynoutria sachalinensis</i> - 20% SC sodium/potassium/ammonium phosphites - 53.6% SL sulphur - 80% WG				<i>Regalia Maxx</i> <i>Phostrol</i>	Marrone Bio Innovations Engage Agro
tebuconazole - 250 g/L EW tebuconazole - 432 g/L SC tebuconazole - 432 g/L SC tebuconazole - 125 g/L EC (F) tebuconazole - 125 g/L EC (F) trifloxystrobin - 150 g/L SC (F) zoxamide - 8.43% WG (F)	prothioconazole - 125 g/L EC (F) prothioconazole - 125 g/L EC (F) prothioconazole - 175 g/L SC (F) mancozeb - 66.7% WG (F)			<i>Cosavet DF Edge</i> <i>Folicur 250EW</i> <i>Hornet 432 F</i> <i>Palliser</i> <i>Prosaro 250EC</i> <i>Prosaro XTR</i> <i>Delaro 325 SC</i> <i>Gavel 75 DF</i>	Engage Agro Bayer Nufarm Agriculture Inc. Bayer Bayer Bayer Bayer Gowan Canada

Seed Treatments

(Component 1) Active Ingredient* - Formulation	(Component 2) Active Ingredient* - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
azoxystrobin - 143 g/L SC azoxystrobin - 1.33% FS azoxystrobin - 1.33% FS	fludioxonil - 143 g/L SC fludioxonil - 3.32% FS fludioxonil - 3.32% FS	difenoconazole - 112 g/L SC metalaxyl-m+s - 2.65% FS metalaxyl-m+s - 2.65% FS	thiabendazole - 26.5% FS thiabendazole: thiamethoxam - 26.5%: 47.6% FS	<i>Stadium</i> <i>Maxim Quattro</i> <i>Cruiser Maxx Corn</i>	Syngenta Canada Syngenta Canada Syngenta Canada
<i>Bacillus subtilis</i> - 26.2% WP captan - 30% FS carbathiin - 87.5 g/L FS	ipconazole - 9.38 g/L FS	ipconazole - 5.0 g/L FS		<i>Serenade OPTI</i> <i>Agrox FL</i> <i>Rancona VRS</i>	Bayer Norac Concepts Arysta LifeScience Canada
carbathiin - 133.33 g/L FS	metalaxyl - 13.33 g/L FS			<i>Rancona Trio</i>	Arysta LifeScience Canada
carbathiin - 15.59% FS	thiram - 13.25% FS			<i>Vitaflor 280</i>	Arysta LifeScience Canada
carbathiin - 15.59% FS carbathiin - 47.6 g/L FS clothianidin - 600 g/L FS	thiram - 13.25% FS thiram - 13.25% FS thiram - 95.3 g/L FS	imidacloprid - 285.7 g/L FS		<i>Vitaflor SP Fungicide</i> <i>Vitaflor Fungicide</i> <i>Gaoucho CSFL</i> <i>Nipsit INSIDE 600</i>	IPCO IPCO Bayer Valent Canada
clothianidin - 600 g/L FS clothianidin - 600 g/L FS clothianidin - 30.7 g/L FS	metalaxyl - 9.24 g/L FS	metconazole - 4.92 g/L FS		<i>Insecticide</i> <i>Poncho 600FS</i> <i>Titan</i> <i>Nipsit SUITE Cereals OF</i> <i>Seed Protectant</i>	BASF Canada BASF Canada Valent Canada
clothianidin - 290 g/L FS clothianidin - 600 g/L FS cyantraniliprole - 200 g/L FS	metalaxyl - 7.15 g/L FS penflufen - 100 g/L FS	trifloxystrobin - 7.15 g/L FS prothioconazole - 18 g/L FS	penflufen - 10.7 g/L FS	<i>Prosper EverGol</i> <i>Titan Erresto</i> <i>Verimark</i>	Bayer Bayer Corteva Agriscience Division of DowDuPont
cyantraniliprole - 600 g/L FS cyantraniliprole - 625 g/L FS				<i>Fortenza</i> <i>Lumiderm</i>	Syngenta Canada Corteva Agriscience
difenoconazole - 19.4 g/L FS difenoconazole - 112 g/L SC difenoconazole - 77.2 g/L FS difenoconazole - 36.8 g/L FS	fludioxonil - 19.4 g/L FS fludioxonil - 143 g/L SC mandipropamid - 154.3 g/L FS fludioxonil - 7.6 g/L FS	azoxystrobin - 143 g/L SC sedaxane - 77.2 g/L FS metalaxyl-m+s - 9.2 g/L FS	sedaxane - 15.4 g/L FS	<i>Maxim D</i> <i>Stadium</i> <i>Vibrance Ultra Potato</i> <i>Vibrance Quattro</i>	Division of DowDuPont Syngenta Canada Syngenta Canada Syngenta Canada Syngenta Canada

(Component 1) Active Ingredient* - Formulation	(Component 2) Active Ingredient* - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
difenoconazole - 123 g/L FS	fludioxonil - 62.5 g/L FS	thiamethoxam - 250 g/L FS	metalaxyl-	Cruiser Maxx Potato Extreme	Syngenta Canada
difenoconazole - 16 g/L FS	fludioxonil - 1.7 g/L FS	thiamethoxam - 269 g/L FS	m+s:sedaxane-5:3.4 g/L FS	Helix Vibrance	Syngenta Canada
difenoconazole - 36.9 g/L FS	fludioxonil - 7.7 g/L FS	thiamethoxam - 61.5 g/L FS	metalaxyl-m+s: sedaxane-9:2: 15.4 g/L FS	Cruiser Vibrance Quattro	Syngenta Canada
difenoconazole 16 g/L FS	metalaxyl-m+s - 5 g/L FS	thiamethoxam - 269 g/L FS	fludioxonil: sedaxane- 1.7 g/L : 3.4 g/L FS	Visivio	Syngenta Canada
dimethyl benzyl ammonium chloride - 10% Liquid				General Storage	Ag-Services
ethaboxam - 383 g/L FS	fludioxonil - 25 g/L FS	metalaxyl-m+s - 37.5 g/L	sedaxane - 50 g/L FS	Disinfectant	Valent Canada
ethaboxam - 383 g/L FS				INTEGO Solo Fungicide Vibrance Maxx with	Syngenta Canada
fenamidone - 500g/L SC				INTEGO Seed Treatment	
fludioxonil - 0.5% DS				Reason 500 SC	Bayer
fludioxonil - 19.4 g/L FS	difenoconazole - 19.4 g/L FS			Maxim PSP	Syngenta Canada
fludioxonil - 143 g/L SC	difenoconazole - 112 g/L SC	azoxystrobin - 143 g/L SC		Maxim D	Syngenta Canada
fludioxonil - 62.5 g/L FS	difenoconazole - 123 g/L FS	thiamethoxam - 250 g/L FS		Stadium	Syngenta Canada
				Cruiser Maxx Potato	Syngenta Canada
fludioxonil - 0.5% DS	mancozeb - 5.7% DS			Extreme	
fludioxonil - 0.73% FS	metalaxyl-m+s - 1.10% FS			Maxim MZ PSP	Syngenta Canada
fludioxonil - 25 g/L FS	metalaxyl-m+s - 37.5 g/L FS	sedaxane - 50 g/L FS		Apron Maxx RTA	Syngenta Canada
fludioxonil - 0.73% FS	metalaxyl-m+s - 1.10% FS	sedaxane - 500 g/L FS		Vibrance Maxx RFC	Syngenta Canada
fludioxonil - 7.6 g/L FS	metalaxyl-m+s - 9.2 g/L FS	sedaxane - 15.4 g/L FS	difenoconazole - 36.8 g/L FS	Vibrance Maxx RTA	Syngenta Canada
fludioxonil - 25 g/L FS	metalaxyl-m+s - 37.5 g/L FS	sedaxane - 50 g/L FS	ethaboxam - 383 g/L FS	Vibrance Quattro	Syngenta Canada
				Vibrance Maxx RFC with INTEGO Seed Treatment	Syngenta Canada
fludioxonil - 25 g/L FS	metalaxyl-m+s - 20 g/L FS	thiabendazole - 150 g/L FS		Apron Advance	Syngenta Canada
fludioxonil - 3.32% FS	metalaxyl-m+s - 2.65% FS	thiabendazole - 26.5% FS		Maxim Quattro	Syngenta Canada
fludioxonil - 1.12% FS	metalaxyl-m+s - 1.7% FS	thiamethoxam - 22.6% FS		Cruiser Maxx Beans	Syngenta Canada
fludioxonil - 3.32% FS	metalaxyl-m+s - 2.65% FS	thiamethoxam - 47.6% FS		Cruiser Maxx Corn	Syngenta Canada
fludioxonil - 0.73% FS	metalaxyl-m+s - 1.1% FS	thiamethoxam - 47.6%		Cruiser Maxx Vibrance Pulses	Syngenta Canada
				Beans	
fludioxonil - 1.12% FS	metalaxyl-m+s - 1.7% FS	thiamethoxam - 22.6% FS		Helix Vibrance	Syngenta Canada
fludioxonil - 1.7 g/L FS	metalaxyl-m+s - 5 g/L FS	thiamethoxam - 269 g/L FS		Cruiser Vibrance Quattro	Syngenta Canada
fludioxonil - 7.7 g/L FS	metalaxyl-m+s - 9.2 g/L FS	thiamethoxam - 61.5 g/L FS		Visivio	Syngenta Canada
fludioxonil - 1.7 g/L FS	metalaxyl-m+s - 5 g/L FS	thiamethoxam - 269 g/L FS			
fluxapyroxad - 17 g/L FS	metalaxyl - 10 g/L FS	pyraclostrobin - 17 g/L FS		Insure Pulse	BASF Canada
fluxapyroxad - 8.35g/L FS	pyraclostrobin - 16.7 g/L FS	triticonazole - 16.7 g/L FS	metalaxyl - 10 g/L FS	Insure Cereal FX4	BASF Canada
hydrogen peroxide - 27% LS				StorOx	Biosafe Systems
imidacloprid - 240 g/L FS				Admire SPT	Bayer
imidacloprid - 240 g/L FS				Alias 240SC	ADAMA Canada
imidacloprid - 600 g/L FS				Sombriero 600FS	ADAMA Canada
imidacloprid - 600 g/L FS				Stress Shield 600	Bayer
imidacloprid - 285.7 g/L FS	carbathiin - 47.6 g/L FS	thiram - 95.3 g/L FS	prothioconazole-15.4 g/L FS	Gaucha CSFL	Bayer
imidacloprid - 600 g/L FS	metalaxyl - 6.2 g/L FS	tebuconazole - 3.0 g/L FS	trifloxystrobin - 154 g/L FS	Raxil PRO Shield	Bayer CropScience
imidacloprid - 600 g/L FS	metalaxyl - 317 g/L FS	penflufen - 154 g/L FS		Triflex EverGo! Shield	Bayer

(Component 1) Active Ingredient* - Formulation	(Component 2) Active Ingredient* - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
ipiconazole - 9.38 g/L FS	carbathiin - 87.5 g/L FS			Rancona VRS	Arysta LifeScience Canada
ipiconazole - 5.0 g/L FS	carbathiin - 133.33 g/L FS	metalaxyl - 13.33 g/L FS		Rancona Trio	Arysta LifeScience Canada
ipiconazole - 4.61 g/L FS	metalaxyl - 6.15 g/L FS			Cover 2	Loveland Products Canada Inc.
ipiconazole - 4.61 g/L FS	metalaxyl - 6.15 g/L FS			Rancona Pinnacle	Arysta LifeScience Canada
mancozeb - 16% DS				Solan MZ	Norac Concepts
mancozeb - 16% DS				Tube Seal	Norac Concepts
mancozeb - 16% DS				Potato ST16	Wilbur-Ellis Co.
mancozeb - 5.7% DS	fludioxonil - 0.5% DS			Maxim MZ PSP	Syngenta Canada
mandipropamid - 250 g/L FS				Revus	Syngenta Canada
mandipropamid - 154.3 g/L FS	difenoconazole - 77.2 g/L FS	sedaxane - 77.2 g/L FS		Vibrance Ultra Potato	Syngenta Canada
metalaxyl - 317 g/L FS				Allegiance FL	Bayer
metalaxyl - 317 g/L FS				Belmont 2.7FS	Arysta LifeScience Canada
metalaxyl - 317 g/L FS	ipiconazole - 4.61 g/L FS			Trilex Component B	Bayer
metalaxyl - 6.15 g/L FS	ipiconazole - 4.61 g/L FS			Cover 2	Loveland Products Canada Inc.
metalaxyl - 6.15 g/L FS	ipiconazole - 4.61 g/L FS			Rancona Pinnacle	Arysta LifeScience Canada
metalaxyl - 13.33 g/L FS	ipiconazole - 5.0 g/L FS	carbathiin - 133.33 g/L FS		Rancona Trio	Arysta LifeScience Canada
metalaxyl - 46.5 g/L FS	metconazole 23.2 g/L FS			Metlock CT	Valent Canada
metalaxyl - 9.24 g/L FS	metconazole - 4.92 g/L FS	clothianidin - 30.7 g/L FS		Nipsit SUITE Cereals OF Seed Protectant	Valent Canada
metalaxyl - 61.4 g/L FS	prothioconazole - 76.8 g/L FS	penflufen - 38.4 g/L FS		EverGol Energy	Bayer
metalaxyl - 10 g/L FS	pyraclostrobin - 17 g/L FS	triticonazole - 17 g/L FS		Insure Cereal	BASF Canada
metalaxyl - 10 g/L FS	pyraclostrobin - 16.7 g/L FS	triticonazole - 16.7 g/L FS	fluzapyroxad - 8.35 g/L FS	Insure Cereal FX4	BASF Canada
metalaxyl - 10 g/L FS	pyraclostrobin - 17 g/L FS	fluzapyroxad - 17 g/L FS		Insure Pulse	BASF Canada
metalaxyl - 7.15 g/L FS	trifloxystrobin - 7.15 g/L FS	clothianidin - 290 g/L FS		Prosper EverGol	Bayer
metalaxyl - 317 g/L FS	trifloxystrobin - 154 g/L FS	penflufen - 154 g/L FS		Trilex EverGol	Bayer
metalaxyl - 317 g/L FS	trifloxystrobin - 154 g/L FS	penflufen - 154 g/L FS		Trilex EverGol Shield	Bayer
metalaxyl - 6.6 g/L FS	tebuconazole - 5 g/L FS			Raxil MD	Bayer
metalaxyl - 6.2 g/L FS	tebuconazole - 3.0 g/L FS	imidacloprid - 600 g/L FS		Raxil PRO Shield	Bayer CropScience
metalaxyl - 6.2 g/L FS	tebuconazole - 3.0 g/L FS	prothioconazole - 15.4 g/L FS		Raxil PRO	Bayer
metalaxyl-m+s - 1.10% FS	fludioxonil - 0.73% FS			Apron Maxx RTA	Syngenta Canada
metalaxyl-m+s - 37.5 g/L FS	fludioxonil - 25 g/L FS	sedaxane - 50 g/L FS		Vibrance Maxx RFC	Syngenta Canada
metalaxyl-m+s - 1.10% FS	fludioxonil - 0.73% FS	sedaxane - 500 g/L FS		Vibrance Maxx RTA	Syngenta Canada
metalaxyl-m+s - 9.2 g/L FS	fludioxonil - 7.6 g/L FS	sedaxane - 15.4 g/L FS	difenoconazole - 36.8 g/L FS	Vibrance Quattro	Syngenta Canada
metalaxyl-m+s - 37.5 g/L FS	fludioxonil - 25 g/L FS	sedaxane - 50 g/L FS	ethaboxam - 383 g/L FS	Vibrance Maxx RFC with INTEGO Seed Treatment	Syngenta Canada Valent Canada
metalaxyl-m+s - 20 g/L FS	fludioxonil - 25 g/L FS	thiabendazole - 150 g/L FS		Apron Advance	Syngenta Canada
metalaxyl-m+s - 2.65% FS	fludioxonil - 3.32% FS	thiabendazole - 26.5% FS		Maxim Quattro	Syngenta Canada
metalaxyl-m+s - 1.7% FS	fludioxonil - 1.12% FS	thiamethoxam - 22.6% FS		Cruiser Maxx Beans	Syngenta Canada
metalaxyl-m+s - 2.65% FS	fludioxonil - 3.32% FS	thiamethoxam - 47.6% FS		Cruiser Maxx Corn	Syngenta Canada
metalaxyl-m+s - 5 g/L FS	fludioxonil - 1.7 g/L FS	thiamethoxam - 269 g/L FS	sedaxane: difenoconazole - 1.33%; 26.5% FS	Helix Vibrance	Syngenta Canada
metalaxyl-m+s - 9.2 g/L FS	fludioxonil - 7.7 g/L FS	thiamethoxam - 61.5 g/L FS	sedaxane: difenoconazole - 3.4; 16 g/L FS	Cruiser Vibrance Quattro	Syngenta Canada

(Component 1) Active Ingredient* - Formulation	(Component 2) Active Ingredient* - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
metalaxyl-m+s - 5 g/L FS	fludioxonil - 1.7 g/L FS	thiamethoxam - 269 g/L FS	difenoconazole: sedaxane - 16 g/L; 3.4 g/L FS sedaxane - 500 g/L FS	Visivio	Syngenta Canada
metalaxyl-m+s - 1.1% FS	fludioxonil - 0.73% FS	thiamethoxam - 47.5% FS		Cruiser Maxx Vibrance Pulses	Syngenta Canada
metalaxyl-m+s - 1.7% FS	fludioxonil - 1.12% FS	thiamethoxam - 22.6% FS	sedaxane - 500 g/L FS	Cruiser Maxx Vibrance Pulses	Syngenta Canada
metconazole - 23.2 g/L FS	metalaxyl - 46.5 g/L FS			Metlock CT	Valent Canada
metconazole - 4.92 g/L FS	metalaxyl - 9.24 g/L FS	clothianidin - 30.7 g/L FS		Nipsit SUITE Cereals OF Seed Protectant Confine Extra	Valent Canada
mono/di-potassium salts of phosphorous acid - 53% SC				Rampart	Winfield Solutions
mono/di-potassium salts of phosphorous acid - 53% SC				Lumisena	Loveland Products Canada
Oxathiapiprolin - 200g/L FS				Prosper EverGol Trilex EverGol	Division of DowDuPont
penflufen - 10.7 g/L FS	metalaxyl - 7.15 g/L FS	trifloxystrobin - 7.15 g/L FS	clothianidin - 290 g/L FS	Trilex EverGol Trilex EverGol Shield	Bayer
penflufen - 154 g/L FS	metalaxyl - 317 g/L FS	trifloxystrobin - 154 g/L FS	imidacloprid - 600 g/L FS	Ernesto Silver	Bayer
penflufen - 154 g/L FS	metalaxyl - 317 g/L FS	trifloxystrobin - 154 g/L FS		EverGol Energy Trilex Component A	Bayer
penflufen - 100 g/L FS	prothioconazole - 18 g/L FS	clothianidin - 600 g/L FS		EverGol Energy Raxil PRO	Bayer
penflufen - 100 g/L FS	prothioconazole - 18 g/L FS	metalaxyl - 61.4 g/L FS		Raxil PRO Ernesto Silver	Bayer
penflufen - 38.4 g/L FS	prothioconazole - 76.8 g/L FS	penflufen - 38.4 g/L FS	imidacloprid - 600 g/L FS	Raxil PRO Ernesto Silver	Bayer
penflufen - 154 g/L FS	trifloxystrobin - 154 g/L FS	tebuconazole - 3.0 g/L FS		Titan Ernesto	Bayer
prothioconazole - 76.8 g/L FS	metalaxyl - 61.4 g/L FS	tebuconazole - 3.0 g/L FS		Insure Cereal	BASF Canada
prothioconazole - 15.4 g/L FS	metalaxyl - 6.2 g/L FS	tebuconazole - 3.0 g/L FS		Insure Pulse	BASF Canada
prothioconazole - 15.4 g/L FS	metalaxyl - 6.2 g/L FS			Insure Cereal FX4	BASF Canada
prothioconazole - 18 g/L FS	penflufen - 100 g/L FS			Heads Up Plant Protectants	Heads Up Plant Protectants
prothioconazole - 18 g/L FS	penflufen - 100 g/L FS			Vibrance 500FS	Syngenta Canada
pyraclostrobin - 17 g/L FS	metalaxyl - 10 g/L FS			Vibrance Quattro	Syngenta Canada
pyraclostrobin - 17 g/L FS	metalaxyl - 10 g/L FS			Helix Vibrance	Syngenta Canada
pyraclostrobin - 17 g/L FS	fluxapyroxad 8:35 g/L FS			Cruiser Vibrance Quattro	Syngenta Canada
pyraclostrobin - 16.7 g/L FS	fluxapyroxad 8:35 g/L FS			Visivio	Syngenta Canada
saponins of <i>Chenopodium quinoa</i> - 63.02% WS				Vibrance Maxx RFC Vibrance Maxx RTA	Syngenta Canada
sedaxane - 500 g/L FS				Vibrance Maxx RFC with INTEGO Seed Treatment	Syngenta Canada
sedaxane - 15.4 g/L FS	metalaxyl-m+s - 9.2 g/L FS	difenoconazole - 36.8 g/L FS	ethaboxam - 383 g/L FS	Vibrance Ultra Potato Cruiser Maxx	Syngenta Canada
sedaxane - 3.4 g/L FS	metalaxyl-m+s - 5 g/L FS	difenoconazole - 16 g/L FS		Vibrance Pulses	Syngenta Canada
sedaxane - 15.4 g/L FS	metalaxyl-m+s - 9.2 g/L FS	difenoconazole - 36.9 g/L FS		Cruiser Maxx	Syngenta Canada
sedaxane - 15.4 g/L FS	metalaxyl-m+s - 5 g/L FS	thiamethoxam - 269 g/L FS		Vibrance Pulses	Syngenta Canada
sedaxane - 15.4 g/L FS	metalaxyl-m+s - 5 g/L FS	thiamethoxam - 269 g/L FS		Vibrance Beans	Syngenta Canada
sedaxane - 3.4 g/L FS	fludioxonil - 1.7 g/L FS			Rascendo Visivio	Syngenta Canada
sedaxane - 50 g/ L FS	metalaxyl-m+s - 37.5 g/L FS	fludioxonil - 25 g/L FS			Syngenta Canada
sedaxane - 500 g/L FS	metalaxyl-m+s - 1.10% FS	fludioxonil - 0.73% FS			Syngenta Canada
sedaxane - 50 g/L FS	metalaxyl-m+s - 37.5 g/L FS	fludioxonil - 25 g/L FS			Syngenta Canada
sedaxane - 77.2 g/L FS	mandipropamid - 154.3 g/L FS	difenoconazole - 77.2 g/L FS			Syngenta Canada
sedaxane - 500 g/L FS	metalaxyl-m+s - 1.1% FS	fludioxonil - 0.73% FS			Syngenta Canada
sedaxane - 500 g/L FS	metalaxyl-m+s - 1.7% FS	fludioxonil - 1.12% FS			Syngenta Canada
sulfoxaflor - 500 g/L FS		fludioxonil: metalaxyl- m+s-1.7:5 g/L FS			Syngenta Canada
sulfoxaflor - 500 g/L FS	difenoconazole - 16 g/L FS				Syngenta Canada

(Component 1) Active Ingredient* - Formulation	(Component 2) Active Ingredient* - Formulation	(Component 3) Active Ingredient* - Formulation	(Component 4) Active Ingredient* - Formulation	Product	Company
tebuconazole - 5 g/L FS	metalaxyl - 6.6 g/L FS	imidacloprid - 600 g/L FS	prothioconazole - 15.4 g/L FS	Raxil MD	Bayer
tebuconazole - 3.0 g/L FS	metalaxyl - 6.2 g/L FS	prothioconazole - 15.4 g/L FS		Raxil PRO Shield	Bayer
tebuconazole - 3.0 g/L FS	metalaxyl - 6.2 g/L FS			Raxil PRO	Bayer
thiabendazole - 500 g/L SC				Mertect SC	Syngenta Canada
thiabendazole - 150 g/L FS	metalaxyl-m+s - 20 g/L FS	fludioxonil - 25 g/L FS	azoxystrobin - 1.33% FS	Apron Advance	Syngenta Canada
thiabendazole - 26.5% FS	metalaxyl-m+s - 2.65% FS	fludioxonil - 3.32% FS	azoxystrobin:	Maxim Quattro	Syngenta Canada
thiabendazole - 26.5% FS	metalaxyl-m+s - 2.65% FS	fludioxonil - 3.32% FS	thiamethoxam - 1.33%; 47.6% FS	Cruiser Maxx Corn	Syngenta Canada
thiamethoxam - 240 g/L FS				Actara 240SC	Syngenta Canada
thiamethoxam - 47.6% FS				Cruiser 5FS	Syngenta Canada
thiamethoxam - 250 g/L FS	fludioxonil - 62.5 g/L FS	difenoconazole - 123 g/L FS		Cruiser Maxx	Syngenta Canada
thiamethoxam - 22.6% FS	fludioxonil - 1.12% FS	metalaxyl-m+s - 1.7% FS	azoxystrobin: thiabendazole	Potato Extreme	Syngenta Canada
thiamethoxam - 47.6% FS	fludioxonil - 3.32% FS	metalaxyl-m+s - 2.65% FS	- 1.33;26.5% FS	Cruiser Maxx Beans	Syngenta Canada
thiamethoxam - 22.6% FS	fludioxonil - 1.12% FS	metalaxyl-m+s - 1.7% FS	sedaxane - 500 g/L FS	Cruiser Maxx Corn	Syngenta Canada
thiamethoxam - 47.6% FS	fludioxonil - 0.73% FS	metalaxyl-m+s - 1.1% FS		Cruiser Maxx	Syngenta Canada
thiamethoxam - 61.5 g/L FS	fludioxonil - 7.7 g/L FS	metalaxyl-m+s - 9.2 g/L FS	sedaxane: difenoconazole - 15.4;36.9 g/L FS	Vibrance Beans	Syngenta Canada
thiamethoxam - 269 g/L FS	fludioxonil - 1.7 g/L FS	metalaxyl-m+s - 5 g/L FS	sedaxane: difenoconazole - 3.4;16 g/L FS	Cruiser Maxx	Syngenta Canada
thiamethoxam - 269 g/L FS	fludioxonil - 1.7 g/L FS	sedaxane - 3.4 g/L FS	difenoconazole: metalaxyl m+s - 16 g/L; 5g/L FS	Vibrance Pulses	Syngenta Canada
thiophanate-methyl - 10% DS				Cruiser Vibrance Quattro	Syngenta Canada
thiram - 13.25% FS	carbathiin - 15.59% FS			Helix Vibrance	Syngenta Canada
thiram - 13.25% FS	carbathiin - 15.59% FS			Visivio	Syngenta Canada
thiram - 95.3 g/L FS	carbathiin - 15.59% FS			Senator PSPT	Nippon Soda Company
trifloxystrobin - 7.15 g/L FS	carbathiin - 47.6 g/L FS	imidacloprid - 285.7 g/L FS		Vitafla 280	Arysta LifeScience
trifloxystrobin - 154 g/L FS	metalaxyl - 7.15 g/L FS	clothianidin - 290 g/L FS		Vitafla SP Fungicide	Canada
trifloxystrobin - 154 g/L FS	metalaxyl - 317 g/L FS	penflufen - 154 g/L FS	penflufen - 10.7 g/L FS	Vitafla Fungicide	IPCO
triconazole - 17 g/L FS	penflufen - 154 g/L FS	metalaxyl - 10 g/L FS		Gaucho CS FL	IPCO
triconazole - 16.7 g/L FS	fluxapyroxad - 8.35g/L FS	metalaxyl - 10 g/L FS		Prosper EverGol	Bayer
		metalaxyl - 10 g/L FS		Trilex EverGol	Bayer
		pyraclostrobin - 16.7 g/L FS		Trilex Component A	Bayer
				Insure Cereal	BASF
				Insure Cereal FX4	BASF Canada

Insecticides

Product	Active Ingredient	Formulation	Company
<i>Actara 240 SC/Actara 25 WG</i>	thiamethoxam	240 g/L SC, 25% WG	Syngenta Canada
<i>Admire 240 / SPT</i>	imidacloprid	240 g/L SC	Bayer CropScience
<i>Alias 240 SC</i>	imidacloprid	240 g/L SC	Loveland Products Canada
<i>Ambush</i>	permethrin	500 g/L EC	Amvac Chemical Corp.
<i>Assail</i>	acetamiprid	70% WP	Engage Agro / Nippon Soda Co. Ltd.
<i>Beleaf</i>	flonicamid	50% WP	FMC Corporation
<i>Capture</i>	bifenthrin	240 g/L	FMC Corporation
<i>Citadel 480EC</i>	chlorpyrifos	480 g/L EC	IPCO
<i>Closer</i>	sulfoxaflor	240 g/L	Dow AgroSciences
<i>Clutch</i>	clothianidin	50% WG	Valent Canada Inc.
<i>Concept</i>	imidacloprid: deltamethrin	75 g/L SC : 10 g/L SC	Bayer CropScience
<i>Coragen</i>	chlorantraniliprole	200 g/L SC	FMC Corporation
<i>Cygon 480EC/Cygon 480-Ag</i>	dimethoate	480 g/L EC	IPCO, FMC Corporation
<i>Decis</i>	deltamethrin	50 g/L EC	Bayer CropScience
<i>Delegate</i>	spinetoram	25% SG	Dow AgroSciences
<i>Dibrom</i>	naled	864 g/L EC	Loveland Products Canada
<i>Dipel 2X DF</i>	<i>Bacillus thuringiensis</i>	32 billion CLU/kg WG	Valent BioSciences
<i>Eco Bran</i>	carbaryl	2% spreadable bran bait	Peacock Industries
<i>Entrust</i>	spinosad	80% WP	Dow AgroSciences
<i>Fulfill</i>	pymetrozine	50% WG	Syngenta Canada
<i>Imidan</i>	phosmet	70% WP	Gowan Canada
<i>Insecto</i>	diatomaceous earth	90%	Natural Insecto Products Inc.
<i>Lagon 480E</i>	dimethoate	480 g/L EC	Loveland Products Canada
<i>Lannate</i>	methomyl	90% SP	E. I. duPont Canada
<i>Lorsban 4E</i>	chlorpyrifos	480 g/L EC	Dow AgroSciences
<i>Mako</i>	cypermethrin	407 g/L EC	Engage Agro Corp.
<i>Malathion 85E / 500</i>	malathion	85%, 500 g/L EC	Loveland Products Canada, IPCO
<i>Matador</i>	lambda-cyhalothrin	120 g/L EC	Syngenta Canada
<i>Minecto Duo</i>	thiamethoxam: cyantraniliprole	20% : 20% WG	Syngenta Canada
<i>Movento 240 SC</i>	spirotetramat	240 g/L SC	Bayer CropScience
<i>MPOWER Krypton</i>	chlorpyrifos	480 g/L	NewAgco Inc.
<i>Nolo Bait</i>	spores of <i>Nosema (Paranosema) locustae</i> Canning	Minimum of 2.2x10 ⁶ on coated wheat bran	M&R Durango, Inc.
<i>Nufos 4E</i>	chlorpyrifos	480 g/L EC	FMC Corporation
<i>Oberon</i>	spiromesifen	240 g/L SC	Bayer CropScience
<i>Orthene</i>	acephate	75% SP	Loveland Products Canada
<i>Perm-UP</i>	permethrin	384 g/L EC	United Phosphorous Inc
<i>Phostoxin</i>	aluminum phosphide	55% tablets	Degesch America Inc
<i>Poleci</i>	Delthamethrin	25 g/L EC	Bayer CropScience
<i>Pounce</i>	permethrin	384 g/L EC	FMC Corporation
<i>Protect-It</i>	diatomaceous earth	90%	Hedley Technologies Ltd
<i>Pyrifos 15G</i>	chlorpyrifos	15% G	Gowan Canada
<i>Pyrinex 480EC</i>	chlorpyrifos	480 g/L EC	ADAMA Canada
<i>Rimon 10 EC</i>	novaluron	10% EC	Platform Specialty Products
<i>Sevin XLR</i>	carbaryl	466 g/L	Univar Canada Ltd.
<i>Sharda chlorpyrifos 480 EC</i>	chlorpyrifos	480 g/L EC	Sharda CropChem Canada
<i>Ship 250 EC</i>	cypermethrin	250 g/L	Sharda CropChem Canada
<i>Silencer 120 EC</i>	lambda-cyhalothrin	120 g/L EC	ADAMA Canada
<i>Sluggo Professional</i>	ferric phosphate	0.76 % granules	Engage Agro Corp.
<i>Success 480 SC</i>	spinosad	480 g/L SC	Dow AgroSciences
<i>Superior 70 oil</i>	mineral oil	99%	Loveland Products Canada, N.M. Bartlett Inc.
<i>Tempo</i>	cyfluthrin	20% WP	Bayer CropScience
<i>Thimet 20G</i>	phorate	20% G	Amvac Chemical
<i>UP-Cyde</i>	cypermethrin	250 g/L EC	United Phosphorous Inc
<i>Voliam Xpress</i>	lambda-cyhalothrin:	50 g/L: 100 g/L	Syngenta Canada
<i>Warhawk</i>	chlorantraniliprole chlorpyrifos	480 g/L EC	Loveland Products Canada

Key to Product Pages

Pesticide Product Name

This field lists the pesticide product name. Where there is only one product the commercial “trade” name is given. Where more than one company sells pesticides with the same combination of active ingredients the “generic” (active ingredient) name is given.

If the active ingredients are all in a common formulation (liquid, granule, etc.) the generic name will appear as ‘Ingredient A/ Ingredient B’ and if the active ingredients are in separate containers to be mixed in the sprayer the names are given as ‘Ingredient A + Ingredient B’.

Pesticide Resistance Group

This area will the pesticide active ingredient(s) to the mode of action that ingredient uses and refer to a page number where more information can be found.

Company:

This section identifies the company (or companies) that manufacture or market this crop protection product (or generic equivalents) in Canada as well as the PCP# for that (those) product(s). See page 7 for more information on PCP numbers. PCP#s are given as ‘(PCP#XXXXX)’ where XXXXX is a four or five digit number unique to that product. In some cases, where there are multiple components with separate PCP numbers, the PCP number will be provided below under ‘Formulation:’

Formulation:

This section gives information on the active ingredient and its concentration in the product as well as information on formulation type and packaging types and configurations. Formulation strength (or concentrations) are given in % by weight for dry formulations and g/L for liquid formulations. PCP numbers may also be give for some products (see above).

Crops and Staging:*

This section indicates on which crops the product may be used and what stage of crop development it should be applied at. Rates may also be included in this section if they vary between crop types or crop stage.

***This section will also indicate which crops are registered under the User Requested Minor Use Label Expansion (URMULE) program. Some companies, as a condition of placing these minor crops on their labels request, that users of their product on these crops do so at their own risk because the registration was approved with information the company did not produce.*

These crops will be flagged separately from the main crops.

Pest (Diseases, Insects, Weeds) and Staging:

This section indicates the pests (Diseases, Insects, or Weeds) that are indicated on the product label as controlled or suppressed, as well as any specifics on the timing of application relative to the pest stage if required. Rates may also be included in this section if they differ for different pests or stage of pest.

Rates:

The rates provided in this section are given in the amount of product required per acre and the number of acres treated per package unit where possible. This section will also indicate any adjuvants that are to be used in conjunction with the product and the rate of that adjuvant.

This section will not be present if rates have been integrated into either of the previous **Crops** or **Pest** sections.

Application Information:

- **Water Volume:** This section indicates the minimum carrier water volume to be used to apply the product. Using less than the recommended minimum carrier application volume can negatively affect pesticide performance, particularly with contact pesticides and when using low drift nozzles.
- **Nozzles and Pressure:** This section indicates if there are any particular nozzles that should or should not be used to apply the product. Pressures indicated reflect those for conventional nozzles. Low drift nozzles may require higher pressures for proper performance. A general statement of “Use nozzles and pressures designed to deliver proper coverage with **ASABE** ___ droplets” indicating the ideal droplet sizes to allow for the combination of lowest drift potential and best performance from the pesticide. ASABE refers to the American Society of Agricultural Engineers who have set standards a series of droplet measurements (in microns or micrometres) that classify droplet sizes from ‘fine’ to ‘very coarse’.

How it Works:

This section typically refers to the page where a general description of the various modes of action of either herbicide, fungicides or insecticides.

Effects of Growing Conditions:

This section summarizes any adverse conditions that will affect the biological function of the crop or the target pest and therefore possibly impact the product’s performance. In most cases both crop and target pest must be growing or functioning normally for pesticides to provide expected performance and/or crop tolerance. Adverse weather conditions such as extreme heat, cold, drought or flooding can slow or stop the biological processes in the crop or pest. These biological processes in the crop allow the pesticide to be degraded quickly. If biological processes that are attacked by the pesticide, and under normal conditions would kill the pest, are not functioning normally the pest may be able to rid itself of the pesticide before dying and recover from the application.

Tank Mixes:

This section indicates which other pesticides the pesticide label indicates are registered for use as tank mix combinations with this pesticide.

Common mixes may include:

Herbicides:

- (Subtitles may indicate specific crops or condition restrictions:)

Insecticides:

Fungicides:

Fertilizers:

There may be additional pesticides that are registered but not listed on this product’s label. Other pesticides may have this product listed as a mix option on their labels. The note below (**in bold**) directs users to a chart inside the back cover that show all available mixes for this pesticide. The product listed on the left column of the chart is the product that supports the mix. Mixes supported by both products are marked with an ‘X’. Mixes supported by only one of the products is indicated by an arrow pointing to the left column.

Included in the tank mix section in non-bolded italics may be any precautions against the mixing of pesticides which will have adverse reactions such as crop injury, reduced pest control or unusual increased danger in the use of the product.

Note: The above mixes are those listed on the pesticide label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Restrictions:

Since most pesticides have a capacity to injure neighboring plants, wildlife or people, they will come with restrictions on their use in order to prevent this unintentional damage. Misuse of pesticides may result in as little as temporary or superficial damage to plants or a slight irritation to the eyes or nose, or could also result in poor performance of the pesticide, severe injury and/or yield loss to very sensitive plants and/or unacceptable residues in agricultural commodities, and/or serious illness or death of non-target organism or people. It is important to comply with product restrictions in order to minimize the impact of the pesticide used on non-target organisms and people. A selection of common restrictions and precautions found on product labels are provided in this section, **but it is important to read the label carefully in order to understand how to use the product properly.**

- **Rainfall:** This section indicates the required delay between application and rainfall to avoid reductions in the performance of the product or the unintentional movement of the product.
- **Re-entry:** This section indicates when it is safe for a person to re-enter treated field following an application of a particular pesticide without the same personal protection used to apply the product.
- **Resistance Management:** This section highlights products where an increased risk of the target pests developing resistance to the group of products (typically fungicides) has been identified. If no specific risk has been identified the reader is referred to a general resistance section. All pesticides have some risk of the target pest developing resistance. Rotating pesticide groups using as many different resistance groups as possible in the rotation is one way to avoid or delay resistance development.
- **Grazing:** This section indicates whether and how soon treated crops may be grazed by livestock or otherwise fed to livestock. This restriction is in place to avoid residues of the pesticide from being detected in milk or meat from animals consuming forage, greenfeed or straw from treated crops or forage.
- **Pre-harvest interval:** Is the time that must be left between application of a pesticide and the harvest of a crop in order to prevent greater than allowable residues of the pesticide in the harvested material. Harvest is the cutting of the crop (i.e. combining or hay cut) or removal of the harvestable material from the plant (i.e. picking fruit or stripper header). Maximum Residue Limits (MRLs) are set for commodities based on registered rates and staging of pesticides used in the production of those commodities. Disregarding these intervals can result in residues over the MRLs, which can lead to market disruptions.
- **Re-cropping:** This section indicates how soon specific crops may be seeded into treated fields. Failure to adhere to these delays could result in injury to the following crop.
- **Aerial Application:** This section indicates whether the product may be applied by aircraft and any special conditions that may be necessary.
- **Labelling:** In addition to other precautions and warnings, seed treatment products will also have statements about how seed treated with the product should be labeled.
- **Storage:** This section indicates how the product must be stored. As a general rule, unused pesticides should always be stored in their original containers in a secure, dry area, away from other pesticides, food or feed.
- **Buffer Zones:** This section will indicate any setback distances that are required from sensitive aquatic or upland habitats. Newer labels may indicate that these distances are from the downwind edge of the boom but older labels may not. Examples of aquatic habitats are lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands. Examples of terrestrial habitats are grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands.

In addition to the set back or 'buffer' distances indicated on product labels, provincial environment departments may also have additional restrictions or requirements for permits to apply pesticides to or near water. Check with the provincial environment department/ministry for more information.

Tank Cleaning:

This section describes the measures that are required to properly clean out spray tanks. A general overview of sprayer cleaning is given on page 12, but products where there is a high risk of crop damage as a result of very low level contamination of the spray solution, will have specific measures indicated.

Hazard Rating:

This section indicates the relative toxicity of the pesticide, formulations or components. For an explanation of the symbols used here see pages 7-8. An additional symbol that is used that is not a standard symbol is the (!) exclamation mark which indicates an otherwise undefined risk factor (i.e. irritation).

Example:



Caution – Eye Irritant

Some older products have not had hazard ratings developed, while other products have very low toxicity and do not have hazard warnings. Even in the absence of a hazard rating users should wear a minimum of nitrile gloves and an apron as well as long sleeved apparel during mixing and avoid unnecessary exposure.