

For the Period July 19 to 25, 2016

Good progress was made with haying this past week. Livestock producers have 50 per cent of the hay crop baled or put into silage and another 20 per cent cut and ready for baling, according to Saskatchewan Agriculture's weekly Crop Report. The five-year average (2011-2015) is 55 per cent baled or put into silage.

Hay quality is rated as five per cent excellent, 70 per cent good, 24 per cent fair and one per cent poor. Hay is slow to dry in the swath due to rain and high humidity.

Crops are generally in good condition. Sixty-two per cent of the spring wheat, winter wheat, flax and canola are in good condition. Wet conditions across much of the lentil-producing area has affected quality. Lentils are rated as 15 per cent excellent, 47 per cent good, 26 per cent fair, 10 per cent poor and two per cent very poor. Peas are rated as 30 per cent excellent, 52 per cent good, 16 per cent fair and two per cent poor.

Most areas of the province received rain during the past week, with the heaviest showers dropping between 50 and 100 mm of precipitation. There were reports of 200 mm of rain in the Elfros region. Heavy rain over the past several weeks has caused crops to lodge, and they remain under flooding stress in some areas. Lentils and peas in many areas of the province are suffering from too much moisture. Diseases and hail have also caused crop damage.

Provincially, cropland topsoil moisture is rated as 15 per cent surplus, 80 per cent adequate and five per cent short. Hay land and pasture topsoil moisture is rated as nine per cent surplus, 83 per cent adequate and eight per cent short.

Harvesting of winter wheat, peas and lentils in some areas is expected to start in the coming week. Farmers are busy haying and getting ready for harvest.

One year ago

Sixty-seven per cent of the hay crop had been baled or put into silage. Harvest was just beginning with desiccation of pulses and some winter cereals combined. Moisture conditions improved with recent rains.

Follow the 2016 Crop Report on
Twitter at @SKAgriculture

For further information, contact Daphne Cruise, PAg,
Crop Management Specialist, Moose Jaw, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3587, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.saskatchewan.ca/crop-report.

Southeastern Saskatchewan:

- Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE – Radville and Lake Alma areas

Fifty per cent of the hay crop is baled and 22 per cent is cut and ready for baling. Haying is furthest advanced in Crop District 2A, which has 65 per cent baled, and least advanced in Crop District 1B, which has 36 per cent baled. Hay quality is rated as 79 per cent good and 21 per cent fair. High humidity and rain showers are causing quality loss in the swath. Dry weather is needed to get the rest of the hay up in good shape. There are a few reports of producers starting to take their second cut of hay.

The southeast recorded varying amounts of rain during the week, although amounts were less than they have been in the previous weeks. Most areas, with the exception of the few that were hit with thunderstorms, reported less than 15 mm of rain. The Carnduff area reported 15 mm, the Grenfell area 28 mm, Weyburn 34mm, Moose Jaw 73 mm and Radville 10 mm. The Maryfield area holds the regional record for greatest amount of precipitation since April 1 (413 mm). The Odessa area has recorded the least amount of rain for the region since April 1 (210 mm).

Cropland topsoil moisture is rated as 16 per cent surplus, 81 per cent adequate and three per cent short. Hay land and pasture topsoil moisture is rated as 10 per cent surplus, 85 per cent adequate and five per cent short. Crop districts 1B and 2B are reporting that 32 and 23 per cent, respectively, of cropland acres have surplus topsoil moisture at this time. Twenty-nine per cent of hay land and pasture in CD 1B has surplus topsoil moisture. There is crop loss where land has flooded or remained wet.

Most crops in the region remain in good condition, although the pulse crops are succumbing to excess moisture stress. Crop damage this week was mainly caused by localized flooding and diseases such as root rot, leaf spots, sclerotinia and botrytis. Hail was reported in the region, which caused significant damage in some areas. Marble- to golf-ball-sized hail was reported in the Indian Head and Moose Jaw regions. Some producers are expecting to start harvest in the next week if the weather cooperates.

Southwestern Saskatchewan:

- Crop District 3ASW – Coronach, Assiniboia and Ogema areas
- Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B – Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 – Consul, Maple Creek and Leader areas

Scattered and sporadic rain showers in the region allowed some producers to make good progress in haying while others were slowed. Fifty per cent of the hay crop has been baled and an additional 19 is cut and ready for baling. Hay quality is rated as three per cent excellent, 66 per cent good and 31 per cent fair. Haying is most advanced in Crop District 4A, which has 65 per cent baled, and least advanced in Crop District 3AW, which has 30 per cent baled. The hay has been slow to dry due to frequent rain showers and high humidity. Rainfall varied greatly in the region, from nothing to 93 mm in the Kyle area. Other areas receiving large amounts of rain include Moose Jaw (64 mm), Big Beaver (30

mm), Shaunavon (28 mm), Swift Current (31 mm) and Leader (52 mm). The Webb area holds the regional record for greatest amount of precipitation since April 1 (433 mm).

Cropland topsoil moisture conditions are rated as 14 per cent surplus, 84 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as six per cent surplus, 89 per cent adequate and five per cent short. Crop District 4B is reporting that 25 per cent of the cropland has surplus topsoil moisture. Crop loss is occurring due to excess moisture in some areas, particularly in low spots and areas in the field that may have been saturated prior to the rain.

The majority of crops in the region are in good condition, although a pulse quality downgrade is imminent due to excess moisture in areas. The majority of crop damage this past week is attributed to localized flooding, strong winds, lodging and hail. There are reports of some roads being washed out in the Kyle area. There are a couple reports of fall rye being combined. Although harvest will be later than anticipated because of the moisture, some pea crops are expected to be harvested in the coming week.

East-Central Saskatchewan:

- Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A – Lumsden, Craik, Watrous and Clavet areas

Good haying progress was made in some areas of the region. Forty-five per cent of the hay crop has been baled and an additional 23 per cent is cut and ready to bale. Crop District 6A has 53 per cent baled, while Crop District 5B has 38 per cent baled. Hay quality is rated as seven per cent excellent, 74 per cent good, 15 per cent fair and four per cent poor. A few days of warm and breezy conditions allowed haying to continue toward the end of the week. Haying has been very slow in the region due to less-than-ideal weather conditions. Frequent rain showers and high humidity have caused quality to deteriorate.

Heavy rain fell on the region during the past week. Amounts ranged from 4 mm (Bulyea area) to 93 mm (Wynyard area). There were reports of over 200 mm falling on the Elfron region within two hours. Roads and fields were flooded and culverts were washed out. Most other areas recorded over 25 mm of rain for the week. The Kelvington area holds the regional record for greatest amount of precipitation since April 1 (365 mm).

Topsoil moisture conditions on cropland are rated as 17 per cent surplus and 83 per cent adequate. Hay land and pasture topsoil moisture is rated as 13 per cent surplus, 86 per cent adequate and one per cent short. Crop District 5B is reporting that 30 per cent of the cropland has surplus topsoil moisture at this time. There is crop loss due to excess moisture in some areas, particularly in low spots and areas in the field that have been saturated for a couple of weeks.

Most crops remain in good condition. Sources of crop damage this week include localized flooding, hail, wind and diseases such as root rot and other pulse diseases. Some pulse crops are expected to be desiccated in the coming week if weather cooperates.

West-Central Saskatchewan:

- Crop Districts 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A – Rosetown, Kindersley, Eston, Major
- Crop District 7B - Kerrobert, Macklin, Wilkie and Biggar areas

Haying continues despite the rain. Forty-seven per cent of the hay crop has been baled or put into silage, and 24 per cent is cut and ready for baling. Continuous rain and high humidity are taking their toll on hay quality, which is now rated as five per cent excellent, 48 per cent good and 47 per cent fair.

Rainfall varied greatly throughout the region, ranging from nothing in the Tramping Lake and Unity areas to 50 mm in the Conquest area. Other areas that received large amounts of rain this past week include Dinsmore (37 mm), Sonningdale (24 mm), Kindersley (48 mm), Biggar (41 mm) and Macklin (37 mm). The Conquest area holds the regional record for greatest amount of precipitation since April 1 (332 mm).

Topsoil moisture conditions on cropland are rated as 15 per cent surplus and 83 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as 10 per cent surplus, 86 per cent adequate and four per cent short. Crop District 7A is reporting that 25 per cent of cropland has surplus topsoil moisture. There is crop loss due to excess moisture in some areas.

The majority of crops are in good condition, but pulses are taking the biggest hit when it comes yield loss due to excess moisture. Crop damage this week is attributed to localized flooding, diseases and hail. Baseball-sized hail and heavy rain was reported in the Outlook and Conquest areas, causing significant damage to crops and property. There are predictions of peas being desiccated this coming week if weather cooperates. Warm and dry weather is needed as harvest approaches if fields in some areas are to be accessed. Despite the excess moisture in most areas of the region, there are few areas that could use more moisture to help fill and finish off the crops.

Northeastern Saskatchewan:

- Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas

The region is furthest ahead in haying, with 59 per cent of the hay crop baled or put into silage. An additional 17 per cent is cut and ready for baling. Ninety-two per cent of the hay is rated as good quality and eight per cent is rated as poor. Although rain was reported, it resulted in very little down time for cutters and balers in most areas. High humidity is causing delays in baling.

Rainfall ranged from nothing in the Kinistino area to 36 mm in the Porcupine Plain and Tisdale areas. Other areas receiving significant rainfall include Star City (20 mm), Birch Hills (15 mm), Alvena (6 mm), and Garrick (21 mm). The Porcupine Plain area holds the regional record for greatest amount of precipitation since April 1 (348 mm).

Cropland topsoil moisture conditions are rated as 27 per cent surplus, 71 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as 36 per

cent surplus and 64 per cent adequate. Crop District 8A is reporting that 46 per cent of cropland and 43 per cent of hay land and pasture have surplus topsoil moisture at this time.

Crops remain are generally in good condition. Crop damage includes hail, lodging and disease. Significant rain in the past few weeks has washed out roads in some areas and is flooding crops, causing loss. Some harvesting will begin in the next week or so. Warm, dry weather is needed to dry the soil enough to support harvest equipment.

Northwestern Saskatchewan:

- Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

Drier weather has allowed for good progress to be made in the hay field. Fifty-three per cent of the hay crop has been baled and another 21 per cent is cut and ready for baling. Hay quality at this time is rated as 14 per cent excellent, 72 per cent good and 14 per cent fair. The region did not receive as much rain as other areas in the province, so haying was able to continue for the most part. Hay that has been cut and baled the last two weeks is of much better quality as little, if any, rain fell on the swaths.

Rainfall ranged from nil to 16 mm (Turtleford area). The Hafford and Lloydminster areas received 12 mm, Debden 7 mm and Barthel and Meadow Lake 10 mm. The Frenchman Butte area holds the regional record for greatest amount of precipitation since April 1 (278 mm). The Speers area has reported the least amount of rainfall since April 1 (100 mm).

Cropland topsoil moisture is rated as one per cent surplus, 71 per cent adequate and 28 per cent short. Hay land and pasture topsoil moisture is rated as 66 per cent adequate, 33 per cent short and one per cent very short. Although most of the region has adequate topsoil moisture, there are areas that are short on moisture and could use a good rain.

Crops in general are in good condition. The region has not received as much precipitation throughout the growing season as the rest of the province. Crop damage this past week is attributed to strong winds, lodging and diseases, particularly in peas.

| Provincial | | | | | | | |
|-------------|--------------|----------|--------------|---------|-----|--------|------------|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 36 | 48 | 31 | 37 | 23 | 27 | 19 |
| % good | 61 | 50 | 62 | 58 | 69 | 65 | 73 |
| % fair | 3 | 2 | 7 | 4 | 7 | 7 | 6 |
| % poor | 0 | 0 | 0 | 1 | 1 | 1 | 1 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| | | | | | | | |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 31 | 31 | 27 | 9 | 30 | 15 | 8 |
| % good | 62 | 61 | 66 | 66 | 52 | 47 | 68 |
| % fair | 6 | 6 | 6 | 20 | 16 | 26 | 8 |
| % poor | 1 | 1 | 1 | 5 | 2 | 10 | 16 |
| % very poor | 0 | 1 | 0 | 0 | 0 | 2 | 0 |

| Southeast | | | | | | | |
|-------------|--------------|----------|--------------|---------|-----|--------|------------|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 48 | 46 | 39 | 35 | 42 | 32 | 23 |
| % good | 51 | 50 | 56 | 62 | 50 | 54 | 71 |
| % fair | 1 | 4 | 5 | 3 | 7 | 13 | 5 |
| % poor | 0 | 0 | 0 | 0 | 1 | 1 | 1 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | | | | | | | |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 35 | 36 | 10 | 8 | 35 | 17 | N/A |
| % good | 60 | 59 | 90 | 64 | 55 | 49 | N/A |
| % fair | 4 | 5 | 0 | 23 | 7 | 21 | N/A |
| % poor | 1 | 0 | 0 | 5 | 2 | 8 | N/A |
| % very poor | 0 | 0 | 0 | 0 | 1 | 5 | N/A |

| Southwest | | | | | | | |
|-------------|--------------|----------|--------------|---------|-----|--------|------------|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 38 | 57 | 37 | 39 | 41 | 27 | 25 |
| % good | 57 | 43 | 58 | 56 | 57 | 70 | 61 |
| % fair | 5 | 0 | 5 | 5 | 2 | 3 | 10 |
| % poor | 0 | 0 | 0 | 0 | 0 | 0 | 3 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| | | | | | | | |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 26 | 44 | 28 | 37 | 25 | 10 | 25 |
| % good | 68 | 51 | 62 | 63 | 48 | 47 | 61 |
| % fair | 5 | 4 | 8 | 0 | 26 | 29 | 10 |
| % poor | 1 | 1 | 2 | 0 | 1 | 13 | 3 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 1 | 1 |

| East-central | | | | | | | |
|--------------|-------------|----------|--------------|-------|-----|--------|------------|
| | Winter whea | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 29 | 27 | 33 | 33 | 28 | 31 | 9 |
| % good | 69 | 70 | 59 | 56 | 66 | 62 | 71 |
| % fair | 2 | 3 | 7 | 11 | 6 | 6 | 20 |
| % poor | 0 | 0 | 1 | 0 | 0 | 1 | 0 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |


| East-central (continued) | | | | | | | |
|--------------------------|------|--------|---------|---------|-----|--------|----------|
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 36 | 33 | N/A | 10 | 35 | 34 | N/A |
| % good | 58 | 58 | N/A | 78 | 55 | 38 | N/A |
| % fair | 5 | 7 | N/A | 10 | 8 | 25 | N/A |
| % poor | 1 | 2 | N/A | 1 | 1 | 3 | N/A |
| % very poor | 0 | 0 | N/A | 1 | 1 | 0 | N/A |

| West-central | | | | | | | |
|--------------|-------------|----------|--------------|---------|-----|--------|------------|
| | Winter whea | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 20 | N/A | 36 | 36 | 35 | 33 | 25 |
| % good | 73 | N/A | 57 | 61 | 56 | 61 | 73 |
| % fair | 7 | N/A | 7 | 3 | 9 | 6 | 2 |
| % poor | 0 | N/A | 0 | 0 | 0 | 0 | 0 |
| % very poor | 0 | N/A | 0 | 0 | 0 | 0 | 0 |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 23 | 38 | 37 | 20 | 34 | 15 | N/A |
| % good | 65 | 58 | 59 | 54 | 50 | 45 | N/A |
| % fair | 11 | 4 | 4 | 26 | 9 | 24 | N/A |
| % poor | 1 | 0 | 0 | 0 | 6 | 11 | N/A |
| % very poor | 0 | 0 | 0 | 0 | 1 | 5 | N/A |

| Northeast | | | | | | | |
|-------------|-------------|----------|--------------|---------|-----|--------|------------|
| | Winter whea | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | N/A | N/A | 18 | 45 | 19 | 33 | 16 |
| % good | N/A | N/A | 73 | 53 | 74 | 61 | 66 |
| % fair | N/A | N/A | 9 | 2 | 7 | 6 | 12 |
| % poor | N/A | N/A | 0 | 0 | 0 | 0 | 6 |
| % very poor | N/A | N/A | 0 | 0 | 0 | 0 | 0 |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 16 | 16 | N/A | N/A | 16 | N/A | N/A |
| % good | 66 | 71 | N/A | N/A | 66 | N/A | N/A |
| % fair | 17 | 7 | N/A | N/A | 14 | N/A | N/A |
| % poor | 1 | 4 | N/A | N/A | 2 | N/A | N/A |
| % very poor | 0 | 2 | N/A | N/A | 2 | N/A | N/A |

| Northwest | | | | | | | |
|-------------|--------------|----------|--------------|---------|-----|--------|------------|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | N/A | N/A | 22 | N/A | 7 | 11 | N/A |
| % good | N/A | N/A | 70 | N/A | 84 | 76 | N/A |
| % fair | N/A | N/A | 8 | N/A | 9 | 13 | N/A |
| % poor | N/A | N/A | 0 | N/A | 0 | 0 | N/A |
| % very poor | N/A | N/A | 0 | N/A | 0 | 0 | N/A |
| | | | | | | | |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | N/A | 25 | N/A | N/A | 32 | N/A | N/A |
| % good | N/A | 67 | N/A | N/A | 59 | N/A | N/A |
| % fair | N/A | 8 | N/A | N/A | 8 | N/A | N/A |
| % poor | N/A | 0 | N/A | N/A | 1 | N/A | N/A |
| % very poor | N/A | 0 | N/A | N/A | 0 | N/A | N/A |


Crop Districts and Rural Municipalities in Saskatchewan



Geomatic Services, Ministry of Agriculture June 10, 2014

Weekly Rainfall


from July 19 to July 25, 2016



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall

from July 19 to July 25, 2016



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period 19-25, 2016


| Crop Dist. | R.M. No. | Name | Past Week | Since 1-Apr | Crop Dist. | R.M. No. | Name | Past Week | Since 1-Apr | Crop Dist. | R.M. No. | Name | Past Week | Since 1-Apr |
|------------|----------|----------------|-----------|-------------|------------|----------|------------------|-----------|-------------|------------|----------|-----------------|-----------|-------------|
| 1A | 2 | Mount Pleasant | 15 | 294 | 4A | 49 | White Valley | 7 | 272 | 7A | 287 | St. Andrews | 31 | 308 |
| | 3 | Enniskillen | 3 | 273 | | 51 | Reno | NIL | 164 | | 288 | Pleasant Valley | N/A | 166 |
| | 33 | Moose Creek | N/A | 239 | | 79 | Arlington | 11 | 326 | | 290 A | Kindersley | 25 | 198 |
| | 34 | Browning | N/A | 335 | | 109 A | Carmichael | N/A | 281 | | 290 B | Kindersley | 48 | 254 |
| | 61 | Antler | N/A | 282 | | 109 B | Carmichael | N/A | 216 | | 290 C | Kindersley | 35 | 239 |
| 1B | 64 | Brock | 2 | 234 | 4B | 110 | Piapot | 3 | 282 | 7B | 292 | Milton | 31 | 314 |
| | 65 | Tecumseh | 11 | 236 | | 111 | Maple Creek | N/A | 208 | | 317 A | Marriott | N/A | 220 |
| | 91 | Maryfield | 6 | 413 | | 139 | Gull Lake | NIL | 320 | | 317 B | Marriott | 16 | 191 |
| | 122 | Martin | 15 | 325 | | 142 | Enterprise | N/A | 108 | | 318 | Mountain View | 41 | 251 |
| | 123 | Silverwood | 7 | 278 | | 169 | Pittville | N/A | 282 | | 320 A | Oakdale | 10 | 219 |
| 2A | 124 A | Kingsley | 9 | 286 | 5A | 231 | Happyland | 52 | 336 | | 320 B | Oakdale | 9 | 264 |
| | 124 B | Kingsley | N/A | NIL | | 183 | Fertile Belt | N/A | 311 | 7B | 321 | Prairiedale | 20 | 280 |
| | 125 A | Chester | 10 | 375 | | 211 | Churchbridge | N/A | 223 | | 347 | Biggar | 25 | 180 |
| | 125 B | Chester | 21 | 294 | | 213 | Saltcoats | 33 | 162 | | 350 A | Mariposa | NIL | 147 |
| | 151 | Rocanville | 10 | 312 | | 241 | Calder | 36 | 189 | | 350 B | Mariposa | N/A | 146 |
| 2B | 154 | Elcapo | N/A | 280 | 5B | 243 | Wallace | N/A | 215 | | 351 | Progress | N/A | 195 |
| | 155 | Wolseley | 28 | 322 | | 244 | Orkney | N/A | 186 | 8A | 352 | Heart's Hill | 18 | 326 |
| | 67 | Weyburn | 7 | 252 | | 245 A | Garry | 46 | 248 | | 377 | Glenside | 7 | 162 |
| | 68 | Brokenshell | 9 | 313 | | 245 B | Garry | 25 | 196 | | 378 | Rosemount | 10 | 187 |
| | 97 | Wellington | N/A | 253 | | 245 C | Garry | N/A | 148 | | 379 | Reford | 12 | 175 |
| 3ASE | 127 A | Francis | 10 | 210 | 6A | 246 | Ituna Bon Accord | 12 | 155 | | 381 | Grass Lake | 4 | 144 |
| | 127 B | Francis | 21 | 212 | | 247 | Kellross | 7 | 161 | 8B | 382 | Eye Hill | 37.5 | 249 |
| | 129 | Bratt's Lake | 10 | 247 | | 248 | Touchwood | 71 | 229 | | 409 | Buffalo | N/A | 125 |
| | 131 A | Baildon | N/A | 304 | | 271 | Cote | N/A | 170 | | 410 | Round Valley | NIL | 136 |
| | 131 B | Baildon | 28 | 375 | | 273 | Sliding Hills | 8 | 128 | | 395 | Porcupine | 36 | 348 |
| 3ASW | 156 A | Indian Head | 36.7 | 271 | 6B | 277 | Emerald | 32 | 232 | | 397 | Barrier Valley | 19.8 | 284 |
| | 156 B | Indian Head | 21 | 293 | | 305 | Invermay | 59 | 319 | 9AE | 428 | Star City | 20 | 275 |
| | 159 | Sherwood | 34 | 230 | | 307 | Elfros | 58 | 324 | | 456 | Arborfield | 18 | 346 |
| | 160 | Pense | 12 | 240 | | 308 A | Big Quill | 93 | 253 | | 457 | Connaught | 36 | 265 |
| | 161 | Moose Jaw | N/A | 271 | | 308 B | Big Quill | N/A | 184 | | 486 | Moose Range | 16 | 291 |
| 3AN | 162 | Caron | 20 | 284 | 6A | 331 | Livingston | 36 | 292 | 9AW | 487 | Nipawin | 21 | 310 |
| | 191 | Marquis | 19 | 276 | | 336 | Sasman | 27 | 227 | | 369 | St. Peter | 9 | 166 |
| | 38 A | Laurier | 10.4 | 356 | | 337 | Lakeview | N/A | 264 | | 370 A | Humboldt | N/A | 205 |
| | 38 B | Laurier | 5 | 284 | | 338 | Lakeside | 34 | 270 | | 370 B | Humboldt | N/A | 186 |
| | 39 | The Gap | N/A | 291 | | 366 | Kelvington | 39 | 365 | 9B | 371 | Bayne | 4 | 252 |
| 3BS | 10 | Happy Valley | 30.5 | 315 | 6A | 367 | Ponass Lake | 10 | 273 | | 372 | Grant | 3.4 | 131 |
| | 12 | Poplar Valley | 2 | 249 | | 190 A | Dufferin | 26 | 321 | | 400 | Three Lakes | 5 | 222 |
| | 40 | Bengough | N/A | N/A | | 190 B | Dufferin | 15 | 289 | | 402 | Fish Creek | 6 | 174 |
| | 42 | Willow Bunch | NIL | 297 | | 190 C | Dufferin | 37 | 260 | | 429 | Flett's Springs | 5 | 155 |
| | 43 | Old Post | NIL | 259 | | 190 D | Dufferin | N/A | 192 | | 459 | Kinistino | NIL | 164 |
| 3BN | 73 A | Stonehenge | NIL | 332 | 6B | 219 A | Longlaketon | N/A | 196 | | 460 | Birch Hills | 14.8 | 166 |
| | 73 B | Stonehenge | N/A | 348 | | 219 B | Longlaketon | 28 | 241 | 9AE | 488 | Torch River | 21 | 272 |
| | 102 | Lake Johnston | NIL | 211 | | 220 | McKillop | 4 | 219 | | 520 | Paddockwood | N/A | 143 |
| | 103 | Sutton | 5 | 267 | | 221 A | Sarnia | 10.7 | 274 | | 521 | Lakeland | N/A | 143 |
| | 132 A | Hillsborough | 9 | 287 | | 221 B | Sarnia | N/A | 222 | 9AW | 406 | Mayfield | 8 | 170 |
| 3BS | 132 B | Hillsborough | 8 | 319 | 6B | 222 | Craik | 19 | 288 | | 435 | Redberry | 12 | 173 |
| | 134 | Shamrock | N/A | 17 | | 251 | Big Arm | 20 | 217 | | 436 | Douglas | NIL | 100 |
| | 193 A | Eyebrow | 7 | 260 | | 252 | Arm River | N/A | 209 | | 463 | Duck Lake | N/A | 136 |
| | 193 B | Eyebrow | N/A | 340 | | 279 | Mount Hope | N/A | 233 | | 467 A | Round Hill | NIL | 220 |
| | 17 | Val Marie | 9.3 | 304 | | 282 | McCraney | 59 | 275 | 9B | 467 B | Round Hill | 2 | 186 |
| 3BN | 18 | Lone Tree | 17 | 358 | 6B | 312 | Morris | 48 | 344 | | 494 | Canwood | 7 | 124 |
| | 75 | Pinto Creek | 4 | 354 | | 313 | Lost River | 16 | 185 | | 438 | Battle River | 4 | 184 |
| | 76 | Auvergne | 3 | 273 | | 339 | Leroy | 31 | 207 | | 440 | Hillsdale | NIL | 99 |
| | 77 | Wise Creek | 4 | 350 | | 340 | Wolverine | 23 | 244 | | 442 | Manitou Lake | NIL | 187 |
| | 78 | Grassy Creek | 3.5 | 265 | | 341 | Viscount | 20 | 196 | | 498 A | Parkdale | 3 | 194 |
| 3BN | 105 | Glenbain | 7 | 347 | 6B | 343 A | Blucher | N/A | 149 | 9B | 498 B | Parkdale | 1 | 41 |
| | 106 | Whiska Creek | 11 | 319 | | 343 B | Blucher | 12 | 157 | | 499 A | Mervin | 16.1 | 207 |
| | 107 | Lac Pelletier | N/A | 340 | | 223 A | Huron | N/A | 236 | | 499 B | Mervin | N/A | N/A |
| | 108 | Bone Creek | 28 | 353 | | 223 B | Huron | 5 | 262 | | 501 A | Frenchman Butte | 8 | 184 |
| | 138 A | Webb | NIL | 433 | | 284 A | Rudy | 46 | 286 | | 501 B | Frenchman Butte | 5 | 177 |
| 3BN | 138 B | Webb | 4 | 319 | 6B | 285 B | Rudy | 32 | 284 | 9B | 501 C | Frenchman Butte | N/A | 278 |
| | 166 | Excelsior | 24 | 341 | | 285 A | Fertile Valley | 50 | 332 | | 502 | Britannia | 12 | 237 |
| | 167 | Sask. Landing | 8.2 | 330 | | 285 B | Fertile Valley | 46 | 323 | | 561 | Loon Lake | 10 | 187 |
| | 168 A | Riverside | 31 | 369 | | 286 | Milden | 37 | 298 | | 588 A | Meadow Lake | 12 | 154 |
| | 168 B | Riverside | 17.8 | 234 | | 314 | Dundurn | N/A | 206 | | 588 B | Meadow Lake | 10 | 164 |
| 3BN | 226 | Victory | 14 | 276 | 6B | 344 | Corman Park | N/A | 111 | 9B | 622 | Beaver River | 12.4 | 184 |
| | 228 | Lacadena | 24 | 329 | | 346 | Perdue | 21 | 280 | | | | | |
| | 257 | Monet | 93 | 433 | | 376 | Eagle Creek | 24 | 196 | | | | | |
| | | | | | | 403 | Rosthern | 2 | 140 | | | | | |
| | | | | | | | | | | | | | | |

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Municipality No: A, B, C and D - more than one reporter

Cumulative Rainfall


from April 1 to July 25, 2016



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Cropland Topsoil Moisture Conditions


July 25, 2016



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Hay and Pasture Topsoil Moisture Conditions

July 25, 2016



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.