

For the Period May 23 to 29, 2017

Saskatchewan producers now have 81 per cent of the 2017 crop in the ground, right in line with the five-year (2012-2016) seeding average for this time of year of 82 per cent. Wet field conditions and frequent rainfall in many northern areas have delayed progress, and producers there will need a couple more weeks of warm and dry weather.

Seeding is most advanced in the southeast, where producers have 95 per cent of the crop in the ground. Ninety-four per cent is seeded in the southwest; 80 per cent in the west-central region; 79 per cent in the east-central region; 76 per cent in the northwest and 43 per cent in the northeast.

One year ago

Producers had seeded 94 per cent of the 2016 crop. Cool and wet weather was delaying seeding and herbicide applications for many producers.

Follow the 2017 Crop Report on
Twitter @SKAgriculture

Seeding Progress May 29, 2017 % Seeded	
Spring wheat	81
Durum	92
Barley	73
Oats	55
Canary seed	71
Flax	80
Canola	76
Mustard	78
Soybeans	91
Lentils	96
Field peas	95
Chickpeas	94

Ninety-six per cent of the lentils, 95 per cent of the field peas, 92 per cent of the durum, 91 per cent of the soybeans, 81 per cent of the spring wheat, 80 per cent of the flax, 78 per cent of the mustard, 76 per cent of the canola and 73 per cent of the barley have now been seeded.

A slowly moving weather system brought significant rainfall to many areas in the north last week, but missed most of the central and southern parts of the province. The Nipawin area received 65 mm of rain, while many parts of central and southern Saskatchewan received nothing. Fields in the north remain very wet, while many fields in other parts of the province are in need of rain to help crops germinate and emerge.

Seeding Progress in SK Per cent seeded All Crops	
May 29, 2017	81
May 30, 2016	94
June 1, 2015	97
June 2, 2014	78
May 27, 2013	67
May 28, 2012	74
5 year avg. (2012-2016)	82
10 year avg. (2007-2016)	81

Provincially, cropland topsoil moisture is rated as 11 per cent surplus, 75 per cent adequate, 12 per cent short and two per cent very short. Hay land and pasture topsoil moisture is rated as five per cent surplus, 70 per cent adequate, 23 per cent short and two per cent very short.

For further information, contact Shannon Friesen, PAg,
Cropping Management Specialist, Moose Jaw, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3592, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.saskatchewan.ca/crop-report.

Overall, emerged crops are in fair-to-excellent condition, but emergence has been delayed in many areas by the cool weather and dry field conditions. The majority of crop damage this week was caused by strong winds, frost, hail, localized flooding and lack of moisture. Flea beetles and cutworms have been reported in canola crops, and some fields have had to be re-seeded.

Producers are busy seeding, controlling weeds and picking rocks.

SaskPower reports that there have been 91 incidents in May of farm equipment coming in contact with power poles or lines, including 15 incidents between May 22 and 27 alone. Weekly and monthly totals are adjusted on an ongoing basis, as some incidents are not reported to SaskPower immediately after they occur. Producers are urged to be especially careful when using equipment around power lines. Safety information is available at www.saskpower.com/safety.

Southeastern Saskatchewan:

- Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE – Radville, Minton and Lake Alma areas

Producers in the region are wrapping up seeding operations and now have 95 per cent of the crop seeded. The five-year (2012-2016) seeding average for this time of year is 80 per cent. However, some producers will need another week or more of good weather to complete seeding.

Rainfall last week ranged from nil to 15 mm in the Whitewood area. The Frobisher and Odessa areas reported 9 mm of rain, the Lampman and Tantallon areas 14 mm, the Weyburn area 5 mm, the Briercrest area 1 mm and the Radville area 2 mm. The Moose Jaw area has reported receiving the most precipitation (69 mm) in the region since April 1.

Topsoil moisture conditions continue to deteriorate in the region due to strong winds. Cropland topsoil moisture is rated as one per cent surplus, 63 per cent adequate, 26 per cent short and 10 per cent very short. Hay land and pasture topsoil moisture is rated as one per cent surplus, 55 per cent adequate, 35 per cent short and nine per cent very short. Crop District 2A is reporting that 30 per cent of cropland and 44 per cent of hay land and pasture are very short of topsoil moisture. A significant rain will be needed soon to help crops germinate and emerge and pastures grow. There are concerns that crop and hay yields will be negatively affected if rain is not received soon.

Overall, emerged crops are in fair-to-excellent condition, but emergence has been delayed in many areas by the cool weather and dry field conditions. The majority of crop damage this past week was caused by lack of moisture, frost, hail and strong winds. Flea beetles and cutworms are causing damage in canola fields and some fields are being re-seeded as moisture permits. Frost damage has also resulted in some flax and canola fields being re-seeded. Strong winds last week have not only delayed herbicide applications but have caused soil to erode and drift in many fields. There was also damage reported to buildings and trees.

Farmers are busy seeding, controlling weeds and rolling pulses.

Southwestern Saskatchewan:

- Crop District 3ASW – Coronach, Assiniboia and Ogema areas
- Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B – Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 – Consul, Maple Creek and Leader areas

Many producers in the region have wrapped up their seeding efforts. Ninety-four per cent of the crop is now in the ground, well ahead of the five-year (2012-2016) seeding average for this time of year of 88 per cent. While many producers have completed seeding, others will need another week or more.

Rainfall last week ranged from nil to 9 mm in the Old Wives area. The Rockglen and Limerick areas reported 3 mm of rain, the Eyebrow area 2 mm, the Hazenmore, Gull Lake and Tyner areas 7 mm, the Tompkins area 8 mm and the Shaunavon and Rush Lake areas 4 mm. The Gull Lake area has reported receiving the most precipitation (69 mm) in the region since April 1.

Strong winds continue to dry out topsoil in the region. Cropland topsoil moisture is rated as one per cent surplus, 79 per cent adequate and 20 per cent short. Hay land and pasture topsoil moisture is rated as 60 per cent adequate and 40 per cent short. Crop District 4B is reporting that 33 per cent of cropland and 67 per cent of hay land and pasture are short of topsoil moisture. A significant rain will be needed soon to help crops germinate and emerge and pastures grow. There are concerns that crop and hay yields will be negatively affected if rain is not received soon.

Overall, emerged crops are in fair-to-excellent condition, but emergence has been delayed in many areas by the cool weather and dry field conditions. Most of the crop damage this past week was caused by frost, lack of moisture and strong winds. Flea beetles, cutworms and wireworms have also caused damage in some areas. Strong winds have not only delayed herbicide applications but have caused soil to erode and drift in many fields.

Farmers are busy seeding, controlling weeds and moving cattle.

East-Central Saskatchewan:

- Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A – Lumsden, Craik, Watrous and Clavet areas

Significant seeding progress was made last week thanks to warm and relatively dry weather. The east-central region now has 79 per cent of the crop seeded, well ahead of the five-year (2012-2016) seeding average for this time of year of 74 per cent. Producers expect to wrap up seeding operations in the coming weeks if field conditions remain ideal. Some areas continue to deal with excess moisture, so warm and dry weather will be needed to help dry fields up.

Rainfall last week ranged from trace amounts to 23 mm in the Kamsack area. The Langenburg area reported 14 mm of rain, the Ituna area 15 mm, the Quill Lake area 18 mm, the Kelvington

area 20 mm, the Earl Grey area 10 mm, the Holdfast area 8 mm and the Semans area 12 mm. The Foam Lake area has reported receiving the most precipitation (75 mm) in the region since April 1.

Although recent winds have helped to dry out some wet fields, it has caused topsoil moisture to deteriorate in other areas. Cropland topsoil moisture is rated as nine per cent surplus, 82 per cent adequate and nine per cent short. Hay land and pasture topsoil moisture is rated as five per cent surplus, 81 per cent adequate and 14 per cent short.

Crop District 5B is reporting that 21 per cent of cropland acres have surplus topsoil moisture, while CD 6A is reporting that 20 per cent of cropland acres are short of topsoil moisture.

Crops are just beginning to emerge and are in fair-to-excellent condition. Emergence has been slower than normal due to cooler temperatures. There were reports of light frost last week, although damage has been minimal so far. Flea beetles and cutworms caused some damage, forcing some producers to re-seed canola crops. Strong winds caused some damage to buildings and have eroded topsoil in many fields.

Farmers are busy seeding, fixing fences and controlling weeds.

West-Central Saskatchewan:

- Crop Districts 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A – Rosetown, Kindersley, Eston, Major
- Crop District 7B – Kerrobert, Macklin, Wilkie and Biggar areas

Seeding continues to rapidly advance in the region and producers now have 80 per cent of the crop in the ground. This is just slightly behind the five-year (2012-2016) seeding average of 83 per cent for this time of year. While some producers have wrapped up seeding, many more will need a couple more weeks of optimal field conditions.

Varying amounts of rainfall were reported this past week, ranging from trace amounts to 50 mm in the Phippen area. The Biggar area reported 25 mm of rain, the Tramping Lake area 31 mm, the Outlook area 9 mm, the Langham area 28 mm, the Harris area 14 mm and the Netherhill area 10 mm. The Battleford area has reported receiving the most precipitation (127 mm) in the region since April 1.

Cropland topsoil moisture is rated as seven per cent surplus, 89 per cent adequate and four per cent short. Hay land and pasture topsoil moisture is rated as five per cent surplus, 89 per cent adequate and six per cent short. Some areas could use a good rain to help crops emerge and pastures grow, while other areas need warm and dry weather.

Crops are slowly emerging and are in fair-to-excellent condition at this time. Some fields are being sprayed for cutworms and flea beetles, while others are expected to be re-seeded in the coming week. Strong winds and localized flooding have caused some damage as well as delaying herbicide applications. Last year's crop continues to be harvested, depending on field conditions, while unsalvageable fields are being burned.

Farmers are busy seeding, rolling pulses and controlling weeds.

Northeastern Saskatchewan:

- Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas

The northeastern region continues to struggle with excess moisture and frequent rainfall, which is delaying seeding. Just 43 per cent of the crop has been seeded, up from 25 per cent last week. The five-year (2012-2016) seeding average for this time of year is 80 per cent. Several weeks of warm and dry weather is needed to help dry fields up. Producers in some areas have not turned a wheel on their equipment, and crop reporters have indicated that there will likely be a large number of acres left unseeded due to excess moisture.

Large amounts of rain were reported in much of the region. The Nipawin area received 65 mm, the Prince Albert area 22 mm of rain, the Tisdale area 40 mm and the Arborfield area 17 mm. On the other hand, the Humboldt area received only 7 mm and the Bruno area 5 mm. The Nipawin area has reported receiving the most precipitation (200 mm) in the region since April 1.

Cropland topsoil moisture is rated as 46 per cent surplus and 54 per cent adequate. Hay land and pasture topsoil moisture is rated as 34 per cent surplus and 66 per cent adequate. Crop districts 8A and 9AE are reporting that 73 per cent and 28 per cent, respectively, of cropland has surplus topsoil moisture at this time. Warm, dry and windy weather is needed to help dry fields up.

Some crops are just starting to emerge and are in fair-to-excellent condition. Fields have been slow to dry, making it difficult to harvest what is left of last year's crop. Harvest operations continue, depending on field conditions, while unsalvageable fields are being burned. Hail, localized flooding and strong winds have caused some damage to fields and emerged crops. There are some reports of flea beetles and cutworms in early-seeded canola.

Farmers are busy seeding, working fields, controlling weeds and trying to finish last year's harvest.

Northwestern Saskatchewan:

- Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

Despite recent heavy rainfall in some areas, producers in the northwestern region now have 76 per cent of the crop seeded. The five-year (2012-2016) seeding average for this time of year is 85 per cent. Many fields remain wet, but others are in optimal condition. Several more weeks of warm and dry weather will be needed for producers to finish seeding operations.

Rainfall in the region ranged from 6 mm in the Duck Lake area to 52 mm in the Frenchman Butte area. The Neilburg area reported 29 mm, the Meadow Lake area 30 mm, the Hafford

area 35 mm, the North Battleford area 13 mm and the Debden area 9 mm. The St. Walburg area has reported receiving the most precipitation (160 mm) in the region since April 1.

Topsoil moisture conditions have improved with the recent rainfall. Cropland topsoil moisture is rated as 16 per cent surplus, 82 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as 12 per cent surplus, 86 per cent adequate and two per cent short. Crop District 9B is reporting that 24 per cent of cropland and 15 per cent of hay land and pasture have surplus topsoil moisture.

Crops are slowly emerging and are in fair-to-excellent condition. Localized flooding, frost, strong winds and insects such as cutworms and flea beetles have caused the majority of crop damage this past week. There are reports of farmers burning residue and unsalvageable crops.

Farmers are busy seeding and controlling weeds.



Crop Conditions - May 29, 2017

Provincial							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	25	7	16	10	15	17	6
% good	57	84	70	65	77	68	68
% fair	15	8	12	23	5	12	24
% poor	3	1	2	2	3	3	2
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	12	17	3	11	15	12	0
% good	56	60	37	75	71	69	63
% fair	26	17	51	14	13	17	34
% poor	4	5	6	0	1	1	3
% very poor	2	1	3	0	0	1	0
Southeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	31	17	16	13	16	22	13
% good	50	68	65	35	68	52	57
% fair	15	13	16	42	14	24	26
% poor	4	2	3	10	2	2	4
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	14	16	15	15	13	26	0
% good	42	42	0	69	67	45	25
% fair	33	26	75	16	19	28	75
% poor	8	10	10	0	1	1	0
% very poor	3	6	0	0	0	0	0
Southwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	3	0	13	8	22	8	15
% good	54	96	69	69	63	78	75
% fair	41	4	17	22	15	14	5
% poor	2	0	1	1	0	0	5
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	2	14	0	0	4	4	0
% good	59	48	32	100	77	79	61
% fair	38	26	56	0	17	14	35
% poor	1	8	7	0	2	2	4
% very poor	0	4	5	0	0	1	0
East-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	14	2	9	12	11	11	0
% good	78	86	68	69	74	63	100
% fair	6	12	18	17	7	17	0
% poor	2	0	5	2	8	9	0
% very poor	0	0	0	0	0	0	0

East-central (continued)							
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	10	13	N/A	1	19	12	0
% good	77	63	N/A	88	69	70	100
% fair	12	16	N/A	11	9	18	0
% poor	1	8	N/A	0	3	0	0
% very poor	0	0	N/A	0	0	0	0
West-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	14	19	13	15	13	11	0
% good	78	77	82	78	86	85	59
% fair	6	4	4	7	1	4	41
% poor	2	0	0	0	0	0	0
% very poor	0	0	1	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	3	14	3	8	25	17	0
% good	81	70	79	89	64	69	100
% fair	16	14	18	3	10	14	0
% poor	0	1	0	0	0	0	0
% very poor	0	1	0	0	1	0	0
Northeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	N/A	4	22	45	15	41	0
% good	N/A	87	66	30	83	56	100
% fair	N/A	6	12	25	2	3	0
% poor	N/A	3	0	0	0	0	0
% very poor	N/A	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	30	31	N/A	0	30	N/A	N/A
% good	70	45	N/A	50	53	N/A	N/A
% fair	0	24	N/A	50	17	N/A	N/A
% poor	0	0	N/A	0	0	N/A	N/A
% very poor	0	0	N/A	0	0	N/A	N/A
Northwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	82	38	29	N/A	22	31	N/A
% good	18	45	64	N/A	78	69	N/A
% fair	0	17	7	N/A	0	0	N/A
% poor	0	0	0	N/A	0	0	N/A
% very poor	0	0	0	N/A	0	0	N/A
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	50	27	N/A	N/A	26	17	N/A
% good	50	73	N/A	N/A	72	83	N/A
% fair	0	0	N/A	N/A	2	0	N/A
% poor	0	0	N/A	N/A	0	0	N/A
% very poor	0	0	N/A	N/A	0	0	N/A

Crop Districts and Rural Municipalities in Saskatchewan

- Regional Service Office
- Regional Satellite Office
- Crop District
- Rural Municipality



Projection: UTM Zone 13 Datum: NAD83




Data Source:
Crop Districts - Saskatchewan Ministry of Agriculture

Geomatic Services, Ministry of Agriculture June 10, 2014

Weekly Rainfall

from May 23 to May 29, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period May 23 to 29, 2017


Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	7	55	4A	49	White Valley	N/A	43	7A	287	St. Andrews	N/A	52
	3	Enniskillen	9	51		51	Reno	NIL	37.5		288	Pleasant Valley	3	55
	31	Storthoaks	N/A	10		79	Arlington	4	46		290 A	Kindersley	10	58
	32	Reciprocity	8	63		109	Carmichael	7	69		290 B	Kindersley	5	35
	33	Moose Creek	13	55		110	Piapot	N/A	43.5		290 C	Kindersley	N/A	24
	34	Browning	14	34		111	Maple Creek	N/A	N/A		292	Milton	10	61
	61	Antler	N/A	5	4B	139 A	Gull Lake	7	34		317 A	Marriott	N/A	N/A
	64	Brock	14	31		139 B	Gull Lake	8	22.5		317 B	Marriott	14	61
	65	Tecumseh	4	36		142	Enterprise	N/A	N/A		318	Mountain View	25	102
1B	91	Maryfield	N/A	26		169	Pittville	N/A	34		320 A	Oakdale	6	42.5
	94	Hazelwood	N/A	N/A		231	Happyland	5	52		320 B	Oakdale	16	52
	122	Martin	7	17	5A	183	Fertile Belt	11	33		321	Prairiedale	N/A	50
	123	Silverwood	15	28		211 A	Churchbridge	14	53	7B	347	Biggar	22	80
	124	Kingsley	12	42		211 B	Churchbridge	11	35		350 A	Mariposa	31	87.5
	125 A	Chester	4	32		211 C	Churchbridge	NIL	0		350 B	Mariposa	14	72
	125 B	Chester	NIL	33		213	Saltcoats	8	42		351	Progress	18	96
	151	Rocanville	14	31		218	Cupar	N/A	N/A		352	Heart's Hill	14	86
	154 A	Elcapo	4	23		241	Calder	16	30		377	Glenside	17	82
	154 B	Elcapo	7	22		243	Wallace	2	13		378	Rosemount	23	107
	155	Wolseley	1	24.3		244	Orkney	5	12		379	Reford	N/A	58
2A	67	Weyburn	2	21		245 A	Garry	3	34		381	Grass Lake	10	64.5
	68	Brokenshell	5	25		245 B	Garry	6	49		382	Eye Hill	17.5	75
	96	Fillmore	2.8	3.8		245 C	Garry	N/A	40.5		409 A	Buffalo	46	127
	97	Wellington	1	13		246 A	Ituna Bon Accord	5	39		409 B	Buffalo	50	101
2B	127 A	Francis	9	42		246 B	Ituna Bon Accord	15	46		410	Round Valley	12.5	101.2
	127 B	Francis	2.5	21.3		247	Kellross	7	49	8A	395	Porcupine	N/A	56
	129	Bratt's Lake	2	19		248	Touchwood	N/A	27.5		397	Barrier Valley	9.9	65.8
	131 A	Baildon	1	59		271	Cote	23	65		428	Star City	5	64
	131 B	Baildon	4	69		273	Sliding Hills	N/A	38		456	Arborfield	17	104
	156 A	Indian Head	3.3	28.8	5B	277	Emerald	16	75		457	Connaught	40	122
	156 B	Indian Head	NIL	55		305	Invermay	17	70		486	Moose Range	27	106
	159	Sherwood	2	26		307	Elfros	9	41		487	Nipawin	65	199.5
	160 A	Pense	NIL	14		308 A	Big Quill	10	27	8B	369	St. Peter	7	39
	160 B	Pense	N/A	18.5		308 B	Big Quill	18	37		370 A	Humboldt	3	53
	161	Moose Jaw	2	27		331	Livingston	16	74		370 B	Humboldt	N/A	5
	162	Caron	1	26.5		335	Hazel Dell	8	26		371	Bayne	5	59
	191	Marquis	N/A	32		336	Sasman	10	47		372	Grant	7.9	88.4
3ASE	38 A	Laurier	2	16.5		337	Lakeview	17.5	46.5		400	Three Lakes	N/A	35
	38 B	Laurier	NIL	16		338	Lakeside	15	49		429 A	Flett's Springs	N/A	43
	39	The Gap	NIL	10		366	Kelvington	20	57		429 B	Flett's Springs	13	77.5
3ASW	10	Happy Valley	N/A	46		367	Ponass Lake	15	53.5		459	Kinistino	3	55
	12	Poplar Valley	NIL	47		368	Spalding	8	47	9AE	460	Birch Hills	3	59.8
	40	Bengough	N/A	N/A	6A	190 A	Dufferin	6	39		488	Torch River	8	90
	42	Willow Bunch	NIL	34		190 B	Dufferin	5	46		491	Buckland	9.5	18.5
	43	Old Post	3	36		190 C	Dufferin	NIL	45		520	Paddockwood	22	122
	73 A	Stonehenge	NIL	41.4		190 D	Dufferin	N/A	39	9AW	521	Lakeland	22	122
	73 B	Stonehenge	3	59		219 A	Longlaketon	10	38		406	Mayfield	13	58
3AN	102	Lake Johnston	N/A	38.7		219 B	Longlaketon	5	53		435	Redberry	35	120
	103	Sutton	N/A	41		220	McKillop	8	42		436	Douglas	10	97
	132 A	Hillsborough	1	24		221 A	Sarnia	8.1	51.0		463	Duck Lake	6	83.6
	132 B	Hillsborough	8.5	45.5		221 B	Sarnia	5	35.6		466	Meeting Lake	9	121
	193	Eyebrow	2	25		222	Craik	4	40		467 A	Round Hill	13	111
	17	Val Marie	N/A	10.1		251	Big Arm	7.5	29.5		467 B	Round Hill	9	85
3BS	18	Lone Tree	NIL	27		252	Arm River	5	33		467 C	Round Hill	9	114
	75	Pinto Creek	7	48		279	Mount Hope	12	32.2		494	Canwood	9	97
	76	Auvergne	N/A	36		282	McCraney	4	51	9B	438	Battle River	26	112
	77	Wise Creek	N/A	46		312	Morris	9	52.5		440	Hillsdale	22	89.5
	78	Grassy Creek	1.5	41.0		313	Lost River	NIL	48		442	Manitou Lake	29.2	108.4
	105	Glenbain	NIL	38		339	Leroy	4.4	33.8		498 A	Parkdale	29	126
	106	Whiska Creek	3	44		340	Wolverine	2	57		498 B	Parkdale	25	81.5
	107	Lac Pelletier	N/A	34		341	Viscount	8	55		499 A	Mervin	27	115
	108	Bone Creek	4	47		343 A	Blucher	NIL	35		501 A	Frenchman Butte	35	160
	138 A	Webb	1	36		223 A	Huron	3	37		501 B	Frenchman Butte	31	95
3BN	138 B	Webb	2	23	6B	223 B	Huron	NIL	32		501 C	Frenchman Butte	52	155
	165	Morse	N/A	N/A		284 A	Rudy	9	80		502	Britannia	38	157
	166	Excelsior	4	45		284 B	Rudy	12	56.5		561	Loon Lake	18	140
	168 A	Riverside	NIL	25		284 C	Rudy	6	41		588 A	Meadow Lake	21	134
	168 B	Riverside	N/A	22.2		285	Fertile Valley	8	53		588 B	Meadow Lake	30	153
	226	Victory	N/A	27		286	Milden	6	61		622	Beaver River	N/A	177.7
	228 A	Lacadena	7	66		314	Dundurn	4	57					
	228 B	Lacadena	N/A	6.5		344 A	Corman Park	28	77					
	257	Monet	NIL	45.5		344 B	Corman Park	N/A	0					
						346	Perdue	9	57					
						376	Eagle Creek	16	87					
						403	Rosthern	4	85					

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Municipality No: A, B, C and D - more than one reporter

Cumulative Rainfall


from April 1 to May 29, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Cropland Topsoil Moisture Conditions

May 29, 2017




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Hay and Pasture Topsoil Moisture Conditions

May 29, 2017



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.